

C H A P T E R 2

Differentiation

Section 2.1	The Derivative and the Tangent Line Problem	95
Section 2.2	Basic Differentiation Rules and Rates of Change	109
Section 2.3	Product and Quotient Rules and Higher-Order Derivatives	120
Section 2.4	The Chain Rule	134
Section 2.5	Implicit Differentiation	147
Section 2.6	Related Rates	161
Review Exercises	173
Problem Solving	184

CHAPTER 2

Differentiation


Section 2.1 The Derivative and the Tangent Line Problem

1. (a) At (x_1, y_1) , slope = 0.

At (x_2, y_2) , slope $\approx \frac{5}{2}$.

(b) At (x_1, y_1) , slope $\approx -\frac{5}{2}$.

At (x_2, y_2) , slope ≈ 2 .



$$(c) y = \frac{f(4) - f(1)}{4 - 1}(x - 1) + f(1)$$

$$= \frac{3}{3}(x - 1) + 2$$

$$= 1(x - 1) + 2$$

$$= x + 1$$

5. $f(x) = 3 - 2x$ is a line. Slope = -2

$$\begin{aligned} 7. \text{ Slope at } (1, -3) &= \lim_{\Delta x \rightarrow 0} \frac{g(1 + \Delta x) - g(1)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(1 + \Delta x)^2 - 4 - (-3)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{1 + 2(\Delta x) + (\Delta x)^2 - 1}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} [2 + (\Delta x)] = 2 \end{aligned}$$

$$\begin{aligned} 9. \text{ Slope at } (0, 0) &= \lim_{\Delta t \rightarrow 0} \frac{f(0 + \Delta t) - f(0)}{\Delta t} \\ &= \lim_{\Delta t \rightarrow 0} \frac{3(\Delta t) - (\Delta t)^2 - 0}{\Delta t} \\ &= \lim_{\Delta t \rightarrow 0} (3 - \Delta t) = 3 \end{aligned}$$

2. (a) At (x_1, y_1) , slope $\approx \frac{2}{3}$.

At (x_2, y_2) , slope $\approx -\frac{2}{5}$.

(b) At (x_1, y_1) , slope $\approx \frac{4}{3}$.

At (x_2, y_2) , slope $\approx \frac{5}{4}$.

$$4. (a) \frac{f(4) - f(1)}{4 - 1} = \frac{5 - 2}{3} = 1$$

$$\frac{f(4) - f(3)}{4 - 3} \approx \frac{5 - 4.75}{1} = 0.25$$

$$\text{Thus, } \frac{f(4) - f(1)}{4 - 1} > \frac{f(4) - f(3)}{4 - 3}.$$

(b) The slope of the tangent line at $(1, 2)$ equals $f'(1)$. This slope is steeper than the slope of the line through $(1, 2)$ and $(4, 5)$. Thus,

$$\frac{f(4) - f(1)}{4 - 1} < f'(1).$$

6. $g(x) = \frac{3}{2}x + 1$ is a line. Slope = $\frac{3}{2}$

$$\begin{aligned} 8. \text{ Slope at } (2, 1) &= \lim_{\Delta x \rightarrow 0} \frac{g(2 + \Delta x) - g(2)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{5 - (2 + \Delta x)^2 - 1}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{5 - 4 - 4(\Delta x) - (\Delta x)^2 - 1}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (-4 - \Delta x) = -4 \end{aligned}$$

$$\begin{aligned} 10. \text{ Slope at } (-2, 7) &= \lim_{\Delta t \rightarrow 0} \frac{h(-2 + \Delta t) - h(-2)}{\Delta t} \\ &= \lim_{\Delta t \rightarrow 0} \frac{(-2 + \Delta t)^2 + 3 - 7}{\Delta t} \\ &= \lim_{\Delta t \rightarrow 0} \frac{4 - 4(\Delta t) + (\Delta t)^2 - 4}{\Delta t} \\ &= \lim_{\Delta t \rightarrow 0} (-4 + \Delta t) = -4 \end{aligned}$$

11. $f(x) = 3$

$$\begin{aligned}f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{3 - 3}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} 0 = 0\end{aligned}$$

14. $f(x) = 3x + 2$

$$\begin{aligned}f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{[3(x + \Delta x) + 2] - [3x + 2]}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{3\Delta x}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} 3 = 3\end{aligned}$$

16. $f(x) = 9 - \frac{1}{2}x$

$$\begin{aligned}f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{[9 - (1/2)(x + \Delta x)] - [9 - (1/2)x]}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \left(-\frac{1}{2} \right) = -\frac{1}{2}\end{aligned}$$

17. $f(x) = 2x^2 + x - 1$

$$\begin{aligned}f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{[2(x + \Delta x)^2 + (x + \Delta x) - 1] - [2x^2 + x - 1]}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{(2x^2 + 4x\Delta x + 2(\Delta x)^2 + x + \Delta x - 1) - (2x^2 + x - 1)}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{4x\Delta x + 2(\Delta x)^2 + \Delta x}{\Delta x} = \lim_{\Delta x \rightarrow 0} (4x + 2\Delta x + 1) = 4x + 1\end{aligned}$$

18. $f(x) = 1 - x^2$

$$\begin{aligned}f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{[1 - (x + \Delta x)^2] - [1 - x^2]}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{1 - x^2 - 2x\Delta x - (\Delta x)^2 - 1 + x^2}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{-2x\Delta x - (\Delta x)^2}{\Delta x} = \lim_{\Delta x \rightarrow 0} (-2x - \Delta x) = -2x\end{aligned}$$

12. $g(x) = -5$

$$\begin{aligned}g'(x) &= \lim_{\Delta x \rightarrow 0} \frac{g(x + \Delta x) - g(x)}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{-5 - (-5)}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{0}{\Delta x} = 0\end{aligned}$$

13. $f(x) = -5x$

$$\begin{aligned}f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{-5(x + \Delta x) - (-5x)}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} -5 = -5\end{aligned}$$

15. $h(s) = 3 + \frac{2}{3}s$

$$\begin{aligned}h'(s) &= \lim_{\Delta s \rightarrow 0} \frac{h(s + \Delta s) - h(s)}{\Delta s} \\&= \lim_{\Delta s \rightarrow 0} \frac{3 + \frac{2}{3}(s + \Delta s) - (3 + \frac{2}{3}s)}{\Delta s} \\&= \lim_{\Delta s \rightarrow 0} \frac{\frac{2}{3}\Delta s}{\Delta s} = \frac{2}{3}\end{aligned}$$

19. $f(x) = x^3 - 12x$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{[(x + \Delta x)^3 - 12(x + \Delta x)] - [x^3 - 12x]}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^3 + 3x^2 \Delta x + 3x(\Delta x)^2 + (\Delta x)^3 - 12x - 12\Delta x - x^3 + 12x}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{3x^2 \Delta x + 3x(\Delta x)^2 + (\Delta x)^3 - 12\Delta x}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (3x^2 + 3x\Delta x + (\Delta x)^2 - 12) = 3x^2 - 12 \end{aligned}$$

20. $f(x) = x^3 + x^2$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{[(x + \Delta x)^3 + (x + \Delta x)^2] - [x^3 + x^2]}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^3 + 3x^2 \Delta x + 3x(\Delta x)^2 + (\Delta x)^3 + x^2 + 2x\Delta x + (\Delta x)^2 - x^3 - x^2}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{3x^2 \Delta x + 3x(\Delta x)^2 + (\Delta x)^3 + 2x\Delta x + (\Delta x)^2}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (3x^2 + 3x\Delta x + (\Delta x)^2 + 2x + (\Delta x)) = 3x^2 + 2x \end{aligned}$$

21. $f(x) = \frac{1}{x-1}$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\frac{1}{x + \Delta x - 1} - \frac{1}{x-1}}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x-1) - (x + \Delta x - 1)}{\Delta x(x + \Delta x - 1)(x-1)} \\ &= \lim_{\Delta x \rightarrow 0} \frac{-\Delta x}{\Delta x(x + \Delta x - 1)(x-1)} \\ &= \lim_{\Delta x \rightarrow 0} \frac{-1}{(x + \Delta x - 1)(x-1)} \\ &= -\frac{1}{(x-1)^2} \end{aligned}$$

22. $f(x) = \frac{1}{x^2}$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\frac{1}{(x + \Delta x)^2} - \frac{1}{x^2}}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^2 - (x + \Delta x)^2}{\Delta x(x + \Delta x)^2 x^2} \\ &= \lim_{\Delta x \rightarrow 0} \frac{-2x\Delta x - (\Delta x)^2}{\Delta x(x + \Delta x)^2 x^2} \\ &= \lim_{\Delta x \rightarrow 0} \frac{-2x - \Delta x}{(x + \Delta x)^2 x^2} \\ &= \frac{-2x}{x^4} \\ &= -\frac{2}{x^3} \end{aligned}$$

23. $f(x) = \sqrt{x+1}$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\sqrt{x + \Delta x + 1} - \sqrt{x+1}}{\Delta x} \cdot \left(\frac{\sqrt{x + \Delta x + 1} + \sqrt{x+1}}{\sqrt{x + \Delta x + 1} + \sqrt{x+1}} \right) \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x + 1) - (x+1)}{\Delta x [\sqrt{x + \Delta x + 1} + \sqrt{x+1}]} \\ &= \lim_{\Delta x \rightarrow 0} \frac{1}{\sqrt{x + \Delta x + 1} + \sqrt{x+1}} = \frac{1}{\sqrt{x+1} + \sqrt{x+1}} = \frac{1}{2\sqrt{x+1}} \end{aligned}$$

24. $f(x) = \frac{4}{\sqrt{x}}$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\frac{4}{\sqrt{x + \Delta x}} - \frac{4}{\sqrt{x}}}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{4\sqrt{x} - 4\sqrt{x + \Delta x}}{\Delta x \sqrt{x} \sqrt{x + \Delta x}} \cdot \left(\frac{\sqrt{x} + \sqrt{x + \Delta x}}{\sqrt{x} + \sqrt{x + \Delta x}} \right) \\ &= \lim_{\Delta x \rightarrow 0} \frac{4x - 4(x + \Delta x)}{\Delta x \sqrt{x} \sqrt{x + \Delta x} (\sqrt{x} + \sqrt{x + \Delta x})} \\ &= \lim_{\Delta x \rightarrow 0} \frac{-4}{\sqrt{x} \sqrt{x + \Delta x} (\sqrt{x} + \sqrt{x + \Delta x})} \\ &= \frac{-4}{\sqrt{x} \sqrt{x} (\sqrt{x} + \sqrt{x})} = \frac{-2}{x \sqrt{x}} \end{aligned}$$

25. (a) $f(x) = x^2 + 1$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{[(x + \Delta x)^2 + 1] - [x^2 + 1]}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{2x \Delta x + (\Delta x)^2}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (2x + \Delta x) = 2x \end{aligned}$$


At $(2, 5)$, the slope of the tangent line is $m = 2(2) = 4$.
The equation of the tangent line is

$$y - 5 = 4(x - 2)$$

$$y - 5 = 4x - 8$$

$$y = 4x - 3.$$

(b)



26. (a) $f(x) = x^2 + 2x + 1$


$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{[(x + \Delta x)^2 + 2(x + \Delta x) + 1] - [x^2 + 2x + 1]}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{2x \Delta x + (\Delta x)^2 + 2 \Delta x}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (2x + \Delta x + 2) = 2x + 2 \end{aligned}$$

At $(-3, 4)$, the slope of the tangent line is $m = 2(-3) + 2 = -4$.
The equation of the tangent line is

$$y - 4 = -4(x + 3)$$

$$y = -4x - 8.$$

(b)



27. (a) $f(x) = x^3$


$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^3 - x^3}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{3x^2 \Delta x + 3x(\Delta x)^2 + (\Delta x)^3}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (3x^2 + 3x \Delta x + (\Delta x)^2) = 3x^2 \end{aligned}$$

At $(2, 8)$, the slope of the tangent is $m = 3(2)^2 = 12$.
The equation of the tangent line is

$$y - 8 = 12(x - 2)$$

$$y = 12x - 16.$$

(b)




28. (a) $f(x) = x^3 + 1$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{[(x + \Delta x)^3 + 1] - (x^3 + 1)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^3 + 3x^2(\Delta x) + 3x(\Delta x)^2 + (\Delta x)^3 + 1 - x^3 - 1}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} [3x^2 + 3x(\Delta x) + (\Delta x)^2] = 3x^2 \end{aligned}$$

At $(1, 2)$, the slope of the tangent line is $m = 3(1)^2 = 3$.
The equation of the tangent line is

$$y - 2 = 3(x - 1)$$

$$y = 3x - 1.$$



29. (a) $f(x) = \sqrt{x}$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\sqrt{x + \Delta x} - \sqrt{x}}{\Delta x} \cdot \frac{\sqrt{x + \Delta x} + \sqrt{x}}{\sqrt{x + \Delta x} + \sqrt{x}} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x) - x}{\Delta x(\sqrt{x + \Delta x} + \sqrt{x})} \\ &= \lim_{\Delta x \rightarrow 0} \frac{1}{\sqrt{x + \Delta x} + \sqrt{x}} = \frac{1}{2\sqrt{x}} \end{aligned}$$


At $(1, 1)$, the slope of the tangent line is

$$m = \frac{1}{2\sqrt{1}} = \frac{1}{2}.$$

The equation of the tangent line is

$$y - 1 = \frac{1}{2}(x - 1)$$

$$y = \frac{1}{2}x + \frac{1}{2}.$$



30. (a) $f(x) = \sqrt{x - 1}$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\sqrt{x + \Delta x - 1} - \sqrt{x - 1}}{\Delta x} \cdot \left(\frac{\sqrt{x + \Delta x - 1} + \sqrt{x - 1}}{\sqrt{x + \Delta x - 1} + \sqrt{x - 1}} \right) \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x - 1) - (x - 1)}{\Delta x(\sqrt{x + \Delta x - 1} + \sqrt{x - 1})} \\ &= \lim_{\Delta x \rightarrow 0} \frac{1}{\sqrt{x + \Delta x - 1} + \sqrt{x - 1}} = \frac{1}{2\sqrt{x - 1}} \end{aligned}$$


At $(5, 2)$, the slope of the tangent line is

$$m = \frac{1}{2\sqrt{5 - 1}} = \frac{1}{4}.$$

The equation of the tangent line is

$$y - 2 = \frac{1}{4}(x - 5)$$

$$y = \frac{1}{4}x + \frac{3}{4}$$



31. (a) $f(x) = x + \frac{4}{x}$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x) + \frac{4}{x + \Delta x} - \left(x + \frac{4}{x}\right)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x(x + \Delta x)(x + \Delta x) + 4x - x^2(x + \Delta x) - 4(x + \Delta x)}{x(\Delta x)(x + \Delta x)} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^3 + 2x^2(\Delta x) + x(\Delta x)^2 - x^3 - x^2(\Delta x) - 4(\Delta x)}{x(\Delta x)(x + \Delta x)} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^2(\Delta x) + x(\Delta x)^2 - 4(\Delta x)}{x(\Delta x)(x + \Delta x)} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^2 + x(\Delta x) - 4}{x(x + \Delta x)} \\ &= \frac{x^2 - 4}{x^2} = 1 - \frac{4}{x^2} \end{aligned}$$

At $(4, 5)$, the slope of the tangent line is

$$m = 1 - \frac{4}{16} = \frac{3}{4}.$$

The equation of the tangent line is

$$y - 5 = \frac{3}{4}(x - 4)$$

$$y = \frac{3}{4}x + 2.$$

32. (a) $f(x) = \frac{1}{x+1}$



$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\frac{1}{x + \Delta x + 1} - \frac{1}{x + 1}}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + 1) - (x + \Delta x + 1)}{\Delta x(x + \Delta x + 1)(x + 1)} \\ &= \lim_{\Delta x \rightarrow 0} -\frac{1}{(x + \Delta x + 1)(x + 1)} \\ &= -\frac{1}{(x + 1)^2} \end{aligned}$$

At $(0, 1)$, the slope of the tangent line is

$$m = -\frac{1}{(0 + 1)^2} = -1.$$

The equation of the tangent line is $y = -x + 1$.

(b)



33. From Exercise 27 we know that $f'(x) = 3x^2$. Since the slope of the given line is 3, we have

$$3x^2 = 3$$

$$x = \pm 1.$$

Therefore, at the points $(1, 1)$ and $(-1, -1)$ the tangent lines are parallel to $3x - y + 1 = 0$. These lines have equations

$$y - 1 = 3(x - 1) \quad \text{and} \quad y + 1 = 3(x + 1)$$

$$y = 3x - 2$$

$$y = 3x + 2.$$

34. Using the limit definition of derivative, $f'(x) = 3x^2$. Since the slope of the given line is 3, we have

$$3x^2 = 3$$

$$x^2 = 1 \Rightarrow x = \pm 1.$$

Therefore, at the points $(1, 3)$ and $(-1, 1)$ the tangent lines are parallel to $3x - y - 4 = 0$. These lines have equations

$$y - 3 = 3(x - 1) \quad \text{and} \quad y - 1 = 3(x + 1)$$

$$y = 3x$$

$$y = 3x + 4.$$

35. Using the limit definition of derivative,

$$f'(x) = \frac{-1}{2x\sqrt{x}}.$$

Since the slope of the given line is $-\frac{1}{2}$, we have

$$-\frac{1}{2x\sqrt{x}} = -\frac{1}{2}$$

$$x = 1.$$

Therefore, at the point $(1, 1)$ the tangent line is parallel to $x + 2y - 6 = 0$. The equation of this line is

$$y - 1 = -\frac{1}{2}(x - 1)$$

$$y - 1 = -\frac{1}{2}x + \frac{1}{2}$$

$$y = -\frac{1}{2}x + \frac{3}{2}$$

37. $f(x) = x \Rightarrow f'(x) = 1$ Matches (b).


39. $f(x) = \sqrt{x} \Rightarrow f'(x)$ Matches (a).

(decreasing slope as $x \rightarrow \infty$)


41. $g(5) = 2$ because the tangent line passes through $(5, 2)$.

$$g'(5) = \frac{2 - 0}{5 - 9} = \frac{2}{-4} = -\frac{1}{2}$$

43. The slope of the graph of f is 1 $\Rightarrow f'(x) = 1$.




44. The slope of the graph of f is 0 $\Rightarrow f'(x) = 0$.




42. $h(-1) = 4$ because the tangent line passes through $(-1, 4)$.

$$h'(-1) = \frac{6 - 4}{3 - (-1)} = \frac{2}{4} = \frac{1}{2}$$

45. The slope of the graph of f is negative for $x < 4$, positive for $x > 4$, and 0 at $x = 4$.




46. The slope of the graph of f is -1 for $x < 4$, 1 for $x > 4$, and undefined at $x = 4$.



49. $f(x) = 5 - 3x$ and $c = 1$

53. $f(0) = 2$ and $f'(x) = -3$,
 $-\infty < x < \infty$


$f(x) = -3x + 2$



50. $f(x) = x^3$ and $c = -2$

54. $f(0) = 4$, $f'(0) = 0$; $f'(x) < 0$ for
 $x < 0$, $f'(x) > 0$ for $x > 0$

$f(x) = x^2 + 4$




51. $f(x) = -x^2$ and $c = 6$

52. $f(x) = 2\sqrt{x}$ and $c = 9$

55. $f(0) = 0$; $f'(0) = 0$; $f'(x) > 0$ if
 $x \neq 0$

$f(x) = x^3$



56. (a) If $f'(c) = 3$ and f is odd, then $f'(-c) = f'(c) = 3$.

- (b) If $f'(c) = 3$ and f is even, then $f'(-c) = -f'(c) = -3$.

57. Let (x_0, y_0) be a point of tangency on the graph of f . By the limit definition for the derivative, $f'(x) = 4 - 2x$. The slope of the line through $(2, 5)$ and (x_0, y_0) equals the derivative of f at x_0 :

$$\frac{5 - y_0}{2 - x_0} = 4 - 2x_0$$

$$5 - y_0 = (2 - x_0)(4 - 2x_0)$$

$$5 - (4x_0 - x_0^2) = 8 - 8x_0 + 2x_0^2$$

$$0 = x_0^2 - 4x_0 + 3$$


$$0 = (x_0 - 1)(x_0 - 3) \Rightarrow x_0 = 1, 3$$

Therefore, the points of tangency are $(1, 3)$ and $(3, 3)$, and the corresponding slopes are 2 and -2 . The equations of the tangent lines are:

$$y - 5 = 2(x - 2) \quad y - 5 = -2(x - 2)$$

$$y = 2x + 1$$

$$y = -2x + 9$$



58. Let (x_0, y_0) be a point of tangency on the graph of f . By the limit definition for the derivative, $f'(x) = 2x$. The slope of the line through $(1, -3)$ and (x_0, y_0) equals the derivative of f at x_0 :

$$\frac{-3 - y_0}{1 - x_0} = 2x_0$$

$$-3 - y_0 = (1 - x_0)2x_0$$

$$-3 - x_0^2 = 2x_0 - 2x_0^2$$

$$x_0^2 - 2x_0 - 3 = 0$$


$$(x_0 - 3)(x_0 + 1) = 0 \Rightarrow x_0 = 3, -1$$

Therefore, the points of tangency are $(3, 9)$ and $(-1, 1)$, and the corresponding slopes are 6 and -2 . The equations of the tangent lines are:

$$y + 3 = 6(x - 1) \quad y + 3 = -2(x - 1)$$

$$y = 6x - 9$$

$$y = -2x - 1$$



59. (a) $g'(0) = -3$

(b) $g'(3) = 0$

(c) Because $g'(1) = -\frac{8}{3}$, g is decreasing (falling) at $x = 1$.

(d) Because $g'(-4) = \frac{7}{3}$, g is increasing (rising) at $x = -4$.

(e) Because $g'(4)$ and $g'(6)$ are both positive, $g(6)$ is greater than $g(4)$, and $g(6) - g(4) > 0$.

(f) No, it is not possible. All you can say is that g is decreasing (falling) at $x = 2$.

60. (a) $f(x) = x^2$


$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^2 - x^2}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^2 + 2x(\Delta x) + (\Delta x)^2 - x^2}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\Delta x(2x + \Delta x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (2x + \Delta x) = 2x \end{aligned}$$

At $x = -1$, $f'(-1) = -2$ and the tangent line is

$$y - 1 = -2(x + 1) \quad \text{or} \quad y = -2x - 1.$$

At $x = 0$, $f'(0) = 0$ and the tangent line is $y = 0$.

At $x = 1$, $f'(1) = 2$ and the tangent line is $y = 2x - 1$.




For this function, the slopes of the tangent lines are always distinct for different values of x .

61. $f(x) = \frac{1}{4}x^3$

By the limit definition of the derivative we have $f'(x) = \frac{3}{4}x^2$.

x	-2	-1.5	-1	-0.5	0	0.5	1	1.5	2
$f(x)$	-2	$-\frac{27}{32}$	$-\frac{1}{4}$	$-\frac{1}{32}$	0	$\frac{1}{32}$	$\frac{1}{4}$	$\frac{27}{32}$	2
$f'(x)$	3	$\frac{27}{16}$	$\frac{3}{4}$	$\frac{3}{16}$	0	$\frac{3}{16}$	$\frac{3}{4}$	$\frac{27}{16}$	3




For this function, the slopes of the tangent lines are sometimes the same.


62. $f(x) = \frac{1}{2}x^2$

By the limit definition of the derivative we have $f'(x) = x$.

x	-2	-1.5	-1	-0.5	0	0.5	1	1.5	2
$f(x)$	2	1.125	0.5	0.125	0	0.125	0.5	1.125	2
$f'(x)$	-2	-1.5	-1	-0.5	0	0.5	1	1.5	2




63. $g(x) = \frac{f(x+0.01) - f(x)}{0.01}$
 $= [2(x+0.01) - (x+0.01)^2 - 2x + x^2]100$
 $= 2 - 2x - 0.01$



The graph of $g(x)$ is approximately the graph of $f'(x) = 2 - 2x$.

64. $g(x) = \frac{f(x+0.01) - f(x)}{0.01}$
 $= (3\sqrt{x+0.01} - 3\sqrt{x})100$




The graph of $g(x)$ is approximately the graph of $f'(x) = \frac{3}{2\sqrt{x}}$.

65. $f(2) = 2(4-2) = 4$, $f(2.1) = 2.1(4-2.1) = 3.99$
 $f'(2) \approx \frac{3.99-4}{2.1-2} = -0.1$ [Exact: $f'(2) = 0$]


66. $f(2) = \frac{1}{4}(2^3) = 2$, $f(2.1) = 2.31525$
 $f'(2) \approx \frac{2.31525-2}{2.1-2} = 3.1525$ [Exact: $f'(2) = 3$]

67. $f(x) = \frac{1}{\sqrt{x}}$ and $f'(x) = \frac{-1}{2x^{3/2}}$.



As $x \rightarrow \infty$, f is nearly horizontal and thus $f' \approx 0$.

68. $f(x) = \frac{x^3}{4} - 3x$ and $f'(x) = \frac{3}{4}x^2 - 3$




69. $f(x) = 4 - (x-3)^2$

$$\begin{aligned} S_{\Delta x}(x) &= \frac{f(2+\Delta x) - f(2)}{\Delta x}(x-2) + f(2) \\ &= \frac{4 - (2+\Delta x-3)^2 - 3}{\Delta x}(x-2) + 3 = \frac{1 - (\Delta x-1)^2}{\Delta x}(x-2) + 3 = (-\Delta x+2)(x-2) + 3 \end{aligned}$$

(a) $\Delta x = 1$: $S_{\Delta x} = (x-2) + 3 = x + 1$

$\Delta x = 0.5$: $S_{\Delta x} = \left(\frac{3}{2}\right)(x-2) + 3 = \frac{3}{2}x$

$\Delta x = 0.1$: $S_{\Delta x} = \left(\frac{19}{10}\right)(x-2) + 3 = \frac{19}{10}x - \frac{4}{5}$



(b) As $\Delta x \rightarrow 0$, the line approaches the tangent line to f at $(2, 3)$.


70. $f(x) = x + \frac{1}{x}$

$$\begin{aligned} S_{\Delta x}(x) &= \frac{f(2+\Delta x) - f(2)}{\Delta x}(x-2) + f(2) = \frac{(2+\Delta x) + \frac{1}{2+\Delta x} - \frac{5}{2}}{\Delta x}(x-2) + \frac{5}{2} \\ &= \frac{2(2+\Delta x)^2 + 2 - 5(2+\Delta x)}{2(2+\Delta x)\Delta x}(x-2) + \frac{5}{2} = \frac{(2\Delta x+3)}{2(2+\Delta x)}(x-2) + \frac{5}{2} \end{aligned}$$

(a) $\Delta x = 1$: $S_{\Delta x} = \frac{5}{6}(x-2) + \frac{5}{2} = \frac{5}{6}x + \frac{5}{6}$

$\Delta x = 0.5$: $S_{\Delta x} = \frac{4}{5}(x-2) + \frac{5}{2} = \frac{4}{5}x + \frac{9}{10}$

$\Delta x = 0.1$: $S_{\Delta x} = \frac{16}{21}(x-2) + \frac{5}{2} = \frac{16}{21}x + \frac{41}{42}$



(b) As $\Delta x \rightarrow 0$, the line approaches the tangent line to f at $(2, \frac{5}{2})$.

71. $f(x) = x^2 - 1, c = 2$

$$f'(2) = \lim_{x \rightarrow 2} \frac{f(x) - f(2)}{x - 2} = \lim_{x \rightarrow 2} \frac{(x^2 - 1) - 3}{x - 2} = \lim_{x \rightarrow 2} \frac{(x - 2)(x + 2)}{x - 2} = \lim_{x \rightarrow 2} (x + 2) = 4$$

72. $g(x) = x(x - 1) = x^2 - x, c = 1$

$$g'(1) = \lim_{x \rightarrow 1} \frac{g(x) - g(1)}{x - 1} = \lim_{x \rightarrow 1} \frac{x^2 - x - 0}{x - 1} = \lim_{x \rightarrow 1} \frac{x(x - 1)}{x - 1} = \lim_{x \rightarrow 1} x = 1$$

73. $f(x) = x^3 + 2x^2 + 1, c = -2$

$$f'(-2) = \lim_{x \rightarrow -2} \frac{f(x) - f(-2)}{x + 2} = \lim_{x \rightarrow -2} \frac{(x^3 + 2x^2 + 1) - 1}{x + 2} = \lim_{x \rightarrow -2} \frac{x^2(x + 2)}{x + 2} = \lim_{x \rightarrow -2} x^2 = 4$$

74. $f(x) = x^3 + 2x, c = 1$

$$f'(1) = \lim_{x \rightarrow 1} \frac{f(x) - f(1)}{x - 1} = \lim_{x \rightarrow 1} \frac{x^3 + 2x - 3}{x - 1} = \lim_{x \rightarrow 1} \frac{(x - 1)(x^2 + x + 3)}{x - 1} = \lim_{x \rightarrow 1} (x^2 + x + 3) = 5$$

75. $g(x) = \sqrt{|x|}, c = 0$

$$g'(0) = \lim_{x \rightarrow 0} \frac{g(x) - g(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{\sqrt{|x|}}{x}. \text{ Does not exist.}$$

$$\text{As } x \rightarrow 0^-, \frac{\sqrt{|x|}}{x} = \frac{-1}{\sqrt{|x|}} \rightarrow -\infty.$$

$$\text{As } x \rightarrow 0^+, \frac{\sqrt{|x|}}{x} = \frac{1}{\sqrt{x}} \rightarrow \infty.$$

76. $f(x) = \frac{1}{x}, c = 3$

$$f'(3) = \lim_{x \rightarrow 3} \frac{f(x) - f(3)}{x - 3} = \lim_{x \rightarrow 3} \frac{(1/x) - (1/3)}{x - 3} = \lim_{x \rightarrow 3} \frac{3 - x}{3x} \cdot \frac{1}{x - 3} = \lim_{x \rightarrow 3} \left(-\frac{1}{3x} \right) = -\frac{1}{9}$$

77. $f(x) = (x - 6)^{2/3}, c = 6$

$$f'(6) = \lim_{x \rightarrow 6} \frac{f(x) - f(6)}{x - 6} = \lim_{x \rightarrow 6} \frac{(x - 6)^{2/3} - 0}{x - 6} = \lim_{x \rightarrow 6} \frac{1}{(x - 6)^{1/3}}$$

Does not exist.

78. $g(x) = (x + 3)^{1/3}, c = -3$

$$g'(-3) = \lim_{x \rightarrow -3} \frac{g(x) - g(-3)}{x - (-3)} = \lim_{x \rightarrow -3} \frac{(x + 3)^{1/3} - 0}{x + 3} = \lim_{x \rightarrow -3} \frac{1}{(x + 3)^{2/3}}$$

Does not exist.

79. $h(x) = |x + 5|, c = -5$

$$h'(-5) = \lim_{x \rightarrow -5} \frac{h(x) - h(-5)}{x - (-5)} = \lim_{x \rightarrow -5} \frac{|x + 5| - 0}{x + 5} = \lim_{x \rightarrow -5} \frac{|x + 5|}{x + 5}$$

Does not exist.

80. $f(x) = |x - 4|, c = 4$

$$f'(4) = \lim_{x \rightarrow 4} \frac{f(x) - f(4)}{x - 4} = \lim_{x \rightarrow 4} \frac{|x - 4| - 0}{x - 4} = \lim_{x \rightarrow 4} \frac{|x - 4|}{x - 4}$$

Does not exist.


81. $f(x)$ is differentiable everywhere except at $x = -1$.
(Discontinuity)

82. $f(x)$ is differentiable everywhere except at $x = \pm 3$.
(Sharp turns in the graph)


84. $f(x)$ is differentiable everywhere except at $x = \pm 2$.
(Discontinuities)

86. $f(x)$ is differentiable everywhere except at $x = 0$. (Discontinuity)


87. $f(x) = |x + 3|$ is differentiable for all $x \neq -3$. There is a sharp corner at $x = -3$.




88. $f(x) = \frac{2x}{x-1}$ is differentiable for all $x \neq 1$. f is not defined at $x = 1$. (Vertical asymptote)



89. $f(x) = x^{2/5}$ is differentiable for all $x \neq 0$. There is a sharp corner at $x = 0$.



90. f is differentiable for all $x \neq 1$.
 f is not continuous at $x = 1$.



91. $f(x) = |x - 1|$

The derivative from the left is

$$\lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{x - 1} = \lim_{x \rightarrow 1^-} \frac{|x - 1| - 0}{x - 1} = \lim_{x \rightarrow 1^-} \frac{|x - 1|}{x - 1} = -1.$$

The derivative from the right is

$$\lim_{x \rightarrow 1^+} \frac{f(x) - f(1)}{x - 1} = \lim_{x \rightarrow 1^+} \frac{|x - 1| - 0}{x - 1} = \lim_{x \rightarrow 1^+} \frac{|x - 1|}{x - 1} = 1.$$

The one-sided limits are not equal. Therefore, f is not differentiable at $x = 1$.

92. $f(x) = \sqrt{1 - x^2}$

The derivative from the left does not exist because

$$\lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{x - 1} = \lim_{x \rightarrow 1^-} \frac{\sqrt{1 - x^2} - 0}{x - 1} = \lim_{x \rightarrow 1^-} \frac{\sqrt{1 - x^2}}{x - 1} \cdot \frac{\sqrt{1 - x^2}}{\sqrt{1 - x^2}} = \lim_{x \rightarrow 1^-} -\frac{1 + x}{\sqrt{1 - x^2}} = -\infty. \quad (\text{Vertical tangent})$$

The limit from the right does not exist since f is undefined for $x > 1$. Therefore, f is not differentiable at $x = 1$.

93. $f(x) = \begin{cases} (x - 1)^3, & x \leq 1 \\ (x - 1)^2, & x > 1 \end{cases}$

The derivative from the left is

$$\begin{aligned} \lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{x - 1} &= \lim_{x \rightarrow 1^-} \frac{(x - 1)^3 - 0}{x - 1} \\ &= \lim_{x \rightarrow 1^-} (x - 1)^2 = 0. \end{aligned}$$

The derivative from the right is

$$\begin{aligned} \lim_{x \rightarrow 1^+} \frac{f(x) - f(1)}{x - 1} &= \lim_{x \rightarrow 1^+} \frac{(x - 1)^2 - 0}{x - 1} \\ &= \lim_{x \rightarrow 1^+} (x - 1) = 0. \end{aligned}$$

These one-sided limits are equal. Therefore, f is differentiable at $x = 1$. ($f'(1) = 0$)

94. $f(x) = \begin{cases} x, & x \leq 1 \\ x^2, & x > 1 \end{cases}$

The derivative from the left is

$$\lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{x - 1} = \lim_{x \rightarrow 1^-} \frac{x - 1}{x - 1} = \lim_{x \rightarrow 1^-} 1 = 1.$$

The derivative from the right is

$$\lim_{x \rightarrow 1^+} \frac{f(x) - f(1)}{x - 1} = \lim_{x \rightarrow 1^+} \frac{x^2 - 1}{x - 1} = \lim_{x \rightarrow 1^+} (x + 1) = 2.$$

These one-sided limits are not equal. Therefore, f is not differentiable at $x = 1$.

95. Note that f is continuous at $x = 2$. $f(x) = \begin{cases} x^2 + 1, & x \leq 2 \\ 4x - 3, & x > 2 \end{cases}$

The derivative from the left is $\lim_{x \rightarrow 2^-} \frac{f(x) - f(2)}{x - 2} = \lim_{x \rightarrow 2^-} \frac{(x^2 + 1) - 5}{x - 2} = \lim_{x \rightarrow 2^-} (x + 2) = 4$.

The derivative from the right is $\lim_{x \rightarrow 2^+} \frac{f(x) - f(2)}{x - 2} = \lim_{x \rightarrow 2^+} \frac{(4x - 3) - 5}{x - 2} = \lim_{x \rightarrow 2^+} 4 = 4$.

The one-sided limits are equal. Therefore, f is differentiable at $x = 2$. ($f'(2) = 4$)

96. Note that f is continuous at $x = 2$. $f(x) = \begin{cases} \frac{1}{2}x + 1, & x < 2 \\ \sqrt{2x}, & x \geq 2 \end{cases}$

The derivative from the left is

$$\lim_{x \rightarrow 2^-} \frac{f(x) - f(2)}{x - 2} = \lim_{x \rightarrow 2^-} \frac{\left(\frac{1}{2}x + 1\right) - 2}{x - 2} = \lim_{x \rightarrow 2^-} \frac{\frac{1}{2}(x - 2)}{x - 2} = \frac{1}{2}.$$

The derivative from the right is



$$\begin{aligned} \lim_{x \rightarrow 2^+} \frac{f(x) - f(2)}{x - 2} &= \lim_{x \rightarrow 2^+} \frac{\sqrt{2x} - 2}{x - 2} \cdot \frac{\sqrt{2x} + 2}{\sqrt{2x} + 2} \\ &= \lim_{x \rightarrow 2^+} \frac{2x - 4}{(x - 2)(\sqrt{2x} + 2)} = \lim_{x \rightarrow 2^+} \frac{2(x - 2)}{(x - 2)(\sqrt{2x} + 2)} = \lim_{x \rightarrow 2^+} \frac{2}{\sqrt{2x} + 2} = \frac{1}{2}. \end{aligned}$$

The one-sided limits are equal. Therefore, f is differentiable at $x = 2$. ($f'(2) = \frac{1}{2}$)

97. (a) The distance from $(3, 1)$ to the line $mx - y + 4 = 0$ is


$$\begin{aligned} d &= \frac{|Ax_1 + By_1 + C|}{\sqrt{A^2 + B^2}} \\ &= \frac{|m(3) - 1(1) + 4|}{\sqrt{m^2 + 1}} = \frac{|3m + 3|}{\sqrt{m^2 + 1}}. \end{aligned}$$

(b)




The function d is not differentiable at $m = -1$. This corresponds to the line $y = -x + 4$, which passes through the point $(3, 1)$.

98. (a) $f(x) = x^2$ and $f'(x) = 2x$



(b) $g(x) = x^3$ and $g'(x) = 3x^2$



- (c) The derivative is a polynomial of degree 1 less than the original function. If $h(x) = x^n$, then $h'(x) = nx^{n-1}$.

(d) If $f(x) = x^4$, then

$$\begin{aligned}f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^4 - x^4}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{x^4 + 4x^3(\Delta x) + 6x^2(\Delta x)^2 + 4x(\Delta x)^3 + (\Delta x)^4 - x^4}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} \frac{\Delta x(4x^3 + 6x^2(\Delta x) + 4x(\Delta x)^2 + (\Delta x)^3)}{\Delta x} \\&= \lim_{\Delta x \rightarrow 0} (4x^3 + 6x^2(\Delta x) + 4x(\Delta x)^2 + (\Delta x)^3) = 4x^3.\end{aligned}$$

Hence, if $f(x) = x^4$, then $f'(x) = 4x^3$ which is consistent with the conjecture. However, this is not a proof since you must verify the conjecture for all integer values of n , $n \geq 2$.

99. False. The slope is $\lim_{\Delta x \rightarrow 0} \frac{f(2 + \Delta x) - f(2)}{\Delta x}$.

100. False. $y = |x - 2|$ is continuous at $x = 2$, but is not differentiable at $x = 2$. (Sharp turn in the graph)

101. False. If the derivative from the left of a point does not equal the derivative from the right of a point, then the derivative does not exist at that point. For example, if $f(x) = |x|$, then the derivative from the left at $x = 0$ is -1 and the derivative from the right at $x = 0$ is 1 . At $x = 0$, the derivative does not exist.

102. True—see Theorem 2.1.

103. $f(x) = \begin{cases} x \sin(1/x), & x \neq 0 \\ 0, & x = 0 \end{cases}$

Using the Squeeze Theorem, we have $-|x| \leq x \sin(1/x) \leq |x|$, $x \neq 0$. Thus, $\lim_{x \rightarrow 0} x \sin(1/x) = 0 = f(0)$ and f is continuous at $x = 0$. Using the alternative form of the derivative, we have

$$\lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{x \sin(1/x) - 0}{x - 0} = \lim_{x \rightarrow 0} \left(\sin \frac{1}{x} \right).$$

Since this limit does not exist ($\sin(1/x)$ oscillates between -1 and 1), the function is not differentiable at $x = 0$.


$$g(x) = \begin{cases} x^2 \sin(1/x), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

Using the Squeeze Theorem again, we have $-x^2 \leq x^2 \sin(1/x) \leq x^2$, $x \neq 0$. Thus, $\lim_{x \rightarrow 0} x^2 \sin(1/x) = 0 = g(0)$ and g is continuous at $x = 0$. Using the alternative form of the derivative again, we have

$$\lim_{x \rightarrow 0} \frac{g(x) - g(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{x^2 \sin(1/x) - 0}{x - 0} = \lim_{x \rightarrow 0} x \sin \frac{1}{x} = 0.$$

Therefore, g is differentiable at $x = 0$, $g'(0) = 0$.

104.



As you zoom in, the graph of $y_1 = x^2 + 1$ appears to be locally the graph of a horizontal line, whereas the graph of $y_2 = |x| + 1$ always has a sharp corner at $(0, 1)$. y_2 is not differentiable at $(0, 1)$.

Section 2.2 Basic Differentiation Rules and Rates of Change

1. (a) $y = x^{1/2}$

$y' = \frac{1}{2}x^{-1/2}$

$y'(1) = \frac{1}{2}$

(b) $y = x^3$

$y' = 3x^2$

$y'(1) = 3$

2. (a) $y = x^{-1/2}$

$y' = -\frac{1}{2}x^{-3/2}$

$y'(1) = -\frac{1}{2}$

(b) $y = x^{-1}$

$y' = -x^{-2}$

$y'(1) = -1$

3. $y = 8$

$y' = 0$

4. $f(x) = -2$

$f'(x) = 0$

5. $y = x^6$

$y' = 6x^5$

6. $y = x^8$

$y' = 8x^7$

7. $y = \frac{1}{x^7} = x^{-7}$

$y' = -7x^{-8} = \frac{-7}{x^8}$

8. $y = \frac{1}{x^8} = x^{-8}$

$y' = -8x^{-9} = \frac{-8}{x^9}$

9. $y = \sqrt[5]{x} = x^{1/5}$

$y' = \frac{1}{5}x^{-4/5} = \frac{1}{5x^{4/5}}$

10. $y = \sqrt[4]{x} = x^{1/4}$

$y' = \frac{1}{4}x^{-3/4} = \frac{1}{4x^{3/4}}$

11. $f(x) = x + 1$

$f'(x) = 1$

12. $g(x) = 3x - 1$

$g'(x) = 3$

13. $f(t) = -2t^2 + 3t - 6$

$f'(t) = -4t + 3$

14. $y = t^2 + 2t - 3$

$y' = 2t + 2$

15. $g(x) = x^2 + 4x^3$

$g'(x) = 2x + 12x^2$

16. $y = 8 - x^3$

$y' = -3x^2$

17. $s(t) = t^3 - 2t + 4$

$s'(t) = 3t^2 - 2$

18. $f(x) = 2x^3 - x^2 + 3x$

$f'(x) = 6x^2 - 2x + 3$

19. $y = \frac{\pi}{2} \sin \theta - \cos \theta$

$y' = \frac{\pi}{2} \cos \theta + \sin \theta$

20. $g(t) = \pi \cos t$

$g'(t) = -\pi \sin t$

21. $y = x^2 - \frac{1}{2} \cos x$

$y' = 2x + \frac{1}{2} \sin x$

22. $y = 5 + \sin x$

$y' = \cos x$

23. $y = \frac{1}{x} - 3 \sin x$

$y' = -\frac{1}{x^2} - 3 \cos x$

24. $y = \frac{5}{(2x)^3} + 2 \cos x = \frac{5}{8}x^{-3} + 2 \cos x$

$y' = \frac{5}{8}(-3)x^{-4} - 2 \sin x = \frac{-15}{8x^4} - 2 \sin x$

FunctionRewriteDifferentiateSimplify

25. $y = \frac{5}{2x^2}$

$y = \frac{5}{2}x^{-2}$

$y' = -5x^{-3}$

$y' = \frac{-5}{x^3}$

26. $y = \frac{2}{3x^2}$

$y = \frac{2}{3}x^{-2}$

$y' = -\frac{4}{3}x^{-3}$

$y' = -\frac{4}{3x^3}$

27. $y = \frac{3}{(2x)^3}$

$y = \frac{3}{8}x^{-3}$

$y' = -\frac{9}{8}x^{-4}$

$y' = \frac{-9}{8x^4}$

<u>Function</u>	<u>Rewrite</u>	<u>Differentiate</u>	<u>Simplify</u>
28. $y = \frac{\pi}{(3x)^2}$	$y = \frac{\pi}{9}x^{-2}$	$y' = -\frac{2\pi}{9}x^{-3}$	$y' = -\frac{2\pi}{9x^3}$
29. $y = \frac{\sqrt{x}}{x}$	$y = x^{-1/2}$	$y' = -\frac{1}{2}x^{-3/2}$	$y' = -\frac{1}{2x^{3/2}}$
30. $y = \frac{4}{x^{-3}}$	$y = 4x^3$	$y' = 12x^2$	$y' = 12x^2$
31. $f(x) = \frac{3}{x^2} = 3x^{-2}, (1, 3)$	32. $f(t) = 3 - \frac{3}{5t}, \left(\frac{3}{5}, 2\right)$	33. $f(x) = -\frac{1}{2} + \frac{7}{5}x^3, \left(0, -\frac{1}{2}\right)$	
$f'(x) = -6x^{-3} = \frac{-6}{x^3}$	$f'(t) = \frac{3}{5t^2}$	$f'(x) = \frac{21}{5}x^2$	
$f'(1) = -6$	$f'\left(\frac{3}{5}\right) = \frac{5}{3}$	$f'(0) = 0$	
34. $y = 3x^3 - 6, (2, 18)$	35. $y = (2x + 1)^2, (0, 1)$	36. $f(x) = 3(5 - x)^2, (5, 0)$	
$y' = 9x^2$	$= 4x^2 + 4x + 1$	$= 3x^2 - 30x + 75$	
$y'(2) = 36$	$y' = 8x + 4$	$f'(x) = 6x - 30$	
	$y'(0) = 4$	$f'(5) = 0$	
37. $f(\theta) = 4 \sin \theta - \theta, (0, 0)$	38. $g(t) = 2 + 3 \cos t, (\pi, -1)$	39. $f(x) = x^2 + 5 - 3x^{-2}$	
$f'(\theta) = 4 \cos \theta - 1$	$g'(t) = -3 \sin t$	$f'(x) = 2x + 6x^{-3} = 2x + \frac{6}{x^3}$	
$f'(0) = 4(1) - 1 = 3$	$g'(\pi) = 0$		
40. $f(x) = x^2 - 3x - 3x^{-2}$	41. $g(t) = t^2 - \frac{4}{t^3} = t^2 - 4t^{-3}$	42. $f(x) = x + x^{-2}$	
$f'(x) = 2x - 3 + 6x^{-3}$	$g'(t) = 2t + 12t^{-4} = 2t + \frac{12}{t^4}$	$f'(x) = 1 - 2x^{-3}$	
$= 2x - 3 + \frac{6}{x^3}$		$= 1 - \frac{2}{x^3}$	
43. $f(x) = \frac{x^3 - 3x^2 + 4}{x^2} = x - 3 + 4x^{-2}$	44. $h(x) = \frac{2x^2 - 3x + 1}{x} = 2x - 3 + x^{-1}$		
$f'(x) = 1 - \frac{8}{x^3} = \frac{x^3 - 8}{x^3}$	$h'(x) = 2 - \frac{1}{x^2} = \frac{2x^2 - 1}{x^2}$		
45. $y = x(x^2 + 1) = x^3 + x$	46. $y = 3x(6x - 5x^2) = 18x^2 - 15x^3$		
$y' = 3x^2 + 1$	$y' = 36x - 45x^2$		
47. $f(x) = \sqrt{x} - 6\sqrt[3]{x} = x^{1/2} - 6x^{1/3}$	48. $f(x) = \sqrt[3]{x} + \sqrt[5]{x} = x^{1/3} + x^{1/5}$		
$f'(x) = \frac{1}{2}x^{-1/2} - 2x^{-2/3} = \frac{1}{2\sqrt{x}} - \frac{2}{x^{2/3}}$	$f'(x) = \frac{1}{3}x^{-2/3} + \frac{1}{5}x^{-4/5} = \frac{1}{3x^{2/3}} + \frac{1}{5x^{4/5}}$		
49. $h(s) = s^{4/5} - s^{2/3}$	50. $f(t) = t^{2/3} - t^{1/3} + 4$		
$h'(s) = \frac{4}{5}s^{-1/5} - \frac{2}{3}s^{-1/3} = \frac{4}{5s^{1/5}} - \frac{2}{3s^{1/3}}$	$f'(t) = \frac{2}{3}t^{-1/3} - \frac{1}{3}t^{-2/3} = \frac{2}{3t^{1/3}} - \frac{1}{3t^{2/3}}$		

51. $f(x) = 6\sqrt{x} + 5 \cos x = 6x^{1/2} + 5 \cos x$
 $f'(x) = 3x^{-1/2} - 5 \sin x = \frac{3}{\sqrt{x}} - 5 \sin x$


53. (a) $y = x^4 - 3x^2 + 2$

$$y' = 4x^3 - 6x$$

$$\text{At } (1, 0): y' = 4(1)^3 - 6(1) = -2$$

$$\text{Tangent line: } y - 0 = -2(x - 1)$$

$$2x + y - 2 = 0$$



55. (a) $f(x) = \frac{2}{\sqrt[4]{x^3}} = 2x^{-3/4}$


$$f'(x) = \frac{-3}{2}x^{-7/4} = \frac{-3}{2x^{7/4}}$$

$$\text{At } (1, 2): f'(1) = \frac{-3}{2}$$

$$\text{Tangent line: } y - 2 = -\frac{3}{2}(x - 1)$$

$$y = -\frac{3}{2}x + \frac{7}{2}$$

$$3x + 2y - 7 = 0$$



57. $y = x^4 - 8x^2 + 2$

$$y' = 4x^3 - 16x$$

$$= 4x(x^2 - 4)$$

$$= 4x(x - 2)(x + 2)$$

$$y' = 0 \Rightarrow x = 0, \pm 2$$

Horizontal tangents: $(0, 2), (2, -14), (-2, -14)$

59. $y = \frac{1}{x^2} = x^{-2}$

$$y' = -2x^{-3} = \frac{-2}{x^3} \text{ cannot equal zero.}$$

Therefore, there are no horizontal tangents.

52. $f(x) = \frac{2}{\sqrt[3]{x}} + 3 \cos x = 2x^{-1/3} + 3 \cos x$

$$f'(x) = \frac{-2}{3}x^{-4/3} - 3 \sin x = \frac{-2}{3x^{4/3}} - 3 \sin x$$


54. (a) $y = x^3 + x$

$$y' = 3x^2 + 1$$

$$\text{At } (-1, -2): y' = 3(-1)^2 + 1 = 4$$

$$\text{Tangent line: } y + 2 = 4(x + 1)$$

$$4x - y + 2 = 0$$



56. (a) $y = (x^2 + 2x)(x + 1)$


$$= x^3 + 3x^2 + 2x$$

$$y' = 3x^2 + 6x + 2$$

$$\text{At } (1, 6): y' = 3(1)^2 + 6(1) + 2 = 11$$

$$\text{Tangent line: } y - 6 = 11(x - 1)$$

$$11x - y - 5 = 0$$



58. $y = x^3 + x$

$$y' = 3x^2 + 1 > 0 \text{ for all } x.$$

Therefore, there are no horizontal tangents.

60. $y = x^2 + 1$

$$y' = 2x = 0 \Rightarrow x = 0$$

$$\text{At } x = 0, y = 1.$$

Horizontal tangent: $(0, 1)$

61. $y = x + \sin x, 0 \leq x < 2\pi$

$$y' = 1 + \cos x = 0$$

$$\cos x = -1 \Rightarrow x = \pi$$

$$\text{At } x = \pi: y = \pi$$

$$\text{Horizontal tangent: } (\pi, \pi)$$

62. $y = \sqrt{3}x + 2 \cos x, 0 \leq x < 2\pi$

$$y' = \sqrt{3} - 2 \sin x = 0$$

$$\sin x = \frac{\sqrt{3}}{2} \Rightarrow x = \frac{\pi}{3} \text{ or } \frac{2\pi}{3}$$

$$\text{At } x = \frac{\pi}{3}: y = \frac{\sqrt{3}\pi + 3}{3}$$

$$\text{At } x = \frac{2\pi}{3}: y = \frac{2\sqrt{3}\pi - 3}{3}$$

$$\text{Horizontal tangents: } \left(\frac{\pi}{3}, \frac{\sqrt{3}\pi + 3}{3}\right), \left(\frac{2\pi}{3}, \frac{2\sqrt{3}\pi - 3}{3}\right)$$

63. $x^2 - kx = 4x - 9$ Equate functions.

$$2x - k = 4 \quad \text{Equate derivatives.}$$

$$\text{Hence, } k = 2x - 4 \text{ and}$$

$$x^2 - (2x - 4)x = 4x - 9 \Rightarrow -x^2 = -9 \Rightarrow x = \pm 3.$$

$$\text{For } x = 3, k = 2 \text{ and for } x = -3, k = -10.$$

65. $\frac{k}{x} = -\frac{3}{4}x + 3 \quad \text{Equate functions.}$

$$-\frac{k}{x^2} = -\frac{3}{4} \quad \text{Equate derivatives.}$$

$$\text{Hence, } k = \frac{3}{4}x^2 \text{ and}$$

$$\frac{\frac{3}{4}x^2}{x} = -\frac{3}{4}x + 3 \Rightarrow \frac{3}{4}x = -\frac{3}{4}x + 3$$

$$\Rightarrow \frac{3}{2}x = 3 \Rightarrow x = 2 \Rightarrow k = 3.$$

64. $k - x^2 = -4x + 7 \quad \text{Equate functions.}$

$$-2x = -4 \quad \text{Equate derivatives.}$$

$$\text{Hence, } x = 2 \text{ and } k - 4 = -8 + 7 \Rightarrow k = 3.$$

66. $k\sqrt{x} = x + 4 \quad \text{Equate functions.}$

$$\frac{k}{2\sqrt{x}} = 1 \quad \text{Equate derivatives.}$$


$$\text{Hence, } k = 2\sqrt{x} \text{ and}$$

$$(2\sqrt{x})\sqrt{x} = x + 4 \Rightarrow 2x = x + 4 \Rightarrow x = 4 \Rightarrow k = 4.$$


67. (a) The slope appears to be steepest between A and B.

(b) The average rate of change between A and B is greater than the instantaneous rate of change at B.

(c)




68. The graph of a function f such that $f' > 0$ for all x and the rate of change of the function is decreasing (i.e., $f'' < 0$) would, in general, look like the graph at the right.




69. $g(x) = f(x) + 6 \Rightarrow g'(x) = f'(x)$

70. $g(x) = -5f(x) \Rightarrow g'(x) = -5f'(x)$

71.



72.



If f is linear then its derivative is a constant function.

$$f(x) = ax + b$$

$$f'(x) = a$$

If f is quadratic, then its derivative is a linear function.

$$f(x) = ax^2 + bx + c$$

$$f'(x) = 2ax + b$$

73. Let (x_1, y_1) and (x_2, y_2) be the points of tangency on $y = x^2$ and $y = -x^2 + 6x - 5$, respectively.

The derivatives of these functions are:

$$y' = 2x \Rightarrow m = 2x_1 \text{ and } y' = -2x + 6 \Rightarrow m = -2x_2 + 6$$

$$m = 2x_1 = -2x_2 + 6$$

$$x_1 = -x_2 + 3$$

Since $y_1 = x_1^2$ and $y_2 = -x_2^2 + 6x_2 - 5$:

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{(-x_2^2 + 6x_2 - 5) - (x_1^2)}{x_2 - x_1} = -2x_2 + 6$$

$$\frac{(-x_2^2 + 6x_2 - 5) - (-x_2 + 3)^2}{x_2 - (-x_2 + 3)} = -2x_2 + 6$$

$$(-x_2^2 + 6x_2 - 5) - (x_2^2 - 6x_2 + 9) = (-2x_2 + 6)(2x_2 - 3)$$

$$-2x_2^2 + 12x_2 - 14 = -4x_2^2 + 18x_2 - 18$$

$$2x_2^2 - 6x_2 + 4 = 0$$

$$2(x_2 - 2)(x_2 - 1) = 0$$

$$x_2 = 1 \text{ or } 2$$

$$x_2 = 1 \Rightarrow y_2 = 0, x_1 = 2 \text{ and } y_1 = 4$$



Thus, the tangent line through $(1, 0)$ and $(2, 4)$ is

$$y - 0 = \left(\frac{4 - 0}{2 - 1}\right)(x - 1) \Rightarrow y = 4x - 4.$$

$$x_2 = 2 \Rightarrow y_2 = 3, x_1 = 1 \text{ and } y_1 = 1$$

Thus, the tangent line through $(2, 3)$ and $(1, 1)$ is

$$y - 1 = \left(\frac{3 - 1}{2 - 1}\right)(x - 1) \Rightarrow y = 2x - 1.$$



74. m_1 is the slope of the line tangent to $y = x$. m_2 is the slope of the line tangent to $y = 1/x$. Since

$$y = x \Rightarrow y' = 1 \Rightarrow m_1 = 1 \text{ and } y = \frac{1}{x} \Rightarrow y' = \frac{-1}{x^2} \Rightarrow m_2 = \frac{-1}{x^2}.$$

The points of intersection of $y = x$ and $y = 1/x$ are

$$x = \frac{1}{x} \Rightarrow x^2 = 1 \Rightarrow x = \pm 1.$$

At $x = \pm 1$, $m_2 = -1/m_1$. Since $m_2 = -1/m_1$, these tangent lines are perpendicular at the points of intersection.

75. $f(x) = 3x + \sin x + 2$

$f'(x) = 3 + \cos x$

Since $|\cos x| \leq 1$, $f'(x) \neq 0$ for all x and f does not have a horizontal tangent line.

77. $f(x) = \sqrt{x}, (-4, 0)$

$f'(x) = \frac{1}{2}x^{-1/2} = \frac{1}{2\sqrt{x}}$

$\frac{1}{2\sqrt{x}} = \frac{0-y}{-4-x}$

$4+x = 2\sqrt{xy}$

$4+x = 2\sqrt{x}\sqrt{x}$

$4+x = 2x$

$x = 4, y = 2$

The point $(4, 2)$ is on the graph of f .

Tangent line: $y - 2 = \frac{0-2}{-4-4}(x - 4)$

$4y - 8 = x - 4$

$0 = x - 4y + 4$

76. $f(x) = x^5 + 3x^3 + 5x$

$f'(x) = 5x^4 + 9x^2 + 5$

Since $5x^4 + 9x^2 \geq 0$, $f'(x) \geq 5$. Thus, f does not have a tangent line with a slope of 3.

78. $f(x) = \frac{2}{x}, (5, 0)$

$f'(x) = -\frac{2}{x^2}$

$-\frac{2}{x^2} = \frac{0-y}{5-x}$

$-10 + 2x = -x^2y$

$-10 + 2x = -x^2\left(\frac{2}{x}\right)$

$-10 + 2x = -2x$

$4x = 10$

$x = \frac{5}{2}, y = \frac{4}{5}$


The point $(\frac{5}{2}, \frac{4}{5})$ is on the graph of f . The slope of the tangent line is $f'(\frac{5}{2}) = -\frac{8}{25}$.

Tangent line: $y - \frac{4}{5} = -\frac{8}{25}\left(x - \frac{5}{2}\right)$


$25y - 20 = -8x + 20$

$8x + 25y - 40 = 0$

79. $f'(1) = -1$



80. $f'(4) = 1$



81. (a) One possible secant is between $(3.9, 7.7019)$ and $(4, 8)$:

$y - 8 = \frac{8 - 7.7019}{4 - 3.9}(x - 4)$


$y - 8 = 2.981(x - 4)$

$y = S(x) = 2.981x - 3.924$

(b) $f'(x) = \frac{3}{2}x^{1/2} \Rightarrow f'(4) = \frac{3}{2}(2) = 3$


$T(x) = 3(x - 4) + 8 = 3x - 4$

$S(x)$ is an approximation of the tangent line $T(x)$.



81. —CONTINUED—

- (c) As you move further away from $(4, 8)$, the accuracy of the approximation T gets worse.



(d)


Δx	-3	-2	-1	-0.5	-0.1	0	0.1	0.5	1	2	3
$f(4 + \Delta x)$	1	2.828	5.196	6.548	7.702	8	8.302	9.546	11.180	14.697	18.520
$T(4 + \Delta x)$	-1	2	5	6.5	7.7	8	8.3	9.5	11	14	17

82. (a) Nearby point: $(1.0073138, 1.0221024)$

$$\text{Secant line: } y - 1 = \frac{1.0221024 - 1}{1.0073138 - 1}(x - 1)$$

$$y = 3.022(x - 1) + 1$$


(Answers will vary.)



$$(b) f''(x) = 3x^2$$

$$T(x) = 3(x - 1) + 1 = 3x - 2$$

- (c) The accuracy worsens as you move away from $(1, 1)$.



(d)

Δx	-3	-2	-1	-0.5	-0.1	0	0.1	0.5	1	2	3
$f(x)$	-8	-1	0	0.125	0.729	1	1.331	3.375	8	27	64
$T(x)$	-8	-5	-2	-0.5	0.7	1	1.3	2.5	4	7	10

The accuracy decreases more rapidly than in Exercise 81 because $y = x^3$ is less “linear” than $y = x^{3/2}$.

83. False. Let $f(x) = x^2$ and $g(x) = x^2 + 4$. Then $f'(x) = g'(x) = 2x$, but $f(x) \neq g(x)$.

84. True. If $f(x) = g(x) + c$, then $f'(x) = g'(x) + 0 = g'(x)$.

85. False. If $y = \pi^2$, then $dy/dx = 0$. (π^2 is a constant.)

86. True. If $y = x/\pi = (1/\pi) \cdot x$, then $dy/dx = (1/\pi)(1) = 1/\pi$.

87. True. If $g(x) = 3f(x)$, then $g'(x) = 3f'(x)$.

88. False. If $f(x) = \frac{1}{x^n} = x^{-n}$, then $f'(x) = -nx^{-n-1} = \frac{-n}{x^{n+1}}$.

89. $f(t) = 2t + 7, [1, 2]$

90. $f(t) = t^2 - 3, [2, 2.1]$

$$f'(t) = 2$$

$$f'(t) = 2t$$

Instantaneous rate of change is the constant 2.

Instantaneous rate of change:

Average rate of change:

$$(2, 1) \Rightarrow f'(2) = 2(2) = 4$$

$$(2.1, 1.41) \Rightarrow f'(2.1) = 4.2$$

(These are the same because f is a line of slope 2.)

Average rate of change:

$$\frac{f(2) - f(1)}{2 - 1} = \frac{[2(2) + 7] - [2(1) + 7]}{1} = 2$$

$$\frac{f(2.1) - f(2)}{2.1 - 2} = \frac{1.41 - 1}{0.1} = 4.1$$

91. $f(x) = -\frac{1}{x}$, $[1, 2]$

$$f'(x) = \frac{1}{x^2}$$

Instantaneous rate of change:

$$(1, -1) \Rightarrow f'(1) = 1$$

$$\left(2, -\frac{1}{2}\right) \Rightarrow f'(2) = \frac{1}{4}$$

Average rate of change:

$$\frac{f(2) - f(1)}{2 - 1} = \frac{(-1/2) - (-1)}{2 - 1} = \frac{1}{2}$$

93. (a) $s(t) = -16t^2 + 1362$

$$v(t) = -32t$$

(b) $\frac{s(2) - s(1)}{2 - 1} = 1298 - 1346 = -48$ ft/sec

(c) $v(t) = s'(t) = -32t$

When $t = 1$: $v(1) = -32$ ft/sec

When $t = 2$: $v(2) = -64$ ft/sec

(d) $-16t^2 + 1362 = 0$

$$t^2 = \frac{1362}{16} \Rightarrow t = \frac{\sqrt{1362}}{4} \approx 9.226 \text{ sec}$$

(e) $v\left(\frac{\sqrt{1362}}{4}\right) = -32\left(\frac{\sqrt{1362}}{4}\right)$

$$= -8\sqrt{1362} \approx -295.242 \text{ ft/sec}$$

95. $s(t) = -4.9t^2 + v_0 t + s_0$

$$= -4.9t^2 + 120t$$

$$v(t) = -9.8t + 120$$

$$v(5) = -9.8(5) + 120 = 71 \text{ m/sec}$$


$$v(10) = -9.8(10) + 120 = 22 \text{ m/sec}$$

97. From $(0, 0)$ to $(4, 2)$, $s(t) = \frac{1}{2}t \Rightarrow v(t) = \frac{1}{2} \text{ mi/min.}$

$$v(t) = \frac{1}{2}(60) = 30 \text{ mph for } 0 < t < 4$$

Similarly, $v(t) = 0$ for $4 < t < 6$. Finally, from $(6, 2)$ to $(10, 6)$,

$$s(t) = t - 4 \Rightarrow v(t) = 1 \text{ mi/in} = 60 \text{ mph.}$$



92. $f(x) = \sin x$, $\left[0, \frac{\pi}{6}\right]$

$$f'(x) = \cos x$$

Instantaneous rate of change:

$$(0, 0) \Rightarrow f'(0) = 1$$

$$\left(\frac{\pi}{6}, \frac{1}{2}\right) \Rightarrow f'\left(\frac{\pi}{6}\right) = \frac{\sqrt{3}}{2} \approx 0.866$$

Average rate of change:

$$\frac{f(\pi/6) - f(0)}{(\pi/6) - 0} = \frac{(1/2) - 0}{(\pi/6) - 0} = \frac{3}{\pi} \approx 0.955$$

94. $s(t) = -16t^2 - 22t + 220$

$$v(t) = -32t - 22$$

$$v(3) = -118 \text{ ft/sec}$$

$$s(t) = -16t^2 - 22t + 220$$

$$= 112 \text{ (height after falling 108 ft)}$$

$$-16t^2 - 22t + 108 = 0$$

$$-2(t - 2)(8t + 27) = 0$$

$$t = 2$$

$$v(2) = -32(2) - 22$$


$$= -86 \text{ ft/sec}$$

96. $s(t) = -4.9t^2 + v_0 t + s_0$

$$= -4.9t^2 + s_0 = 0 \text{ when } t = 6.8.$$

$$s_0 = 4.9t^2 = 4.9(6.8)^2 \approx 226.6 \text{ m}$$

98.



(The velocity has been converted to miles per hour.)

99. $v = 40 \text{ mph} = \frac{2}{3} \text{ mi/min}$


$$\left(\frac{2}{3} \text{ mi/min}\right)(6 \text{ min}) = 4 \text{ mi}$$

$$v = 0 \text{ mph} = 0 \text{ mi/min}$$


$$(0 \text{ mi/min})(2 \text{ min}) = 0 \text{ mi}$$

$$v = 60 \text{ mph} = 1 \text{ mi/min}$$

$$(1 \text{ mi/min})(2 \text{ min}) = 2 \text{ mi}$$



100. This graph corresponds with Exercise 97.



101. (a) Using a graphing utility,


$$R = 0.417v - 0.02.$$

(b) Using a graphing utility,

$$B = 0.00557v^2 + 0.0014v + 0.04.$$

(c) $T = R + B = 0.00557v^2 + 0.418v + 0.02$

(d)



(e) $\frac{dT}{dv} = 0.01114v + 0.418$

For $v = 40$, $T'(40) \approx 0.86$.

For $v = 80$, $T'(80) \approx 1.31$.

For $v = 100$, $T'(100) \approx 1.53$.

(f) For increasing speeds, the total stopping distance increases.

102. $C = (\text{gallons of fuel used})(\text{cost per gallon})$

$$= \left(\frac{15,000}{x}\right)(1.55) = \frac{23,250}{x}$$

$$\frac{dC}{dx} = -\frac{23,250}{x^2}$$

x	10	15	20	25	30	35	40
C	2325	1550	1163	930	775	664	581
$\frac{dC}{dx}$	-233	-103	-58	-37	-26	-19	-15

The driver who gets 15 miles per gallon would benefit more. The rate of change at $x = 15$ is larger in absolute value than that at $x = 35$.

103. $V = s^3$, $\frac{dV}{ds} = 3s^2$

When $s = 4 \text{ cm}$, $\frac{dV}{ds} = 48 \text{ cm}^2 \text{ per cm change in } s$.

104. $A = s^2$, $\frac{dA}{ds} = 2s$

When $s = 4 \text{ m}$,

$$\frac{dA}{ds} = 8 \text{ square meters per meter change in } s.$$

105. $s(t) = -\frac{1}{2}at^2 + c$ and $s'(t) = -at$

Average velocity: $\frac{s(t_0 + \Delta t) - s(t_0 - \Delta t)}{(t_0 + \Delta t) - (t_0 - \Delta t)} = \frac{[-(1/2)a(t_0 + \Delta t)^2 + c] - [-(1/2)a(t_0 - \Delta t)^2 + c]}{2\Delta t}$

$$= \frac{-(1/2)a(t_0^2 + 2t_0\Delta t + (\Delta t)^2) + (1/2)a(t_0^2 - 2t_0\Delta t + (\Delta t)^2)}{2\Delta t}$$

$$= \frac{-2at_0\Delta t}{2\Delta t}$$

$$= -at_0$$

$$= s'(t_0) \text{ instantaneous velocity at } t = t_0$$

106. $C = \frac{1,008,000}{Q} + 6.3Q$

$$\frac{dC}{dQ} = -\frac{1,008,000}{Q^2} + 6.3$$

$$C(351) - C(350) \approx 5083.095 - 5085 \approx -\$1.91$$

When $Q = 350$, $\frac{dC}{dQ} \approx -\$1.93$.

107. $N = f(p)$

(a) $f'(1.479)$ is the rate of change of gallons of gasoline sold when the price is \$1.479 per gallon.

(b) $f'(1.479)$ is usually negative. As prices go up, sales go down.

108. $\frac{dT}{dt} = K(T - T_a)$

109. $y = ax^2 + bx + c$

Since the parabola passes through $(0, 1)$ and $(1, 0)$, we have:

$$(0, 1): 1 = a(0)^2 + b(0) + c \Rightarrow c = 1$$

$$(1, 0): 0 = a(1)^2 + b(1) + 1 \Rightarrow b = -a - 1$$

Thus, $y = ax^2 + (-a - 1)x + 1$. From the tangent line $y = x - 1$, we know that the derivative is 1 at the point $(1, 0)$.

$$y' = 2ax + (-a - 1)$$

$$1 = 2a(1) + (-a - 1)$$

$$1 = a - 1$$

$$a = 2$$

$$b = -a - 1 = -3$$

Therefore, $y = 2x^2 - 3x + 1$.

110. $y = \frac{1}{x}, x > 0$

$$y' = -\frac{1}{x^2}$$


At (a, b) , the equation of the tangent line is

$$y - \frac{1}{a} = -\frac{1}{a^2}(x - a) \quad \text{or} \quad y = -\frac{x}{a^2} + \frac{2}{a}$$

The x -intercept is $(2a, 0)$.

The y -intercept is $(0, \frac{2}{a})$.

The area of the triangle is $A = \frac{1}{2}bh = \frac{1}{2}(2a)\left(\frac{2}{a}\right) = 2$.



111. $y = x^3 - 9x$

$$y' = 3x^2 - 9$$

Tangent lines through $(1, -9)$:

$$y + 9 = (3x^2 - 9)(x - 1)$$

$$(x^3 - 9x) + 9 = 3x^3 - 3x^2 - 9x + 9$$

$$0 = 2x^3 - 3x^2 = x^2(2x - 3)$$

$$x = 0 \text{ or } x = \frac{3}{2}$$

The points of tangency are $(0, 0)$ and $(\frac{3}{2}, -\frac{81}{8})$. At $(0, 0)$, the slope is $y'(0) = -9$. At $(\frac{3}{2}, -\frac{81}{8})$, the slope is $y'(\frac{3}{2}) = -\frac{9}{4}$.

Tangent lines:

$$y - 0 = -9(x - 0) \quad \text{and} \quad y + \frac{81}{8} = -\frac{9}{4}(x - \frac{3}{2})$$

$$y = -9x$$

$$y = -\frac{9}{4}x - \frac{27}{4}$$

$$9x + y = 0$$

$$9x + 4y + 27 = 0$$

112. $y = x^2$

$$y' = 2x$$

(a) Tangent lines through $(0, a)$:

$$y - a = 2x(x - 0)$$

$$x^2 - a = 2x^2$$

$$-a = x^2$$

$$\pm\sqrt{-a} = x$$

The points of tangency are $(\pm\sqrt{-a}, -a)$. At $(\sqrt{-a}, -a)$, the slope is $y'(\sqrt{-a}) = 2\sqrt{-a}$. At $(-\sqrt{-a}, -a)$, the slope is $y'(-\sqrt{-a}) = -2\sqrt{-a}$.

Tangent lines: $y + a = 2\sqrt{-a}(x - \sqrt{-a})$ and $y + a = -2\sqrt{-a}(x + \sqrt{-a})$

$$y = 2\sqrt{-a}x + a$$

$$y = -2\sqrt{-a}x + a$$

Restriction: a must be negative.

(b) Tangent lines through $(a, 0)$:

$$y - 0 = 2x(x - a)$$

$$x^2 = 2x^2 - 2ax$$

$$0 = x^2 - 2ax = x(x - 2a)$$

The points of tangency are $(0, 0)$ and $(2a, 4a^2)$. At $(0, 0)$, the slope is $y'(0) = 0$. At $(2a, 4a^2)$, the slope is $y'(2a) = 4a$.

Tangent lines: $y - 0 = 0(x - 0)$ and $y - 4a^2 = 4a(x - 2a)$

$$y = 0$$

$$y = 4ax - 4a^2$$

Restriction: None, a can be any real number.

113. $f(x) = \begin{cases} ax^3, & x \leq 2 \\ x^2 + b, & x > 2 \end{cases}$

f must be continuous at $x = 2$ to be differentiable at $x = 2$.

$$\left. \begin{array}{l} \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} ax^3 = 8a \\ \lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} (x^2 + b) = 4 + b \end{array} \right\} \begin{array}{l} 8a = 4 + b \\ 8a - 4 = b \end{array}$$

$$f'(x) = \begin{cases} 3ax^2, & x < 2 \\ 2x, & x > 2 \end{cases}$$

For f to be differentiable at $x = 2$, the left derivative must equal the right derivative.

$$3a(2)^2 = 2(2)$$

$$12a = 4$$

$$a = \frac{1}{3}$$

$$b = 8a - 4 = -\frac{4}{3}$$

114. $f(x) = \begin{cases} \cos x, & x < 0 \\ ax + b, & x \geq 0 \end{cases}$

$$f(0) = b = \cos(0) = 1 \Rightarrow b = 1$$

$$f'(x) = \begin{cases} -\sin x, & x < 0 \\ a, & x > 0 \end{cases}$$

$$\text{Hence, } a = 0.$$

$$\text{Answer: } a = 0, b = 1$$

116. Let $f(x) = \cos x$.

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\cos x \cos \Delta x - \sin x \sin \Delta x - \cos x}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\cos x(\cos \Delta x - 1)}{\Delta x} - \lim_{\Delta x \rightarrow 0} \sin x \left(\frac{\sin \Delta x}{\Delta x} \right) \\ &= 0 - \sin x(1) = -\sin x \end{aligned}$$

115. $f_1(x) = |\sin x|$ is differentiable for all $x \neq n\pi, n$ an integer.

$f_2(x) = \sin|x|$ is differentiable for all $x \neq 0$.

You can verify this by graphing f_1 and f_2 and observing the locations of the sharp turns.

Section 2.3 Product and Quotient Rules and Higher-Order Derivatives

1. $g(x) = (x^2 + 1)(x^2 - 2x)$

$$\begin{aligned} g'(x) &= (x^2 + 1)(2x - 2) + (x^2 - 2x)(2x) \\ &= 2x^3 - 2x^2 + 2x - 2 + 2x^3 - 4x^2 \\ &= 4x^3 - 6x^2 + 2x - 2 \end{aligned}$$

3. $h(t) = \sqrt[3]{t^2 + 4} = t^{1/3}(t^2 + 4)$

$$\begin{aligned} h'(t) &= t^{1/3}(2t) + (t^2 + 4)\frac{1}{3}t^{-2/3} \\ &= 2t^{4/3} + \frac{t^2 + 4}{3t^{2/3}} \\ &= \frac{7t^2 + 4}{3t^{2/3}} \end{aligned}$$

2. $f(x) = (6x + 5)(x^3 - 2)$

$$\begin{aligned} f'(x) &= (6x + 5)(3x^2) + (x^3 - 2)(6) \\ &= 18x^3 + 15x^2 + 6x^3 - 12 \\ &= 24x^3 + 15x^2 - 12 \end{aligned}$$

4. $g(s) = \sqrt{s}(4 - s^2) = s^{1/2}(4 - s^2)$

$$\begin{aligned} g'(s) &= s^{1/2}(-2s) + (4 - s^2)\frac{1}{2}s^{-1/2} \\ &= -2s^{3/2} + \frac{4 - s^2}{2s^{1/2}} \\ &= \frac{4 - 5s^2}{2s^{1/2}} \end{aligned}$$

5. $f(x) = x^3 \cos x$

$$\begin{aligned} f'(x) &= x^3(-\sin x) + \cos x(3x^2) \\ &= 3x^2 \cos x - x^3 \sin x \end{aligned}$$

7. $f(x) = \frac{x}{x^2 + 1}$

$$f'(x) = \frac{(x^2 + 1)(1) - x(2x)}{(x^2 + 1)^2} = \frac{1 - x^2}{(x^2 + 1)^2}$$

9. $h(x) = \frac{\sqrt[3]{x}}{x^3 + 1} = \frac{x^{1/3}}{x^3 + 1}$

$$\begin{aligned} h'(x) &= \frac{(x^3 + 1)\frac{1}{3}x^{-2/3} - x^{1/3}(3x^2)}{(x^3 + 1)^2} \\ &= \frac{(x^3 + 1) - x(9x^2)}{3x^{2/3}(x^3 + 1)^2} \\ &= \frac{1 - 8x^3}{3x^{2/3}(x^3 + 1)^2} \end{aligned}$$

11. $g(x) = \frac{\sin x}{x^2}$

$$g'(x) = \frac{x^2(\cos x) - \sin x(2x)}{(x^2)^2} = \frac{x \cos x - 2 \sin x}{x^3}$$

13. $f(x) = (x^3 - 3x)(2x^2 + 3x + 5)$

$$\begin{aligned} f'(x) &= (x^3 - 3x)(4x + 3) + (2x^2 + 3x + 5)(3x^2 - 3) \\ &= 10x^4 + 12x^3 - 3x^2 - 18x - 15 \end{aligned}$$

$$f'(0) = -15$$

15. $f(x) = \frac{x^2 - 4}{x - 3}$

$$\begin{aligned} f'(x) &= \frac{(x - 3)(2x) - (x^2 - 4)(1)}{(x - 3)^2} \\ &= \frac{2x^2 - 6x - x^2 + 4}{(x - 3)^2} \\ &= \frac{x^2 - 6x + 4}{(x - 3)^2} \end{aligned}$$

$$f'(1) = \frac{1 - 6 + 4}{(1 - 3)^2} = -\frac{1}{4}$$

17. $f(x) = x \cos x$

$$f'(x) = (x)(-\sin x) + (\cos x)(1) = \cos x - x \sin x$$

$$f'\left(\frac{\pi}{4}\right) = \frac{\sqrt{2}}{2} - \frac{\pi}{4}\left(\frac{\sqrt{2}}{2}\right) = \frac{\sqrt{2}}{8}(4 - \pi)$$

6. $g(x) = \sqrt{x} \sin x$

$$g'(x) = \sqrt{x} \cos x + \sin x \left(\frac{1}{2\sqrt{x}}\right) = \sqrt{x} \cos x + \frac{1}{2\sqrt{x}} \sin x$$

8. $g(t) = \frac{t^2 + 2}{2t - 7}$

$$g'(t) = \frac{(2t - 7)(2t) - (t^2 + 2)(2)}{(2t - 7)^2} = \frac{2t^2 - 14t - 4}{(2t - 7)^2}$$

10. $h(s) = \frac{s}{\sqrt{s} - 1}$

$$\begin{aligned} h'(s) &= \frac{(\sqrt{s} - 1)(1) - s\left(\frac{1}{2}s^{-1/2}\right)}{(\sqrt{s} - 1)^2} \\ &= \frac{\sqrt{s} - 1 - \frac{1}{2}\sqrt{s}}{(\sqrt{s} - 1)^2} = \frac{\sqrt{s} - 2}{2(\sqrt{s} - 1)^2} \end{aligned}$$

12. $f(t) = \frac{\cos t}{t^3}$

$$f'(t) = \frac{t^3(-\sin t) - \cos t(3t^2)}{(t^3)^2} = -\frac{t \sin t + 3 \cos t}{t^4}$$

14. $f(x) = (x^2 - 2x + 1)(x^3 - 1)$

$$\begin{aligned} f'(x) &= (x^2 - 2x + 1)(3x^2) + (x^3 - 1)(2x - 2) \\ &= 3x^2(x - 1)^2 + 2(x - 1)^2(x^2 + x + 1) \\ &= (x - 1)^2(5x^2 + 2x + 2) \\ f'(1) &= 0 \end{aligned}$$

16. $f(x) = \frac{x + 1}{x - 1}$

$$\begin{aligned} f'(x) &= \frac{(x - 1)(1) - (x + 1)(1)}{(x - 1)^2} \\ &\equiv \frac{x - 1 - x - 1}{(x - 1)^2} \\ &= -\frac{2}{(x - 1)^2} \end{aligned}$$

$$f'(2) = -\frac{2}{(2 - 1)^2} = -2$$

18. $f(x) = \frac{\sin x}{x}$

$$f'(x) = \frac{(x)(\cos x) - (\sin x)(1)}{x^2}$$

$$= \frac{x \cos x - \sin x}{x^2}$$

$$f'\left(\frac{\pi}{6}\right) = \frac{(\pi/6)\left(\sqrt{3}/2\right) - (1/2)}{\pi^2/36}$$

$$= \frac{3\sqrt{3}\pi - 18}{\pi^2}$$

$$= \frac{3(\sqrt{3}\pi - 6)}{\pi^2}$$

FunctionRewriteDifferentiateSimplify

19. $y = \frac{x^2 + 2x}{3}$

$$y = \frac{1}{3}x^2 + \frac{2}{3}x$$

$$y' = \frac{2}{3}x + \frac{2}{3}$$

$$y' = \frac{2x + 2}{3}$$

20. $y = \frac{5x^2 - 3}{4}$

$$y = \frac{5}{4}x^2 - \frac{3}{4}$$

$$y' = \frac{10}{4}x$$

$$y' = \frac{5x}{2}$$

21. $y = \frac{7}{3x^3}$

$$y = \frac{7}{3}x^{-3}$$

$$y' = -7x^{-4}$$

$$y' = -\frac{7}{x^4}$$

22. $y = \frac{4}{5x^2}$

$$y = \frac{4}{5}x^{-2}$$

$$y' = -\frac{8}{5}x^{-3}$$

$$y' = -\frac{8}{5x^3}$$

23. $y = \frac{4x^{3/2}}{x}$

$$y = 4\sqrt{x}, \quad x > 0$$

$$y' = 2x^{-1/2}$$

$$y' = \frac{2}{\sqrt{x}}$$

24. $y = \frac{3x^2 - 5}{7}$

$$y = \frac{3}{7}x^2 - \frac{5}{7}$$

$$y' = \frac{6x}{7}$$

$$y' = \frac{6}{7}x$$

25. $f(x) = \frac{3 - 2x - x^2}{x^2 - 1}$

$$f'(x) = \frac{(x^2 - 1)(-2 - 2x) - (3 - 2x - x^2)(2x)}{(x^2 - 1)^2}$$

$$= \frac{2x^2 - 4x + 2}{(x^2 - 1)^2} = \frac{2(x - 1)^2}{(x^2 - 1)^2}$$

$$= \frac{2}{(x + 1)^2}, \quad x \neq 1$$

26. $f(x) = \frac{x^3 + 3x + 2}{x^2 - 1}$

$$f'(x) = \frac{(x^2 - 1)(3x^2 + 3) - (x^3 + 3x + 2)(2x)}{(x^2 - 1)^2}$$

$$= \frac{x^4 - 6x^2 - 4x - 3}{(x^2 - 1)^2}$$

27. $f(x) = x\left(1 - \frac{4}{x+3}\right) = x - \frac{4x}{x+3}$

$$f'(x) = 1 - \frac{(x+3)4 - 4x(1)}{(x+3)^2}$$

$$= \frac{(x^2 + 6x + 9) - 12}{(x+3)^2}$$

$$= \frac{x^2 + 6x - 3}{(x+3)^2}$$

28. $f(x) = x^4\left[1 - \frac{2}{x+1}\right] = x^4\left[\frac{x-1}{x+1}\right]$

$$f'(x) = x^4\left[\frac{(x+1) - (x-1)}{(x+1)^2}\right] + \left[\frac{x-1}{x+1}\right](4x^3)$$

$$= x^4\left[\frac{2}{(x+1)^2}\right] + \left[\frac{x^2 - 1}{(x+1)^2}\right](4x^3)$$

$$= 2x^3\left[\frac{2x^2 + x - 2}{(x+1)^2}\right]$$

29. $f(x) = \frac{2x+5}{\sqrt{x}} = 2x^{1/2} + 5x^{-1/2}$

$$f'(x) = x^{-1/2} - \frac{5}{2}x^{-3/2} = x^{-3/2}\left[x - \frac{5}{2}\right] = \frac{2x-5}{2x\sqrt{x}} = \frac{2x-5}{2x^{3/2}}$$

30. $f(x) = \sqrt[3]{x}(\sqrt{x} + 3) = x^{1/3}(x^{1/2} + 3)$

$$f'(x) = x^{1/3}\left(\frac{1}{2}x^{-1/2}\right) + (x^{1/2} + 3)\left(\frac{1}{3}x^{-2/3}\right)$$

$$= \frac{5}{6}x^{-1/6} + x^{-2/3}$$

$$= \frac{5}{6x^{1/6}} + \frac{1}{x^{2/3}}$$

Alternate solution:

$$f(x) = \sqrt[3]{x}(\sqrt{x} + 3) \\ = x^{5/6} + 3x^{1/3}$$

$$f'(x) = \frac{5}{6}x^{-1/6} + x^{-2/3}$$

$$= \frac{5}{6x^{1/6}} + \frac{1}{x^{2/3}}$$

31. $h(s) = (s^3 - 2)^2 = s^6 - 4s^3 + 4$

$$h'(s) = 6s^5 - 12s^2 = 6s^2(s^3 - 2)$$

32. $h(x) = (x^2 - 1)^2 = x^4 - 2x^2 + 1$

$$h'(x) = 4x^3 - 4x = 4x(x^2 - 1)$$

33. $f(x) = \frac{2 - (1/x)}{x - 3} = \frac{2x - 1}{x(x - 3)} = \frac{2x - 1}{x^2 - 3x}$

$$f'(x) = \frac{(x^2 - 3x)2 - (2x - 1)(2x - 3)}{(x^2 - 3x)^2} = \frac{2x^2 - 6x - 4x^2 + 8x - 3}{(x^2 - 3x)^2}$$

$$= \frac{-2x^2 + 2x - 3}{(x^2 - 3x)^2} = -\frac{2x^2 - 2x + 3}{x^2(x - 3)^2}$$

34. $g(x) = x^2\left(\frac{2}{x} - \frac{1}{x+1}\right) = 2x - \frac{x^2}{x+1}$

$$g'(x) = 2 - \frac{(x+1)2x - x^2(1)}{(x+1)^2} = \frac{2(x^2 + 2x + 1) - x^2 - 2x}{(x+1)^2} = \frac{x^2 + 2x + 2}{(x+1)^2}$$

35. $f(x) = (3x^3 + 4x)(x - 5)(x + 1)$

$$\begin{aligned} f'(x) &= (9x^2 + 4)(x - 5)(x + 1) + (3x^3 + 4x)(1)(x + 1) + (3x^3 + 4x)(x - 5)(1) \\ &= (9x^2 + 4)(x^2 - 4x - 5) + 3x^4 + 3x^3 + 4x^2 + 4x + 3x^4 - 15x^3 + 4x^2 - 20x \\ &= 9x^4 - 36x^3 - 41x^2 - 16x - 20 + 6x^4 - 12x^3 + 8x^2 - 16x \\ &= 15x^4 - 48x^3 - 33x^2 - 32x - 20 \end{aligned}$$

36. $f(x) = (x^2 - x)(x^2 + 1)(x^2 + x + 1)$

$$\begin{aligned} f'(x) &= (2x - 1)(x^2 + 1)(x^2 + x + 1) + (x^2 - x)(2x)(x^2 + x + 1) + (x^2 - x)(x^2 + 1)(2x + 1) \\ &= (2x - 1)(x^4 + x^3 + 2x^2 + x + 1) + (x^2 - x)(2x^3 + 2x^2 + 2x) + (x^2 - x)(2x^3 + x^2 + 2x + 1) \\ &= 2x^5 + x^4 + 3x^3 + x^2 - 1 + 2x^5 - 2x^2 + 2x^5 - x^4 + x^3 - x^2 - x \\ &= 6x^5 + 4x^3 - 3x^2 - 1 \end{aligned}$$

37. $f(x) = \frac{x^2 + c^2}{x^2 - c^2}$

$$\begin{aligned}f'(x) &= \frac{(x^2 - c^2)(2x) - (x^2 + c^2)(2x)}{(x^2 - c^2)^2} \\&= \frac{-4xc^2}{(x^2 - c^2)^2}\end{aligned}$$

39. $f(t) = t^2 \sin t$

$$\begin{aligned}f'(t) &= t^2 \cos t + 2t \sin t \\&= t(t \cos t + 2 \sin t)\end{aligned}$$

41. $f(t) = \frac{\cos t}{t}$

$$f'(t) = \frac{-t \sin t - \cos t}{t^2} = -\frac{t \sin t + \cos t}{t^2}$$

43. $f(x) = -x + \tan x$

$$f'(x) = -1 + \sec^2 x = \tan^2 x$$

45. $g(t) = \sqrt[4]{t} + 8 \sec t = t^{1/4} + 8 \sec t$

$$g'(t) = \frac{1}{4}t^{-3/4} + 8 \sec t \tan t = \frac{1}{4t^{3/4}} + 8 \sec t \tan t$$

47. $y = \frac{3(1 - \sin x)}{2 \cos x} = \frac{3}{2}(\sec x - \tan x)$

$$\begin{aligned}y' &= \frac{3}{2}(\sec x \tan x - \sec^2 x) = \frac{3}{2} \sec x (\tan x - \sec x) \\&= \frac{3}{2}(\sec x \tan x - \tan^2 x - 1)\end{aligned}$$

49. $y = -\csc x - \sin x$

$$\begin{aligned}y' &= \csc x \cot x - \cos x \\&= \frac{\cos x}{\sin^2 x} - \cos x \\&= \cos x(\csc^2 x - 1) \\&= \cos x \cot^2 x\end{aligned}$$

51. $f(x) = x^2 \tan x$

$$\begin{aligned}f'(x) &= x^2 \sec^2 x + 2x \tan x \\&= x(x \sec^2 x + 2 \tan x)\end{aligned}$$

53. $y = 2x \sin x + x^2 \cos x$

$$\begin{aligned}y' &= 2x \cos x + 2 \sin x + x^2(-\sin x) + 2x \cos x \\&= 4x \cos x + 2 \sin x - x^2 \sin x\end{aligned}$$

38. $f(x) = \frac{c^2 - x^2}{c^2 + x^2}$

$$\begin{aligned}f'(x) &= \frac{(c^2 + x^2)(-2x) - (c^2 - x^2)(2x)}{(c^2 + x^2)^2} \\&= \frac{-4xc^2}{(c^2 + x^2)^2}\end{aligned}$$

40. $f(\theta) = (\theta + 1) \cos \theta$

$$\begin{aligned}f'(\theta) &= (\theta + 1)(-\sin \theta) + (\cos \theta)(1) \\&= \cos \theta - (\theta + 1) \sin \theta\end{aligned}$$

42. $f(x) = \frac{\sin x}{x}$

$$f'(x) = \frac{x \cos x - \sin x}{x^2}$$

44. $y = x + \cot x$

$$y' = 1 - \csc^2 x = -\cot^2 x$$

46. $h(s) = \frac{1}{s} - 10 \csc s$

$$h'(s) = -\frac{1}{s^2} + 10 \csc s \cot s$$

48. $y = \frac{\sec x}{x}$

$$\begin{aligned}y' &= \frac{x \sec x \tan x - \sec x}{x^2} \\&= \frac{\sec x(x \tan x - 1)}{x^2}\end{aligned}$$

50. $y = x \sin x + \cos x$

$$y' = x \cos x + \sin x - \sin x = x \cos x$$

52. $f(x) = \sin x \cos x$

$$\begin{aligned}f'(x) &= \sin x(-\sin x) + \cos x(\cos x) \\&= \cos 2x\end{aligned}$$

54. $h(\theta) = 5\theta \sec \theta + \theta \tan \theta$

$$h'(\theta) = 5\theta \sec \theta \tan \theta + 5 \sec \theta + \theta \sec^2 \theta + \tan \theta$$

55. $g(x) = \left(\frac{x+1}{x+2}\right)(2x-5)$

$$g'(x) = \frac{2x^2 + 8x - 1}{(x+2)^2} \quad (\text{Form of answer may vary.})$$

57. $g(\theta) = \frac{\theta}{1 - \sin \theta}$

$$g''(\theta) = \frac{1 - \sin \theta + \theta \cos \theta}{(\sin \theta - 1)^2} \quad (\text{Form of answer may vary.})$$

56. $f(x) = \left(\frac{x^2 - x - 3}{x^2 + 1}\right)(x^2 + x + 1)$

$$f'(x) = 2\frac{x^5 + 2x^3 + 2x^2 - 2}{(x^2 + 1)^2} \quad (\text{Form of answer may vary.})$$

59. $y = \frac{1 + \csc x}{1 - \csc x}$

$$y' = \frac{(1 - \csc x)(-\csc x \cot x) - (1 + \csc x)(\csc x \cot x)}{(1 - \csc x)^2} = \frac{-2 \csc x \cot x}{(1 - \csc x)^2}$$

$$y'\left(\frac{\pi}{6}\right) = \frac{-2(2)(\sqrt{3})}{(1 - 2)^2} = -4\sqrt{3}$$

60. $f(x) = \tan x \cot x = 1$

$$f''(x) = 0$$

$$f'(1) = 0$$

61. $h(t) = \frac{\sec t}{t}$

$$h'(t) = \frac{t(\sec t \tan t) - (\sec t)(1)}{t^2} = \frac{\sec t(t \tan t - 1)}{t^2}$$

$$h'(\pi) = \frac{\sec \pi(\pi \tan \pi - 1)}{\pi^2} = \frac{1}{\pi^2}$$

62. $f(x) = \sin x(\sin x + \cos x)$

$$f'(x) = \sin x(\cos x - \sin x) + (\sin x + \cos x)\cos x$$

$$= \sin x \cos x - \sin^2 x + \sin x \cos x + \cos^2 x$$

$$= \sin 2x + \cos 2x$$

$$f'\left(\frac{\pi}{4}\right) = \sin \frac{\pi}{2} + \cos \frac{\pi}{2} = 1$$

63. (a) $f(x) = (x^3 - 3x + 1)(x + 2), (1, -3)$


$$f'(x) = (x^3 - 3x + 1)(1) + (x + 2)(3x^2 - 3)$$

$$= 4x^3 + 6x^2 - 6x - 5$$

$$f'(1) = -1; \text{ Slope at } (1, -3)$$

Tangent line: $y + 3 = -1(x - 1) \Rightarrow y = -x - 2$

(b)




64. (a) $f(x) = (x - 1)(x^2 - 2), (0, 2)$

$$f'(x) = (x - 1)(2x) + (x^2 - 2)(1) = 3x^2 - 2x - 2$$

$$f'(0) = -2; \text{ Slope at } (0, 2)$$

Tangent line: $y - 2 = -2x \Rightarrow y = -2x + 2$

(b)




65. (a) $f(x) = \frac{x}{x - 1}, (2, 2)$

$$f'(x) = \frac{(x - 1)(1) - x(1)}{(x - 1)^2} = \frac{-1}{(x - 1)^2}$$

$$f'(2) = \frac{-1}{(2 - 1)^2} = -1; \text{ Slope at } (2, 2)$$

Tangent line: $y - 2 = -1(x - 2) \Rightarrow y = -x + 4$

(b)




66. (a) $f(x) = \frac{x-1}{x+1}$, $\left(2, \frac{1}{3}\right)$

$$f'(x) = \frac{(x+1)(1) - (x-1)(1)}{(x+1)^2} = \frac{2}{(x+1)^2}$$

$$f'(2) = \frac{2}{9}, \text{ Slope at } \left(2, \frac{1}{3}\right)$$

$$\text{Tangent line: } y - \frac{1}{3} = \frac{2}{9}(x-2) \Rightarrow y = \frac{2}{9}x - \frac{1}{9}$$



68. (a) $f(x) = \sec x$, $\left(\frac{\pi}{3}, 2\right)$

$$f'(x) = \sec x \tan x$$

$$f'\left(\frac{\pi}{3}\right) = 2\sqrt{3}, \text{ Slope at } \left(\frac{\pi}{3}, 2\right)$$

$$\text{Tangent line: } y - 2 = 2\sqrt{3}\left(x - \frac{\pi}{3}\right)$$

$$6\sqrt{3}x - 3y + 6 - 2\sqrt{3}\pi = 0$$

69. $f(x) = \frac{8}{x^2 + 4}$; $(2, 1)$

$$f'(x) = \frac{(x^2 + 4)(0) - 8(2x)}{(x^2 + 4)^2} = \frac{-16x}{(x^2 + 4)^2}$$

$$f'(2) = \frac{-16(2)}{(4+4)^2} = -\frac{1}{2}$$

$$y - 1 = -\frac{1}{2}(x - 2)$$

$$y = -\frac{1}{2}x + 2$$

$$2y + x - 4 = 0$$

71. $f(x) = \frac{16x}{x^2 + 16}$; $\left(-2, -\frac{8}{5}\right)$

$$f'(x) = \frac{(x^2 + 16)(16) - 16x(2x)}{(x^2 + 16)^2} = \frac{256 - 16x^2}{(x^2 + 16)^2}$$

$$f'(-2) = \frac{256 - 16(4)}{20^2} = \frac{12}{25}$$

$$y + \frac{8}{5} = \frac{12}{25}(x + 2)$$

$$y = \frac{12}{25}x - \frac{16}{25}$$

$$25y - 12x + 16 = 0$$

67. (a) $f(x) = \tan x$, $\left(\frac{\pi}{4}, 1\right)$

$$f'(x) = \sec^2 x$$



$$f'\left(\frac{\pi}{4}\right) = 2; \text{ Slope at } \left(\frac{\pi}{4}, 1\right)$$

Tangent line:

$$y - 1 = 2\left(x - \frac{\pi}{4}\right)$$

$$y - 1 = 2x - \frac{\pi}{2}$$

$$4x - 2y - \pi + 2 = 0$$



70. $f(x) = \frac{27}{x^2 + 9}$; $\left(-3, \frac{3}{2}\right)$

$$f'(x) = \frac{(x^2 + 9)(0) - 27(2x)}{(x^2 + 9)^2} = \frac{-54x}{(x^2 + 9)^2}$$

$$f'(-3) = \frac{-54(-3)}{(9+9)^2} = \frac{1}{2}$$

$$y - \frac{3}{2} = \frac{1}{2}(x + 3)$$

$$y = \frac{1}{2}x + 3$$

$$2y - x - 6 = 0$$

72. $f(x) = \frac{4x}{x^2 + 6}$; $\left(2, \frac{4}{5}\right)$

$$f'(x) = \frac{(x^2 + 6)(4) - 4x(2x)}{(x^2 + 6)^2} = \frac{24 - 4x^2}{(x^2 + 6)^2}$$

$$f'(2) = \frac{24 - 16}{10^2} = \frac{2}{25}$$

$$y - \frac{4}{5} = \frac{2}{25}(x - 2)$$

$$y = \frac{2}{25}x + \frac{16}{25}$$

$$25y - 2x - 16 = 0$$

73. $f(x) = \frac{x^2}{x-1}$

$$f'(x) = \frac{(x-1)(2x) - x^2(1)}{(x-1)^2}$$

$$= \frac{x^2 - 2x}{(x-1)^2} = \frac{x(x-2)}{(x-1)^2}$$

$f'(x) = 0$ when $x = 0$ or $x = 2$.

Horizontal tangents are at $(0, 0)$ and $(2, 4)$.

74. $f(x) = \frac{x^2}{x^2 + 1}$

$$f'(x) = \frac{(x^2 + 1)(2x) - (x^2)(2x)}{(x^2 + 1)^2}$$

$$= \frac{2x}{(x^2 + 1)^2}$$

$f'(x) = 0$ when $x = 0$.

Horizontal tangent is at $(0, 0)$.

75. $f(x) = \frac{4x-2}{x^2}$

$$f'(x) = \frac{x^2(4) - (4x-2)(2x)}{x^4}$$

$$= \frac{4x^2 - 8x^2 + 4x}{x^4}$$

$$= \frac{4 - 4x}{x^3}$$

$f'(x) = 0$ for $4 - 4x = 0 \Rightarrow x = 1$.

$$f(1) = 2$$

f has a horizontal tangent at $(1, 2)$.

76. $f(x) = \frac{x-4}{x^2-7}$

$$f'(x) = \frac{(x^2-7)(1) - (x-4)(2x)}{(x^2-7)^2} = \frac{x^2-7-2x^2+8x}{(x^2-7)^2} = -\frac{x^2-8x+7}{(x^2-7)^2} = -\frac{(x-7)(x-1)}{(x^2-7)^2}$$

$$f'(x) = 0 \text{ for } x = 1, 7; f(1) = \frac{1}{2}, f(7) = \frac{1}{14}$$

f has horizontal tangents at $\left(1, \frac{1}{2}\right)$ and $\left(7, \frac{1}{14}\right)$.

77. $f(x) = \frac{x+1}{x-1}$

$$f'(x) = \frac{(x-1) - (x+1)}{(x-1)^2} = \frac{-2}{(x-1)^2}$$

$$2y + x = 6 \Rightarrow y = -\frac{1}{2}x + 3; \text{ Slope: } -\frac{1}{2}$$

$$\frac{-2}{(x-1)^2} = -\frac{1}{2}$$


$$(x-1)^2 = 4$$

$$x-1 = \pm 2$$

$$x = -1, 3; f(-1) = 0, f(3) = 2$$

$$y - 0 = -\frac{1}{2}(x+1) \Rightarrow y = -\frac{1}{2}x - \frac{1}{2}$$

$$y - 2 = -\frac{1}{2}(x-3) \Rightarrow y = -\frac{1}{2}x + \frac{7}{2}$$



78. $f(x) = \frac{x}{x-1}$

$$f'(x) = \frac{(x-1) - x}{(x-1)^2} = \frac{-1}{(x-1)^2}$$

Let $(x, y) = (x, x/(x-1))$ be a point of tangency on the graph off.

$$\frac{5 - (x/(x-1))}{-1 - x} = \frac{-1}{(x-1)^2}$$

$$\frac{4x - 5}{(x-1)(x+1)} = \frac{1}{(x-1)^2}$$

$$(4x-5)(x-1) = x+1$$

$$4x^2 - 10x + 4 = 0$$


$$(x-2)(2x-1) = 0 \Rightarrow x = \frac{1}{2}, 2$$

$$f\left(\frac{1}{2}\right) = -1, f(2) = 2; f'\left(\frac{1}{2}\right) = -4, f'(2) = -1$$

Two tangent lines:

$$y + 1 = -4\left(x - \frac{1}{2}\right) \Rightarrow y = -4x + 1$$

$$y - 2 = -1(x-2) \Rightarrow y = -x + 4$$



$$79. f'(x) = \frac{(x+2)3 - 3x(1)}{(x+2)^2} = \frac{6}{(x+2)^2}$$

$$g'(x) = \frac{(x+2)5 - (5x+4)(1)}{(x+2)^2} = \frac{6}{(x+2)^2}$$

$$g(x) = \frac{5x+4}{(x+2)} = \frac{3x}{(x+2)} + \frac{2x+4}{(x+2)} = f(x) + 2$$

f and g differ by a constant.

$$81. (a) p'(x) = f'(x)g(x) + f(x)g'(x)$$

$$p'(1) = f'(1)g(1) + f(1)g'(1) = 1(4) + 6\left(-\frac{1}{2}\right) = 1$$

$$(b) q'(x) = \frac{g(x)f'(x) - f(x)g'(x)}{g(x)^2}$$

$$q'(4) = \frac{3(-1) - 7(0)}{3^2} = -\frac{1}{3}$$

$$83. \text{Area} = A(t) = (2t+1)\sqrt{t} = 2t^{3/2} + t^{1/2}$$

$$A'(t) = 2\left(\frac{3}{2}t^{1/2}\right) + \frac{1}{2}t^{-1/2}$$

$$= 3t^{1/2} + \frac{1}{2}t^{-1/2}$$

$$= \frac{6t+1}{2\sqrt{t}} \text{ cm}^2/\text{sec}$$

$$85. C = 100\left(\frac{200}{x^2} + \frac{x}{x+30}\right), \quad 1 \leq x$$

$$\frac{dC}{dx} = 100\left(-\frac{400}{x^3} + \frac{30}{(x+30)^2}\right)$$

$$(a) \text{ When } x = 10: \frac{dC}{dx} = -\$38.13$$

$$(b) \text{ When } x = 15: \frac{dC}{dx} = -\$10.37$$

$$(c) \text{ When } x = 20: \frac{dC}{dx} = -\$3.80$$

As the order size increases, the cost per item decreases.

$$87. P(t) = 500\left[1 + \frac{4t}{50+t^2}\right]$$

$$P'(t) = 500\left[\frac{(50+t^2)(4) - (4t)(2t)}{(50+t^2)^2}\right]$$

$$= 500\left[\frac{200-4t^2}{(50+t^2)^2}\right]$$

$$= 2000\left[\frac{50-t^2}{(50+t^2)^2}\right]$$

$$P'(2) \approx 31.55 \text{ bacteria per hour}$$

$$80. f'(x) = \frac{x(\cos x - 3) - (\sin x - 3x)(1)}{x^2} = \frac{x \cos x - \sin x}{x^2}$$

$$g'(x) = \frac{x(\cos x + 2) - (\sin x + 2x)(1)}{x^2} = \frac{x \cos x - \sin x}{x^2}$$

$$g(x) = \frac{\sin x + 2x}{x} = \frac{\sin x - 3x + 5x}{x} = f(x) + 5$$

f and g differ by a constant.

$$82. (a) p'(x) = f'(x)g(x) + f(x)g'(x)$$

$$p'(4) = \frac{1}{2}(8) + 1(0) = 4$$

$$(b) q'(x) = \frac{g(x)f'(x) - f(x)g'(x)}{g(x)^2}$$

$$q'(7) = \frac{4(2) - 4(-1)}{4^2} = \frac{12}{16} = \frac{3}{4}$$

$$84. V = \pi r^2 h = \pi(t+2)\left(\frac{1}{2}\sqrt{t}\right)$$

$$= \frac{1}{2}(t^{3/2} + 2t^{1/2})\pi$$

$$V'(t) = \frac{1}{2}\left(\frac{3}{2}t^{1/2} + t^{-1/2}\right)\pi = \frac{3t+2}{4t^{1/2}}\pi \text{ cubic inches/sec}$$

$$86. P = \frac{k}{V}$$

$$\frac{dP}{dV} = -\frac{k}{V^2}$$

$$88. F = \frac{Gm_1 m_2}{d^2} = Gm_1 m_2 d^{-2}$$

$$\frac{dF}{dd} = F'(d) = \frac{-2 Gm_1 m_2}{d^3}$$

89. (a) $\sec x = \frac{1}{\cos x}$

$$\frac{d}{dx}[\sec x] = \frac{d}{dx}\left[\frac{1}{\cos x}\right] = \frac{(\cos x)(0) - (1)(-\sin x)}{(\cos x)^2} = \frac{\sin x}{\cos x \cos x} = \frac{1}{\cos x} \cdot \frac{\sin x}{\cos x} = \sec x \tan x$$

(b) $\csc x = \frac{1}{\sin x}$

$$\frac{d}{dx}[\csc x] = \frac{d}{dx}\left[\frac{1}{\sin x}\right] = \frac{(\sin x)(0) - (1)(\cos x)}{(\sin x)^2} = -\frac{\cos x}{\sin x \sin x} = -\frac{1}{\sin x} \cdot \frac{\cos x}{\sin x} = -\csc x \cot x$$

(c) $\cot x = \frac{\cos x}{\sin x}$

$$\frac{d}{dx}[\cot x] = \frac{d}{dx}\left[\frac{\cos x}{\sin x}\right] = \frac{\sin x(-\sin x) - (\cos x)(\cos x)}{(\sin x)^2} = -\frac{\sin^2 x + \cos^2 x}{\sin^2 x} = -\frac{1}{\sin^2 x} = -\csc^2 x$$

90. $f(x) = \sec x$

$g(x) = \csc x, [0, 2\pi)$

$f'(x) = g'(x)$




$$\sec x \tan x = -\csc x \cot x \Rightarrow \frac{\sec x \tan x}{\csc x \cot x} = -1 \Rightarrow \frac{\frac{1}{\cos x} \cdot \frac{\sin x}{\cos x}}{\frac{1}{\sin x} \cdot \frac{\cos x}{\sin x}} = -1 \Rightarrow \frac{\sin^3 x}{\cos^3 x} = -1 \Rightarrow \tan^3 x = -1 \Rightarrow \tan x = -1$$

$$x = \frac{3\pi}{4}, \frac{7\pi}{4}$$

91. (a) $n(t) = -3.5806t^3 + 82.577t^2 - 603.60t + 1667.5$

$$v(t) = -0.1361t^3 + 3.165t^2 - 23.02t + 59.8$$

$$(c) A = \frac{v(t)}{n(t)} = \frac{-0.1361t^3 + 3.165t^2 - 23.02t + 59.8}{-3.5806t^3 + 82.577t^2 - 603.60t + 1667.5}$$



(d) $A'(t)$ represents the rate of change of the average retail value per 1000 motor homes.

A represents the average retail value (in millions of dollars) per 1000 motor homes.

92. (a) $\sin \theta = \frac{r}{r+h}$

$$r+h = r \csc \theta$$

$$h = r \csc \theta - r = r(\csc \theta - 1)$$

(b) $h'(\theta) = r(-\csc \theta \cdot \cot \theta)$

$$h'(30^\circ) = h'\left(\frac{\pi}{6}\right)$$

$$= -3960(2 \cdot \sqrt{3}) = -7920\sqrt{3} \text{ mi/radian}$$

93. $f(x) = 4x^{3/2}$

$$f'(x) = 6x^{1/2}$$

$$f''(x) = 3x^{-1/2} = \frac{3}{\sqrt{x}}$$

94. $f(x) = x + \frac{32}{x^2}$

$$f'(x) = 1 - \frac{64}{x^3}$$

$$f''(x) = \frac{192}{x^4}$$

95. $f(x) = \frac{x}{x-1}$

$$f'(x) = \frac{(x-1)(1) - x(1)}{(x-1)^2} = \frac{-1}{(x-1)^2}$$

$$f''(x) = \frac{2}{(x-1)^3}$$

97. $f(x) = 3 \sin x$

$$f'(x) = 3 \cos x$$

$$f''(x) = -3 \sin x$$

99. $f'(x) = x^2$

$$f''(x) = 2x$$

100. $f''(x) = 2 - 2x^{-1}$

$$f'''(x) = 2x^{-2} = \frac{2}{x^2}$$

96. $f(x) = \frac{x^2 + 2x - 1}{x} = x + 2 - \frac{1}{x}$

$$f'(x) = 1 + \frac{1}{x^2}$$

$$f''(x) = -\frac{2}{x^3}$$

98. $f(x) = \sec x$

$$f'(x) = \sec x \tan x$$


$$\begin{aligned} f''(x) &= \sec x (\sec^2 x) + \tan x (\sec x \tan x) \\ &= \sec x (\sec^2 x + \tan^2 x) \end{aligned}$$

101. $f'''(x) = 2\sqrt{x}$

$$\begin{aligned} f^{(4)}(x) &= \frac{1}{2}(2)x^{-1/2} = \frac{1}{\sqrt{x}} & f^{(5)}(x) &= 2 \\ f^{(6)}(x) &= 0 \end{aligned}$$

102. $f^{(4)}(x) = 2x + 1$


103.



$$f(2) = 0$$

One such function is $f(x) = (x-2)^2$.

104. The graph of a differentiable function f such that $f > 0$ and $f' < 0$ for all real numbers x would, in general, look like the graph below.



105. $f(x) = 2g(x) + h(x)$

$$f'(x) = 2g'(x) + h'(x)$$

$$f'(2) = 2g'(2) + h'(2)$$

$$= 2(-2) + 4$$

$$= 0$$

106. $f(x) = 4 - h(x)$

$$f'(x) = -h'(x)$$

$$f'(2) = -h'(2) = -4$$

107. $f(x) = \frac{g(x)}{h(x)}$

$$f'(x) = \frac{h(x)g'(x) - g(x)h'(x)}{[h(x)]^2}$$

$$f'(2) = \frac{h(2)g'(2) - g(2)h'(2)}{[h(2)]^2}$$

$$= \frac{(-1)(-2) - (3)(4)}{(-1)^2}$$

$$= -10$$

108. $f(x) = g(x)h(x)$


$$f'(x) = g(x)h'(x) + h(x)g'(x)$$

$$f'(2) = g(2)h'(2) + h(2)g'(2)$$

$$= (3)(4) + (-1)(-2)$$


$$= 14$$

109.




It appears that f is cubic; so f' would be quadratic and f'' would be linear.

110.




It appears that f is quadratic; so f'' would be linear and f' would be constant.


111.




112.



113.



114.



$$115. v(t) = 36 - t^2, \quad 0 \leq t \leq 6$$

$$a(t) = -2t$$

$$v(3) = 27 \text{ m/sec}$$

$$a(3) = -6 \text{ m/sec}^2$$

The speed of the object is decreasing.

$$116. v(t) = \frac{100t}{2t + 15}$$

$$a(t) = \frac{(2t + 15)(100) - (100t)(2)}{(2t + 15)^2}$$

$$= \frac{1500}{(2t + 15)^2}$$

$$(a) a(5) = \frac{1500}{[2(5) + 15]^2} = 2.4 \text{ ft/sec}^2$$

$$(b) a(10) = \frac{1500}{[2(10) + 15]^2} \approx 1.2 \text{ ft/sec}^2$$

$$(c) a(20) = \frac{1500}{[2(20) + 15]^2} \approx 0.5 \text{ ft/sec}^2$$

$$117. s(t) = -8.25t^2 + 66t$$

$$v(t) = -16.50t + 66$$

$$a(t) = -16.50$$

$t(\text{sec})$	0	1	2	3	4
$s(t)$ (ft)	0	57.75	99	123.75	132
$v(t) = s'(t)$ (ft/sec)	66	49.5	33	16.5	0
$a(t) = v'(t)$ (ft/sec 2)	-16.5	-16.5	-16.5	-16.5	-16.5

Average velocity on:


$$[0, 1] \text{ is } \frac{57.75 - 0}{1 - 0} = 57.75.$$

$$[1, 2] \text{ is } \frac{99 - 57.75}{2 - 1} = 41.25.$$

$$[2, 3] \text{ is } \frac{123.75 - 99}{3 - 2} = 24.75.$$

$$[3, 4] \text{ is } \frac{132 - 123.75}{4 - 3} = 8.25.$$

118. (a)

(b) The particle speeds up on the intervals $(\frac{4}{3}, \frac{8}{3})$ and $(4, 6)$.

Answers will vary.

s position function*v* velocity function*a* acceleration function

119. $f(x) = x^n$

$f^{(n)}(x) = n(n-1)(n-2)\cdots(2)(1) = n!$

Note: $n! = n(n-1)\cdots 3 \cdot 2 \cdot 1$ (read "n factorial")

120. $f(x) = \frac{1}{x}$

$$\begin{aligned}f^{(n)}(x) &= \frac{(-1)^n(n)(n-1)(n-2)\cdots(2)(1)}{x^{n+1}} \\&= \frac{(-1)^n n!}{x^{n+1}}\end{aligned}$$

121. $f(x) = g(x)h(x)$

(a) $f'(x) = g(x)h'(x) + h(x)g'(x)$

$f''(x) = g(x)h''(x) + g'(x)h'(x) + h(x)g''(x) + h'(x)g'(x)$

$= g(x)h''(x) + 2g'(x)h'(x) + h(x)g''(x)$

$f'''(x) = g(x)h'''(x) + g'(x)h''(x) + 2g''(x)h''(x) + 2g''(x)h'(x) + h(x)g'''(x) + h'(x)g''(x)$

$= g(x)h'''(x) + 3g'(x)h''(x) + 3g''(x)h'(x) + g'''(x)h(x)$

$$\begin{aligned}f^{(4)}(x) &= g(x)h^{(4)}(x) + g'(x)h'''(x) + 3g''(x)h'''(x) + 3g''(x)h''(x) + 3g'''(x)h''(x) \\&\quad + g'''(x)h'(x) + g^{(4)}(x)h(x)\end{aligned}$$

$= g(x)h^{(4)}(x) + 4g'(x)h'''(x) + 6g''(x)h''(x) + 4g'''(x)h'(x) + g^{(4)}(x)h(x)$

(b) $f^{(n)}(x) = g(x)h^{(n)}(x) + \frac{n(n-1)(n-2)\cdots(2)(1)}{1[(n-1)(n-2)\cdots(2)(1)]}g'(x)h^{(n-1)}(x) + \frac{n(n-1)(n-2)\cdots(2)(1)}{(2)(1)[(n-2)(n-3)\cdots(2)(1)]}g''(x)h^{(n-2)}(x)$

$+ \frac{n(n-1)(n-2)\cdots(2)(1)}{(3)(2)(1)[(n-3)(n-4)\cdots(2)(1)]}g'''(x)h^{(n-3)}(x) + \cdots$

$+ \frac{n(n-1)(n-2)\cdots(2)(1)}{[(n-1)(n-2)\cdots(2)(1)][1]}g^{(n-1)}(x)h'(x) + g^{(n)}(x)h(x)$

$= g(x)h^{(n)}(x) + \frac{n!}{1!(n-1)!}g'(x)h^{(n-1)}(x) + \frac{n!}{2!(n-2)!}g''(x)h^{(n-2)}(x) + \cdots$

$+ \frac{n!}{(n-1)!1!}g^{(n-1)}(x)h'(x) + g^{(n)}(x)h(x)$

Note: $n! = n(n-1)\cdots 3 \cdot 2 \cdot 1$ (read "n factorial")

122. $[xf(x)]' = xf'(x) + f(x)$

$[xf(x)]'' = xf''(x) + f'(x) + f'(x) = xf''(x) + 2f'(x)$

$[xf(x)]''' = xf'''(x) + f''(x) + 2f''(x) = xf'''(x) + 3f''(x)$

In general, $[xf(x)]^{(n)} = xf^{(n)}(x) + nf^{(n-1)}(x)$.

123. $f(x) = x^n \sin x$

$$\begin{aligned}f'(x) &= x^n \cos x + nx^{n-1} \sin x \\&= x^{n-1}(x \cos x + n \sin x)\end{aligned}$$

When $n = 1$: $f'(x) = x \cos x + \sin x$ When $n = 2$: $f'(x) = x(x \cos x + 2 \sin x)$ When $n = 3$: $f'(x) = x^2(x \cos x + 3 \sin x)$ When $n = 4$: $f'(x) = x^3(x \cos x + 4 \sin x)$ For general n , $f'(x) = x^{n-1}(x \cos x + n \sin x)$.

124. $f(x) = \frac{\cos x}{x^n} = x^{-n} \cos x$

$$\begin{aligned}f'(x) &= -x^{-n} \sin x - nx^{-n-1} \cos x \\&= -x^{-n-1}(x \sin x + n \cos x) \\&= -\frac{x \sin x + n \cos x}{x^{n+1}}\end{aligned}$$

When $n = 1$: $f'(x) = -\frac{x \sin x + \cos x}{x^2}$

When $n = 2$: $f'(x) = -\frac{x \sin x + 2 \cos x}{x^3}$

When $n = 3$: $f'(x) = -\frac{x \sin x + 3 \cos x}{x^4}$

When $n = 4$: $f'(x) = -\frac{x \sin x + 4 \cos x}{x^5}$

For general n , $f'(x) = -\frac{x \sin x + n \cos x}{x^{n+1}}$.

125. $y = \frac{1}{x}, y' = -\frac{1}{x^2}, y'' = \frac{2}{x^3}$

$x^3y'' + 2x^2y' = x^3\left[\frac{2}{x^3}\right] + 2x^2\left[-\frac{1}{x^2}\right] = 2 - 2 = 0$

126.

$y = 2x^3 - 6x + 10$

$y' = 6x^2 - 6$

$y'' = 12x$

$y''' = 12$

$-y''' - xy'' - 2y' = -12 - x(12x) - 2(6x^2 - 6) = -24x^2$

127. $y = 2 \sin x + 3$

$y' = 2 \cos x$

$y'' = -2 \sin x$

$y'' + y = -2 \sin x + (2 \sin x + 3) = 3$

128. $y = 3 \cos x + \sin x$

$y' = -3 \sin x + \cos x$

$y'' = -3 \cos x - \sin x$

$y'' + y = (-3 \cos x - \sin x) + (3 \cos x + \sin x) = 0$

129. False. If $y = f(x)g(x)$, then

$\frac{dy}{dx} = f(x)g'(x) + g(x)f'(x).$

130. True. y is a fourth-degree polynomial.

$\frac{d^n y}{dx^n} = 0 \text{ when } n > 4.$

131. True

$$\begin{aligned}h'(c) &= f(c)g'(c) + g(c)f'(c) \\&= f(c)(0) + g(c)(0) \\&= 0\end{aligned}$$

132. True

133. True

134. True.

If $v(t) = c$ then $a(t) = v'(t) = 0$.

135. $f(x) = ax^2 + bx + c$

$f'(x) = 2ax + b$

 x -intercept at $(1, 0)$: $0 = a + b + c$ $(2, 7)$ on graph: $7 = 4a + 2b + c$ Slope 10 at $(2, 7)$: $10 = 4a + b$ Subtracting the third equation from the second, $-3 = b + c$. Subtracting this equation from the first, $3 = a$. Then, $10 = 4(3) + b \Rightarrow b = -2$. Finally, $-3 = (-2) + c \Rightarrow c = -1$.

$f(x) = 3x^2 - 2x - 1$

136. $f(x) = ax^3 + bx^2 + cx + d, a \neq 0$

$$f'(x) = 3ax^2 + 2bx + c$$

$$f'(x) = 0 \Rightarrow x = \frac{-2b \pm \sqrt{4b^2 - 12ac}}{6a}$$

(a) No horizontal tangents: $f'(x) \neq 0$

$$4b^2 - 12ac < 0$$

Example: $a = c = 1, b = 0$:

$$f(x) = x^3 + x$$

(b) Exactly one horizontal tangent

$$4b^2 - 12ac = 0$$

Example: $a = 1, b = 3, c = 3$:

$$f(x) = x^3 + 3x^2 + 3x$$

(c) Exactly two horizontal tangents

$$4b^2 - 12ac > 0$$

Example: $b = 1, a = 1, c = 0$:

$$f(x) = x^3 + x^2$$

137. $f(x) = x|x| = \begin{cases} x^2, & \text{if } x \geq 0 \\ -x^2, & \text{if } x < 0 \end{cases}$

$$f''(x) = \begin{cases} 2x, & \text{if } x \geq 0 \\ -2x, & \text{if } x < 0 \end{cases} = 2|x|$$

$$f''(x) = \begin{cases} 2, & \text{if } x > 0 \\ -2, & \text{if } x < 0 \end{cases}$$

$f''(0)$ does not exist since the left and right derivatives are not equal.

138. (a) $(fg' - f'g)' = fg'' + f'g' - f'g' - f''g$

$$= fg'' - f''g \quad \text{True}$$

(b) $(fg)'' = (fg' + f'g)'$

$$= fg'' + f'g' + f'g' + f''g$$

$$= fg'' + 2f'g' + f''g$$

$$\neq fg'' + f''g \quad \text{False}$$

Section 2.4 The Chain Rule

$$y = f(g(x))$$

$$u = g(x)$$

$$y = f(u)$$

1. $y = (6x - 5)^4$

$$u = 6x - 5$$

$$y = u^4$$

2. $y = \frac{1}{\sqrt{x+1}}$

$$u = x + 1$$

$$y = u^{-1/2}$$

3. $y = \sqrt{x^2 - 1}$

$$u = x^2 - 1$$

$$y = \sqrt{u}$$

4. $y = 3 \tan(\pi x^2)$

$$u = \pi x^2$$

$$y = 3 \tan u$$

5. $y = \csc^3 x$

$$u = \csc x$$

$$y = u^3$$

6. $y = \cos \frac{3x}{2}$

$$u = \frac{3x}{2}$$

$$y = \cos u$$

7. $y = (2x - 7)^3$

$$y' = 3(2x - 7)^2(2) = 6(2x - 7)^2$$

8. $y = 3(4 - x^2)^5$

$$y' = 15(4 - x^2)^4(-2x) = -30x(4 - x^2)^4$$

9. $g(x) = 3(4 - 9x)^4$

$$g'(x) = 12(4 - 9x)^3(-9) = -108(4 - 9x)^3$$

10. $f(t) = (9t + 2)^{2/3}$

$$f'(t) = \frac{2}{3}(9t + 2)^{-1/3}(9) = \frac{6}{\sqrt[3]{9t + 2}}$$

11. $f(t) = (1 - t)^{1/2}$

$$f'(t) = \frac{1}{2}(1 - t)^{-1/2}(-1) = -\frac{1}{2\sqrt{1-t}}$$

13. $y = (9x^2 + 4)^{1/3}$

$$y' = \frac{1}{3}(9x^2 + 4)^{-2/3}(18x) = \frac{6x}{(9x^2 + 4)^{2/3}}$$

15. $y = 2(4 - x^2)^{1/4}$

$$y' = 2\left(\frac{1}{4}\right)(4 - x^2)^{-3/4}(-2x) = \frac{-x}{\sqrt[4]{(4 - x^2)^3}}$$

17. $y = (x - 2)^{-1}$

$$y' = -1(x - 2)^{-2}(1) = \frac{-1}{(x - 2)^2}$$

20. $y = -\frac{5}{(t + 3)^3}$

$$y = -5(t + 3)^{-3}$$

$$y' = 15(t + 3)^{-4} = \frac{15}{(t + 3)^4}$$

23. $f(x) = x^2(x - 2)^4$

$$\begin{aligned} f'(x) &= x^2[4(x - 2)^3(1)] + (x - 2)^4(2x) \\ &= 2x(x - 2)^3[2x + (x - 2)] \\ &= 2x(x - 2)^3(3x - 2) \end{aligned}$$

25. $y = x\sqrt{1 - x^2} = x(1 - x^2)^{1/2}$

$$\begin{aligned} y' &= x\left[\frac{1}{2}(1 - x^2)^{-1/2}(-2x)\right] + (1 - x^2)^{1/2}(1) \\ &= -x^2(1 - x^2)^{-1/2} + (1 - x^2)^{1/2} \\ &= (1 - x^2)^{-1/2}[-x^2 + (1 - x^2)] \\ &= \frac{1 - 2x^2}{\sqrt{1 - x^2}} \end{aligned}$$

12. $g(x) = \sqrt{5 - 3x} = (5 - 3x)^{1/2}$

$$g'(x) = \frac{1}{2}(5 - 3x)^{-1/2}(-3) = \frac{-3}{2\sqrt{5 - 3x}}$$

14. $g(x) = \sqrt{x^2 - 2x + 1} = \sqrt{(x - 1)^2} = |x - 1|$

$$g'(x) = \begin{cases} 1, & x > 1 \\ -1, & x < 1 \end{cases}$$

16. $f(x) = -3\sqrt[4]{2 - 9x}$

$$f(x) = -3(2 - 9x)^{1/4}$$

$$f'(x) = -\frac{3}{4}(2 - 9x)^{-3/4}(-9) = \frac{27}{4(2 - 9x)^{3/4}}$$

18. $s(t) = \frac{1}{t^2 + 3t - 1}$

$$s(t) = (t^2 + 3t - 1)^{-1}$$

$$s'(t) = -1(t^2 + 3t - 1)^{-2}(2t + 3)$$

$$= \frac{-(2t + 3)}{(t^2 + 3t - 1)^2}$$

19. $f(t) = (t - 3)^{-2}$

$$f'(t) = -2(t - 3)^{-3} = \frac{-2}{(t - 3)^3}$$

22. $g(t) = \sqrt{\frac{1}{t^2 - 2}}$

$$g(t) = (t^2 - 2)^{-1/2}$$

$$g'(t) = -\frac{1}{2}(t^2 - 2)^{-3/2}(2t)$$

$$= -\frac{t}{(t^2 - 2)^{3/2}}$$

24. $f(x) = x(3x - 9)^3$

$$\begin{aligned} f'(x) &= x[3(3x - 9)^2(3)] + (3x - 9)^3(1) \\ &= (3x - 9)^2[9x + 3x - 9] \\ &= 27(x - 3)^2(4x - 3) \end{aligned}$$

26. $y = \frac{1}{2}x^2\sqrt{16 - x^2}$

$$\begin{aligned} y' &= \frac{1}{2}x^2\left(\frac{1}{2}(16 - x^2)^{-1/2}(-2x)\right) + x(16 - x^2)^{1/2} \\ &= \frac{-x^3}{2\sqrt{16 - x^2}} + x\sqrt{16 - x^2} \\ &= \frac{-x(3x^2 - 32)}{2\sqrt{16 - x^2}} \end{aligned}$$

27. $y = \frac{x}{\sqrt{x^2 + 1}} = x(x^2 + 1)^{-1/2}$

$$\begin{aligned}y' &= x \left[-\frac{1}{2}(x^2 + 1)^{-3/2}(2x) \right] + (x^2 + 1)^{-1/2}(1) \\&= -x^2(x^2 + 1)^{-3/2} + (x^2 + 1)^{-1/2} \\&= (x^2 + 1)^{-3/2}[-x^2 + (x^2 + 1)] \\&= \frac{1}{(x^2 + 1)^{3/2}}\end{aligned}$$

29. $g(x) = \left(\frac{x+5}{x^2+2}\right)^2$

$$\begin{aligned}g'(x) &= 2\left(\frac{x+5}{x^2+2}\right)\left(\frac{(x^2+2)-(x+5)(2x)}{(x^2+2)^2}\right) \\&= \frac{2(x+5)(2-10x-x^2)}{(x^2+2)^3}\end{aligned}$$


31. $f(v) = \left(\frac{1-2v}{1+v}\right)^3$

$$\begin{aligned}f'(v) &= 3\left(\frac{1-2v}{1+v}\right)^2 \left(\frac{(1+v)(-2)-(1-2v)}{(1+v)^2}\right) \\&= \frac{-9(1-2v)^2}{(1+v)^4}\end{aligned}$$

33. $y = \frac{\sqrt{x}+1}{x^2+1}$

$$y' = \frac{1-3x^2-4x^{3/2}}{2\sqrt{x}(x^2+1)^2}$$

The zero of y' corresponds to the point on the graph of y where the tangent line is horizontal.



35. $y = \sqrt{\frac{x+1}{x}}$

$$y' = -\frac{\sqrt{(x+1)/x}}{2x(x+1)}$$

y' has no zeros.



28. $y = \frac{x}{\sqrt{x^4+4}}$

$$\begin{aligned}y' &= \frac{(x^4+4)^{1/2}(1)-x^1(x^4+4)^{-1/2}(4x^3)}{x^4+4} \\&= \frac{x^4+4-2x^4}{(x^4+4)^{3/2}} \\&= \frac{4-x^4}{(x^4+4)^{3/2}}\end{aligned}$$

30. $h(t) = \left(\frac{t^2}{t^3+2}\right)^2$

$$\begin{aligned}h'(t) &= 2\left(\frac{t^2}{t^3+2}\right)\left(\frac{(t^3+2)(2t)-t^2(3t^2)}{(t^3+2)^2}\right) \\&= \frac{2t^2(4t-t^4)}{(t^3+2)^3} = \frac{2t^3(4-t^3)}{(t^3+2)^3}\end{aligned}$$


32. $g(x) = \left(\frac{3x^2-2}{2x+3}\right)^3$

$$\begin{aligned}g'(x) &= 3\left(\frac{3x^2-2}{2x+3}\right)^2 \left(\frac{(2x+3)(6x)-(3x^2-2)(2)}{(2x+3)^2}\right) \\&= \frac{3(3x^2-2)^2(6x^2+18x+4)}{(2x+3)^4} \\&= \frac{6(3x^2-2)^2(3x^2+9x+2)}{(2x+3)^4}\end{aligned}$$

34. $y = \sqrt{\frac{2x}{x+1}}$

$$y' = \frac{1}{\sqrt{2x(x+1)^{3/2}}}$$


y' has no zeros.



36. $g(x) = \sqrt{x-1} + \sqrt{x+1}$


$$g'(x) = \frac{1}{2\sqrt{x-1}} + \frac{1}{2\sqrt{x+1}}$$

g' has no zeros.



37. $y = \frac{\cos \pi x + 1}{x}$

$$\begin{aligned} \frac{dy}{dx} &= \frac{-\pi x \sin \pi x - \cos \pi x - 1}{x^2} \\ &= -\frac{\pi x \sin \pi x + \cos \pi x + 1}{x^2} \end{aligned}$$



The zeros of y' correspond to the points on the graph of y where the tangent lines are horizontal.

39. (a) $y = \sin x$

$$y' = \cos x$$

$$y'(0) = 1$$

1 cycle in $[0, 2\pi]$

(b) $y = \sin 2x$

$$y' = 2 \cos 2x$$


$$y'(0) = 2$$

2 cycles in $[0, 2\pi]$

The slope of $\sin ax$ at the origin is a .

38. $y = x^2 \tan \frac{1}{x}$

$$\frac{dy}{dx} = 2x \tan \frac{1}{x} - \sec^2 \frac{1}{x}$$



The zeros of y' correspond to the points on the graph of y where the tangent lines are horizontal.

41. $y = \cos 3x$

$$\frac{dy}{dx} = -3 \sin 3x$$

42. $y = \sin \pi x$

$$\frac{dy}{dx} = \pi \cos \pi x$$

45. $y = \sin(\pi x)^2 = \sin(\pi^2 x^2)$

$$y' = \cos(\pi^2 x^2)[2\pi^2 x] = 2\pi^2 x \cos(\pi^2 x^2)$$

47. $h(x) = \sin 2x \cos 2x$

$$\begin{aligned} h'(x) &= \sin 2x(-2 \sin 2x) + \cos 2x(2 \cos 2x) \\ &= 2 \cos^2 2x - 2 \sin^2 2x \\ &= 2 \cos 4x \end{aligned}$$

Alternate solution: $h(x) = \frac{1}{2} \sin 4x$

$$h'(x) = \frac{1}{2} \cos 4x(4) = 2 \cos 4x$$

49. $f(x) = \frac{\cot x}{\sin x} = \frac{\cos x}{\sin^2 x}$

$$\begin{aligned} f'(x) &= \frac{\sin^2 x(-\sin x) - \cos x(2 \sin x \cos x)}{\sin^4 x} \\ &= \frac{-\sin^2 x - 2 \cos^2 x}{\sin^3 x} = \frac{-1 - \cos^2 x}{\sin^3 x} \end{aligned}$$

51. $y = 4 \sec^2 x$

$$\begin{aligned} y' &= 8 \sec x \cdot \sec x \tan x \\ &= 8 \sec^2 x \tan x \end{aligned}$$

52. $g(t) = 5 \cos^2 \pi t = 5(\cos \pi t)^2$

$$\begin{aligned} g'(t) &= 10 \cos \pi t(-\sin \pi t)(\pi) \\ &= -10\pi(\sin \pi t)(\cos \pi t) \\ &= -5\pi \sin 2\pi t \end{aligned}$$

50. $g(v) = \frac{\cos v}{\csc v} = \cos v \cdot \sin v$

$$\begin{aligned} g'(v) &= \cos v(\cos v) + \sin v(-\sin v) \\ &= \cos^2 v - \sin^2 v = \cos 2v \end{aligned}$$

53. $f(\theta) = \frac{1}{4} \sin^2 2\theta = \frac{1}{4}(\sin 2\theta)^2$

$$\begin{aligned} f'(\theta) &= 2\left(\frac{1}{4}\right)(\sin 2\theta)(\cos 2\theta)(2) \\ &= \sin 2\theta \cos 2\theta = \frac{1}{2} \sin 4\theta \end{aligned}$$

54. $h(t) = 2 \cot^2(\pi t + 2)$

$$\begin{aligned} h'(t) &= 4 \cot(\pi t + 2)(-\csc^2(\pi t + 2)(\pi)) \\ &= -4\pi \cot(\pi t + 2) \csc^2(\pi t + 2) \end{aligned}$$

56. $y = 3x - 5 \cos(\pi x)^2$

$= 3x - 5 \cos(\pi^2 x^2)$

$\frac{dy}{dx} = 3 + 5 \sin(\pi^2 x^2)(2\pi^2 x)$

$= 3 + 10\pi^2 x \sin(\pi x)^2$

58. $y = \sin x^{1/3} + (\sin x)^{1/3}$

$$\begin{aligned} y' &= \cos x^{1/3} \left(\frac{1}{3} x^{-2/3} \right) + \frac{1}{3} (\sin x)^{-2/3} \cos x \\ &= \frac{1}{3} \left[\frac{\cos x^{1/3}}{x^{2/3}} + \frac{\cos x}{(\sin x)^{2/3}} \right] \end{aligned}$$

60. $y = (3x^3 + 4x)^{1/5}, (2, 2)$

$$\begin{aligned} y' &= \frac{1}{5}(3x^3 + 4x)^{-4/5}(9x^2 + 4) \\ &= \frac{9x^2 + 4}{5(3x^3 + 4x)^{4/5}} \end{aligned}$$

$y'(2) = \frac{1}{2}$

62. $f(x) = \frac{1}{(x^2 - 3x)^2} = (x^2 - 3x)^{-2}, \left(4, \frac{1}{16}\right)$

$f'(x) = -2(x^2 - 3x)^{-3}(2x - 3) = \frac{-2(2x - 3)}{(x^2 - 3x)^3}$

$f'(4) = -\frac{5}{32}$

64. $f(x) = \frac{x+1}{2x-3}, (2, 3)$

$f'(x) = \frac{(2x-3)(1)-(x+1)(2)}{(2x-3)^2} = \frac{-5}{(2x-3)^2}$

$f'(2) = -5$

55. $f(t) = 3 \sec^2(\pi t - 1)$

$$\begin{aligned} f'(t) &= 6 \sec(\pi t - 1) \sec(\pi t - 1) \tan(\pi t - 1)(\pi) \\ &= 6\pi \sec^2(\pi t - 1) \tan(\pi t - 1) = \frac{6\pi \sin(\pi t - 1)}{\cos^3(\pi t - 1)} \end{aligned}$$

57. $y = \sqrt{x} + \frac{1}{4} \sin(2x)^2$

$= \sqrt{x} + \frac{1}{4} \sin(4x^2)$

$\frac{dy}{dx} = \frac{1}{2}x^{-1/2} + \frac{1}{4} \cos(4x^2)(8x)$

$= \frac{1}{2\sqrt{x}} + 2x \cos(2x)^2$

59. $s(t) = (t^2 + 2t + 8)^{1/2}, (2, 4)$

$s'(t) = \frac{1}{2}(t^2 + 2t + 8)^{-1/2}(2t + 2)$

$= \frac{t+1}{\sqrt{t^2 + 2t + 8}}$

$s'(2) = \frac{3}{4}$

61. $f(x) = \frac{3}{x^3 - 4} = 3(x^3 - 4)^{-1}, \left(-1, -\frac{3}{5}\right)$

$f'(x) = -3(x^3 - 4)^{-2}(3x^2) = -\frac{9x^2}{(x^3 - 4)^2}$

$f'(-1) = -\frac{9}{25}$

63. $f(t) = \frac{3t+2}{t-1}, (0, -2)$

$f'(t) = \frac{(t-1)(3) - (3t+2)(1)}{(t-1)^2} = \frac{-5}{(t-1)^2}$

$f'(0) = -5$

65. $y = 37 - \sec^3(2x), (0, 36)$

$y' = -3 \sec^2(2x)[2 \sec(2x) \tan(2x)]$

$= -6 \sec^3(2x) \tan(2x)$

$y'(0) = 0$

66. $y = \frac{1}{x} + \sqrt{\cos x}$, $\left(\frac{\pi}{2}, \frac{2}{\pi}\right)$

$$y' = -\frac{1}{x^2} - \frac{\sin x}{2\sqrt{\cos x}}$$

$y''(\pi/2)$ is undefined.

67. (a) $f(x) = \sqrt{3x^2 - 2}$, $(3, 5)$


$$f'(x) = \frac{1}{2}(3x^2 - 2)^{-1/2}(6x) = \frac{3x}{\sqrt{3x^2 - 2}}$$

$$f'(3) = \frac{9}{5}$$

Tangent line:

$$y - 5 = \frac{9}{5}(x - 3) \Rightarrow 9x - 5y - 2 = 0$$

(b)




68. (a) $f(x) = \frac{1}{3}x\sqrt{x^2 + 5}$, $(2, 2)$

$$\begin{aligned} f'(x) &= \frac{1}{3}x\left[\frac{1}{2}(x^2 + 5)^{-1/2}(2x)\right] + \frac{1}{3}(x^2 + 5)^{1/2} \\ &= \frac{x^2}{3\sqrt{x^2 + 5}} + \frac{1}{3}\sqrt{x^2 + 5} \end{aligned}$$

$$f'(2) = \frac{4}{3(3)} + \frac{1}{3}(3) = \frac{13}{9}$$

$$\text{Tangent line: } y - 2 = \frac{13}{9}(x - 2) \Rightarrow 13x - 9y - 8 = 0$$

(b)



69. (a) $y = (2x^3 + 1)^2$, $(-1, 1)$


$$y' = 2(2x^3 + 1)(6x^2) = 12x^2(2x^3 + 1)$$

$$y'(-1) = 12(-1) = -12$$

$$\text{Tangent line: } y - 1 = -12(x + 1)$$

$$y = -12x - 11$$

(b)



70. (a) $f(x) = (9 - x^2)^{2/3}$, $(1, 4)$


$$f'(x) = \frac{2}{3}(9 - x^2)^{-1/3}(-2x) = \frac{-4x}{3(9 - x^2)^{1/3}}$$

$$f'(1) = \frac{-4}{3(8)^{1/3}} = -\frac{2}{3}$$

$$\text{Tangent line: } y - 4 = -\frac{2}{3}(x - 1)$$

$$y = -\frac{2}{3}x + \frac{14}{3}$$

(b)



71. (a) $f(x) = \sin 2x$, $(\pi, 0)$


$$f'(x) = 2 \cos 2x$$

$$f'(\pi) = 2$$

Tangent line:

$$y = 2(x - \pi) \Rightarrow 2x - y - 2\pi = 0$$

(b)




72. (a) $y = \cos 3x, \left(\frac{\pi}{4}, -\frac{\sqrt{2}}{2}\right)$

$$y' = -3 \sin 3x$$

$$y'\left(\frac{\pi}{4}\right) = -3 \sin\left(\frac{3\pi}{4}\right) = \frac{-3\sqrt{2}}{2}$$

$$\text{Tangent line: } y + \frac{\sqrt{2}}{2} = \frac{-3\sqrt{2}}{2}\left(x - \frac{\pi}{4}\right)$$

$$y = \frac{-3\sqrt{2}}{2}x + \frac{3\sqrt{2}\pi}{8} - \frac{\sqrt{2}}{2}$$




74. (a) $y = 2 \tan^3 x, \left(\frac{\pi}{4}, 2\right)$

$$y' = 6 \tan^2 x \cdot \sec^2 x$$

$$y'\left(\frac{\pi}{4}\right) = 6(1)(2) = 12$$

$$\text{Tangent line: } y - 2 = 12\left(x - \frac{\pi}{4}\right)$$

$$y = 12x + 2 - 3\pi$$




76. (a) $f(x) = \sqrt{x}(2-x)^2, (4, 8)$

$$f'(x) = \frac{(x-2)(5x-2)}{2\sqrt{x}}$$

$$f'(4) = 9$$

(b) $y - 8 = 9(x - 4)$

$$y = 9x - 28$$




73. (a) $f(x) = \tan^2 x, \left(\frac{\pi}{4}, 1\right)$

$$f'(x) = 2 \tan x \sec^2 x$$

$$f'\left(\frac{\pi}{4}\right) = 2(1)(2) = 4$$

Tangent line:

$$y - 1 = 4\left(x - \frac{\pi}{4}\right) \Rightarrow 4x - y + (1 - \pi) = 0$$




75. (a) $g(t) = \frac{3t^2}{\sqrt{t^2 + 2t - 1}}, \left(\frac{1}{2}, \frac{3}{2}\right)$

$$g'(t) = \frac{3t(t^2 + 3t - 2)}{(t^2 + 2t - 1)^{3/2}}$$

$$g'\left(\frac{1}{2}\right) = -3$$

(b) $y - \frac{3}{2} = -3\left(x - \frac{1}{2}\right)$

$$y = -3x + 3$$




77. (a) $s(t) = \frac{(4-2t)\sqrt{1+t}}{3}, \left(0, \frac{4}{3}\right)$

$$s'(t) = \frac{-2\sqrt{1+t}}{3} + \frac{2-t}{3\sqrt{1+t}}$$

$$s'(0) = 0$$

(b) $y - \frac{4}{3} = 0(x - 0)$

$$y = \frac{4}{3}$$




78. (a) $y = (t^2 - 9)\sqrt{t+2}$, $(2, -10)$

$$y' = \frac{5t^2 + 8t - 9}{2\sqrt{t+2}}$$

$$y'(2) = \frac{27}{4}$$

(b) $y + 10 = \frac{27}{4}(t - 2)$

$$y = \frac{27}{4}t - \frac{47}{2}$$




80. $f(x) = \frac{|x|}{\sqrt{2-x^2}}$, $(1, 1)$

$$f'(x) = \frac{2}{(2-x^2)^{3/2}} \text{ for } x > 0$$

$$f'(1) = 2$$

$$y - 1 = 2(x - 1)$$

$$y = 2x - 1; \text{ Tangent line}$$



82. $f(x) = \frac{x}{\sqrt{2x-1}}$

$$f'(x) = \frac{(2x-1)^{1/2} - x(2x-1)^{-1/2}}{2x-1}$$

$$= \frac{2x-1-x}{(2x-1)^{3/2}}$$

$$= \frac{x-1}{(2x-1)^{3/2}}$$

$$\frac{x-1}{(2x-1)^{3/2}} = 0 \Rightarrow x = 1$$

Horizontal tangent at $(1, 1)$


79. $f(x) = \sqrt{25-x^2}$, $(3, 4)$

$$f'(x) = \frac{-x}{\sqrt{25-x^2}}$$

$$f'(3) = -\frac{3}{4}$$

$$y - 4 = -\frac{3}{4}(x - 3)$$

$$y = -\frac{3}{4}x + \frac{25}{4}, \text{ Tangent line}$$



81. $f(x) = 2 \cos x + \sin 2x$, $0 < x < 2\pi$

$$f'(x) = -2 \sin x + 2 \cos 2x$$

$$= -2 \sin x + 2 - 4 \sin^2 x = 0$$

$$2 \sin^2 x + \sin x - 1 = 0$$

$$(\sin x + 1)(2 \sin x - 1) = 0$$

$$\sin x = -1 \Rightarrow x = \frac{3\pi}{2}$$

$$\sin x = \frac{1}{2} \Rightarrow x = \frac{\pi}{6}, \frac{5\pi}{6}$$

Horizontal tangents at $x = \frac{\pi}{6}, \frac{3\pi}{2}, \frac{5\pi}{6}$

Horizontal tangent at the points $\left(\frac{\pi}{6}, \frac{3\sqrt{3}}{2}\right)$, $\left(\frac{3\pi}{2}, 0\right)$, and $\left(\frac{5\pi}{6}, -\frac{3\sqrt{3}}{2}\right)$

83. $f(x) = 2(x^2 - 1)^3$

$$f'(x) = 6(x^2 - 1)^2(2x)$$

$$= 12x(x^4 - 2x^2 + 1)$$

$$= 12x^5 - 24x^3 + 12x$$

$$f''(x) = 60x^4 - 72x^2 + 12$$

$$= 12(5x^2 - 1)(x^2 - 1)$$

84. $f(x) = (x - 2)^{-1}$

$$f'(x) = -(x - 2)^{-2} = \frac{-1}{(x - 2)^2}$$

$$f''(x) = 2(x - 2)^{-3} = \frac{2}{(x - 2)^3}$$

86. $f(x) = \sec^2 \pi x$

$$\begin{aligned} f'(x) &= 2 \sec \pi x (\pi \sec \pi x \tan \pi x) \\ &= 2\pi \sec^2 \pi x \tan \pi x \end{aligned}$$

$$\begin{aligned} f''(x) &= 2\pi \sec^2 \pi x (\sec^2 \pi x)(\pi) + 2\pi \tan \pi x (2\pi \sec^2 \pi x \tan \pi x) \\ &= 2\pi^2 \sec^4 \pi x + 4\pi^2 \sec^2 \pi x \tan^2 \pi x \\ &= 2\pi^2 \sec^2 \pi x (\sec^2 \pi x + 2 \tan^2 \pi x) \\ &= 2\pi^2 \sec^2 \pi x (3 \sec^2 \pi x - 2) \end{aligned}$$

87. $h(x) = \frac{1}{9}(3x + 1)^3, \quad \left(1, \frac{64}{9}\right)$

$$h'(x) = \frac{1}{9}3(3x + 1)^2(3) = (3x + 1)^2$$

$$h''(x) = 2(3x + 1)(3) = 6(3x + 1)$$

$$h''(1) = 24$$

85. $f(x) = \sin x^2$

$$f'(x) = 2x \cos x^2$$

$$\begin{aligned} f''(x) &= 2x[2x(-\sin x^2)] + 2 \cos x^2 \\ &= 2[\cos x^2 - 2x^2 \sin x^2] \end{aligned}$$

88.

$$f(x) = \frac{1}{\sqrt{x+4}} = (x+4)^{-1/2}, \quad \left(0, \frac{1}{2}\right)$$

$$f'(x) = -\frac{1}{2}(x+4)^{-3/2}$$

$$f''(x) = \frac{3}{4}(x+4)^{-5/2} = \frac{3}{4(x+4)^{5/2}}$$

$$f''(0) = \frac{3}{128}$$

89. $f(x) = \cos(x^2), \quad (0, 1)$

$$f'(x) = -\sin(x^2)(2x) = -2x \sin(x^2)$$

$$\begin{aligned} f''(x) &= -2x \cos(x^2)(2x) - 2 \sin(x^2) \\ &= -4x^2 \cos(x^2) - 2 \sin(x^2) \end{aligned}$$

$$f''(0) = 0$$


90. $g(t) = \tan(2t), \quad \left(\frac{\pi}{6}, \sqrt{3}\right)$

$$g'(t) = 2 \sec^2(2t)$$

$$\begin{aligned} g''(t) &= 4 \sec(2t) \cdot \sec(2t) \tan(2t) 2 \\ &= 8 \sec^2(2t) \tan(2t) \end{aligned}$$


$$g''\left(\frac{\pi}{6}\right) = 32\sqrt{3}$$

91.




The zeros of f' correspond to the points where the graph of f has horizontal tangents.

92.




f is decreasing on $(-\infty, -1)$ so f' must be negative there.
 f is increasing on $(1, \infty)$ so f' must be positive there.

93.



The zeros of f' correspond to the points where the graph of f has horizontal tangents.

94.



The zeros of f' correspond to the points where the graph of f has horizontal tangents.

95. $g(x) = f(3x)$

$$g'(x) = f'(3x)(3) \Rightarrow g'(x) = 3f'(3x)$$

97. (a) $f(x) = g(x)h(x)$

$$f'(x) = g(x)h'(x) + g'(x)h(x)$$

$$f'(5) = (-3)(-2) + (6)(3) = 24$$

(c) $f(x) = \frac{g(x)}{h(x)}$

$$f'(x) = \frac{h(x)g'(x) - g(x)h'(x)}{[h(x)]^2}$$

$$f'(5) = \frac{(3)(6) - (-3)(-2)}{(3)^2} = \frac{12}{9} = \frac{4}{3}$$

98. (a) $g(x) = f(x) - 2 \Rightarrow g'(x) = f'(x)$

(b) $h(x) = 2f(x) \Rightarrow h'(x) = 2f'(x)$

(c) $r(x) = f(-3x) \Rightarrow r'(x) = f'(-3x)(-3) = -3f'(-3x)$

Hence, you need to know $f'(-3x)$.

$$r'(0) = -3f'(0) = (-3)\left(-\frac{1}{3}\right) = 1$$

$$r'(-1) = -3f'(3) = (-3)(-4) = 12$$

(d) $s(x) = f(x+2) \Rightarrow s'(x) = f'(x+2)$

Hence, you need to know $f'(x+2)$.

$$s'(-2) = f'(0) = -\frac{1}{3}, \text{ etc.}$$

99. (a) $h(x) = f(g(x)), g(1) = 4, g'(1) = -\frac{1}{2}, f'(4) = -1, h'(x) = f'(g(x))g'(x)$

$$h'(1) = f'(g(1))g'(1)$$

$$= f'(4)g'(1)$$

$$= (-1)\left(-\frac{1}{2}\right) = \frac{1}{2}$$

(b) $s(x) = g(f(x)), f(5) = 6, f'(5) = -1, g'(6)$ does not exist.

$$s'(x) = g'(f(x))f'(x)$$

$$s'(5) = g'(f(5))f'(5) = g'(6)(-1)$$

Since $g'(6)$ does not exist, $s'(5)$ is not defined.

96. $g(x) = f(x^2)$

$$g'(x) = f'(x^2)(2x) \Rightarrow g'(x) = 2xf'(x^2)$$

(b) $f(x) = g(h(x))$

$$f'(x) = g'(h(x))h'(x)$$

$$f'(5) = g'(3)(-2) = -2g'(3)$$

Need $g'(3)$ to find $f'(5)$.

(d) $f(x) = [g(x)]^3$

$$f'(x) = 3[g(x)]^2g'(x)$$

$$f'(5) = 3(-3)^2(6) = 162$$

x	-2	-1	0	1	2	3
$f'(x)$	4	$\frac{2}{3}$	$-\frac{1}{3}$	-1	-2	-4
$g'(x)$	4	$\frac{2}{3}$	$-\frac{1}{3}$	-1	-2	-4
$h'(x)$	8	$\frac{4}{3}$	$-\frac{2}{3}$	-2	-4	-8
$r'(x)$		12	1			
$s'(x)$	$-\frac{1}{3}$	-1	-2	-4		

100. (a) $h(x) = f(g(x))$
 $h'(x) = f'(g(x))g'(x)$
 $h'(3) = f'(g(3))g'(3)$
 $= f'(5)(1)$
 $= \frac{1}{2}$

(b) $s(x) = g(f(x))$
 $s'(x) = g'(f(x))f'(x)$
 $s'(9) = g'(f(9))f'(9)$
 $= g'(8)(2)$
 $= (-1)(2)$
 $= -2$

101. (a) $F = 132,400(331 - v)^{-1}$
 $F' = (-1)(132,400)(331 - v)^{-2}(-1)$
 $= \frac{132,400}{(331 - v)^2}$

When $v = 30, F' \approx 1.461$.

(b) $F = 132,400(331 + v)^{-1}$
 $F' = (-1)(132,400)(331 + v)^{-2}(1)$
 $= \frac{-132,400}{(331 + v)^2}$

When $v = 30, F' \approx -1.016$.

102. $y = \frac{1}{3} \cos 12t - \frac{1}{4} \sin 12t$

$$\begin{aligned}v &= y' = \frac{1}{3}[-12 \sin 12t] - \frac{1}{4}[12 \cos 12t] \\&= -4 \sin 12t - 3 \cos 12t\end{aligned}$$

When $t = \pi/8$, $y = 0.25$ feet and $v = 4$ feet per second.

104. $y = A \cos \omega t$

(a) Amplitude: $A = \frac{3.5}{2} = 1.75$

$$y = 1.75 \cos \omega t$$


Period: $10 \Rightarrow \omega = \frac{2\pi}{10} = \frac{\pi}{5}$

$$y = 1.75 \cos \frac{\pi t}{5}$$

(b) $v = y' = 1.75 \left[-\frac{\pi}{5} \sin \frac{\pi t}{5} \right] = -0.35\pi \sin \frac{\pi t}{5}$

106. (a) Using a graphing utility, or trial and error, you obtain a model similar to

$$T(t) = 65 + 21 \sin \left(\frac{\pi t}{6} - 2.1 \right).$$



103. $\theta = 0.2 \cos 8t$

The maximum angular displacement is $\theta = 0.2$ (since $-1 \leq \cos 8t \leq 1$).

$$\frac{d\theta}{dt} = 0.2[-8 \sin 8t] = -1.6 \sin 8t$$

When $t = 3$, $d\theta/dt = -1.6 \sin 24 \approx 1.4489$ radians per second.


105. $S = C(R^2 - r^2)$

$$\frac{dS}{dt} = C \left(2R \frac{dR}{dt} - 2r \frac{dr}{dt} \right)$$

Since r is constant, we have $dr/dt = 0$ and

$$\begin{aligned}\frac{dS}{dt} &= (1.76 \times 10^5)(2)(1.2 \times 10^{-2})(10^{-5}) \\&= 4.224 \times 10^{-2} = 0.04224.\end{aligned}$$

(c) $T'(t) \approx 11 \cos \left(\frac{\pi t}{6} - 2.1 \right)$



(d) The temperature changes most rapidly in the spring (April–June) and fall (September–November).

107. (a) $x = -1.6372t^3 + 19.3120t^2 - 0.5082t - 0.6162$

(b) $C = 60x + 1350$

$$= 60(-1.6372t^3 + 19.3120t^2 - 0.5082t - 0.6162) + 1350$$

$$\frac{dC}{dt} = 60(-4.9116t^2 + 38.624t - 0.5082)$$

$$= -294.696t^2 + 2317.44t - 30.492$$

The function dC/dt is quadratic, not linear. The cost function levels off at the end of the day, perhaps due to fatigue.

108. $f(x) = \sin \beta x$

(a) $f'(x) = \beta \cos \beta x$

$$f''(x) = -\beta^2 \sin \beta x$$

$$f'''(x) = -\beta^3 \cos \beta x$$

$$f^{(4)} = \beta^4 \sin \beta x$$

(b) $f''(x) + \beta^2 f(x) = -\beta^2 \sin \beta x + \beta^2(\sin \beta x) = 0$

(c) $f^{(2k)}(x) = (-1)^k \beta^{2k} \sin \beta x$

$$f^{(2k-1)}(x) = (-1)^{k+1} \beta^{2k-1} \cos \beta x$$

110. (a) $r'(x) = f'(g(x))g'(x)$

$$r'(1) = f'(g(1))g'(1)$$

Note that $g(1) = 4$ and $f'(4) = \frac{5-0}{6-2} = \frac{5}{4}$.

Also, $g'(1) = 0$. Thus, $r'(1) = 0$.

111. (a) $g(x) = \sin^2 x + \cos^2 x = 1 \Rightarrow g'(x) = 0$

$$g'(x) = 2 \sin x \cos x + 2 \cos x(-\sin x) = 0$$

109. $f(x+p) = f(x)$ for all x .

(a) Yes, $f'(x+p) = f'(x)$, which shows that f' is periodic as well.

(b) Yes, let $g(x) = f(2x)$, so $g'(x) = 2f'(2x)$. Since f' is periodic, so is g' .

(b) $s'(x) = g'(f(x))f'(x)$

$$s'(4) = g'(f(4))f'(4)$$

Note that $f(4) = \frac{5}{2}$, $g'(\frac{5}{2}) = \frac{6-4}{6-2} = \frac{1}{2}$ and

$$f'(4) = \frac{5}{4}. \text{ Thus, } s'(4) = \frac{1}{2}(\frac{5}{4}) = \frac{5}{8}.$$

(b) $\tan^2 x + 1 = \sec^2 x$

$$g(x) + 1 = f(x)$$

Taking derivatives of both sides, $g'(x) = f'(x)$. Equivalently, $f'(x) = 2 \sec x \cdot \sec x \cdot \tan x$ and $g'(x) = 2 \tan x \cdot \sec^2 x$, which are the same.

112. (a) If $f(-x) = -f(x)$, then

$$\frac{d}{dx}[f(-x)] = \frac{d}{dx}[-f(x)]$$

$$f'(-x)(-1) = -f'(x)$$

$$f'(-x) = f'(x).$$

Thus, $f'(x)$ is even.

(b) If $f(-x) = f(x)$, then

$$\frac{d}{dx}[f(-x)] = \frac{d}{dx}[f(x)]$$

$$f'(-x)(-1) = f'(x)$$

$$f'(-x) = -f'(x).$$

Thus, f' is odd.

113. $|u| = \sqrt{u^2}$

$$\frac{d}{dx}[|u|] = \frac{d}{dx}[\sqrt{u^2}] = \frac{1}{2}(u^2)^{-1/2}(2uu')$$

$$= \frac{uu'}{\sqrt{u^2}} = u' \frac{u}{|u|}, \quad u \neq 0$$

114. $g(x) = |2x - 3|$

$$g'(x) = 2 \left(\frac{2x-3}{|2x-3|} \right), \quad x \neq \frac{3}{2}$$

115. $f(x) = |x^2 - 4|$

$$f'(x) = 2x \left(\frac{x^2 - 4}{|x^2 - 4|} \right), \quad x \neq \pm 2$$

116. $h(x) = |x| \cos x$

$$h'(x) = -|x| \sin x + \frac{x}{|x|} \cos x, \quad x \neq 0$$

117. $f(x) = |\sin x|$

$$f'(x) = \cos x \left(\frac{\sin x}{|\sin x|} \right), \quad x \neq k\pi$$

118. (a) $f(x) = \tan \frac{\pi x}{4}$

$$f(1) = 1$$

$$f'(x) = \frac{\pi}{4} \sec^2 \frac{\pi x}{4}$$

$$f'(1) = \frac{\pi}{4}(2) = \frac{\pi}{2}$$


$$f''(x) = \frac{\pi}{2} \sec^2 \frac{\pi x}{4} \cdot \tan \frac{\pi x}{4} \left(\frac{\pi}{4} \right)$$

$$f''(1) = \frac{\pi^2}{8}(2)(1) = \frac{\pi^2}{4}$$

$$P_1(x) = f'(1)(x - 1) + f(1) = \frac{\pi}{2}(x - 1) + 1$$

$$P_2(x) = \frac{1}{2} \left(\frac{\pi^2}{4} \right) (x - 1)^2 + f'(1)(x - 1) + f(1) = \frac{\pi^2}{8}(x - 1)^2 + \frac{\pi}{2}(x - 1) + 1$$

(b)

(c) P_2 is a better approximation than P_1 .(d) The accuracy worsens as you move away from $x = c = 1$.

119. (a) $f(x) = \sec(2x)$

$$f'(x) = 2(\sec 2x)(\tan 2x)$$

$$\begin{aligned} f''(x) &= 2[2(\sec 2x)(\tan 2x)] \tan 2x + 2(\sec 2x)(\sec^2 2x)(2) \\ &= 4[(\sec 2x)(\tan^2 2x) + \sec^3 2x] \end{aligned}$$

$$f'\left(\frac{\pi}{6}\right) = \sec\left(\frac{\pi}{3}\right) = 2$$

$$f'\left(\frac{\pi}{6}\right) = 2 \sec\left(\frac{\pi}{3}\right) \tan\left(\frac{\pi}{3}\right) = 4\sqrt{3}$$


$$f''\left(\frac{\pi}{6}\right) = 4[2(3) + 2^3] = 56$$

$$P_1(x) = 4\sqrt{3}\left(x - \frac{\pi}{6}\right) + 2$$

$$P_2(x) = \frac{1}{2}(56)\left(x - \frac{\pi}{6}\right)^2 + 4\sqrt{3}\left(x - \frac{\pi}{6}\right) + 2$$

$$= 28\left(x - \frac{\pi}{6}\right)^2 + 4\sqrt{3}\left(x - \frac{\pi}{6}\right) + 2$$

(b)

(c) P_2 is a better approximation than P_1 .(d) The accuracy worsens as you move away from $x = \pi/6$.120. False. If $y = (1 - x)^{1/2}$, then

$$y' = \frac{1}{2}(1 - x)^{-1/2}(-1).$$

121. False. If $f(x) = \sin^2 2x$, then

$$f'(x) = 2(\sin 2x)(2 \cos 2x).$$

122. True

123. $f(x) = a_1 \sin x + a_2 \sin 2x + \cdots + a_n \sin nx$
 $f'(x) = a_1 \cos x + 2a_2 \cos 2x + \cdots + na_n \cos nx$
 $f'(0) = a_1 + 2a_2 + \cdots + na_n$
 $|a_1 + 2a_2 + \cdots + na_n| = |f'(0)|$

$$\begin{aligned} &= \lim_{x \rightarrow 0} \left| \frac{f(x) - f(0)}{x - 0} \right| \\ &= \lim_{x \rightarrow 0} \left| \frac{f(x)}{\sin x} \right| \cdot \left| \frac{\sin x}{x} \right| \\ &= \lim_{x \rightarrow 0} \left| \frac{f(x)}{\sin x} \right| \leq 1 \end{aligned}$$

124. $\frac{d}{dx} \left[\frac{P_n(x)}{(x^k - 1)^{n+1}} \right] = \frac{(x^k - 1)^{n+1} P_n'(x) - P_n(x)(n+1)(x^k - 1)^n kx^{k-1}}{(x^k - 1)^{2n+2}}$
 $= \frac{(x^k - 1)P_n'(x) - (n+1)kx^{k-1}P_n(x)}{(x^k - 1)^{n+2}}$

$P_n(x) = (x^k - 1)^{n+1} \frac{d^n}{dx^n} \left[\frac{1}{x^k - 1} \right] \Rightarrow$

$P_{n+1}(x) = (x^k - 1)^{n+2} \frac{d}{dx} \left[\frac{d^n}{dx^n} \left[\frac{1}{x^k - 1} \right] \right] = (x^k - 1)P_n'(x) - (n+1)kx^{k-1}P_n(x)$

$P_{n+1}(1) = -(n+1)kP_n(1)$

$\text{For } n = 1, \frac{d}{dx} \left[\frac{1}{x^k - 1} \right] = \frac{-kx^{k-1}}{(x^k - 1)^2} = \frac{P_1(x)}{(x^k - 1)^2} \Rightarrow P_1(1) = -k. \text{ Also, } P_0(1) = 1.$

We now use mathematical induction to verify that $P_n(1) = (-k)^n n!$ for $n \geq 0$. Assume true for n . Then

$$\begin{aligned} P_{n+1}(1) &= -(n+1)k(-k)^n n! \\ &= -(n+1)k(-k)^n n! \\ &= (-k)^{n+1}(n+1)!. \end{aligned}$$

Section 2.5 Implicit Differentiation

1. $x^2 + y^2 = 36$

$2x + 2yy' = 0$

$y' = \frac{-x}{y}$

2. $x^2 - y^2 = 16$

$2x - 2yy' = 0$

$y' = \frac{x}{y}$

3. $x^{1/2} + y^{1/2} = 9$

$\frac{1}{2}x^{-1/2} + \frac{1}{2}y^{-1/2}y' = 0$

$$\begin{aligned} y' &= -\frac{x^{-1/2}}{y^{-1/2}} \\ &= -\sqrt{\frac{y}{x}} \end{aligned}$$

4. $x^3 + y^3 = 8$

$3x^2 + 3y^2y' = 0$

$y' = -\frac{x^2}{y^2}$

5. $x^3 - xy + y^2 = 4$

$3x^2 - xy' - y + 2yy' = 0$

$(2y - x)y' = y - 3x^2$

$y' = \frac{y - 3x^2}{2y - x}$

6. $x^2y + y^2x = -2$

$x^2y' + 2xy + y^2 + 2xyy' = 0$

$(x^2 + 2xy)y' = -(y^2 + 2xy)$

$y' = \frac{-y(y+2x)}{x(x+2y)}$

7. $x^3y^3 - y - x = 0$

$3x^3y^2y' + 3x^2y^3 - y' - 1 = 0$

$(3x^3y^2 - 1)y' = 1 - 3x^2y^3$

$y' = \frac{1 - 3x^2y^3}{3x^3y^2 - 1}$

8. $(xy)^{1/2} - x + 2y = 0$

$\frac{1}{2}(xy)^{-1/2}(xy' + y) - 1 + 2y' = 0$

$\frac{x}{2\sqrt{xy}}y' + \frac{y}{2\sqrt{xy}} - 1 + 2y' = 0$

$xy' + y - 2\sqrt{xy} + 4\sqrt{xy}y' = 0$

$y' = \frac{2\sqrt{xy} - y}{4\sqrt{xy} + x}$

10. $2 \sin x \cos y = 1$

$2[\sin x(-\sin y)y' + \cos y(\cos x)] = 0$

$y' = \frac{\cos x \cos y}{\sin x \sin y}$

$= \cot x \cot y$

9. $x^3 - 3x^2y + 2xy^2 = 12$

$3x^2 - 3x^2y' - 6xy + 4xyy' + 2y^2 = 0$

$(4xy - 3x^2)y' = 6xy - 3x^2 - 2y^2$

$y' = \frac{6xy - 3x^2 - 2y^2}{4xy - 3x^2}$

12. $(\sin \pi x + \cos \pi y)^2 = 2$

$2(\sin \pi x + \cos \pi y)[\pi \cos \pi x - \pi(\sin \pi y)y'] = 0$

$\pi \cos \pi x - \pi(\sin \pi y)y' = 0$

$y' = \frac{\cos \pi x}{\sin \pi y}$

13. $\sin x = x(1 + \tan y)$

$\cos x = x(\sec^2 y)y' + (1 + \tan y)(1)$

$y' = \frac{\cos x - \tan y - 1}{x \sec^2 y}$

14. $\cot y = x - y$

$(-\csc^2 y)y' = 1 - y'$

$y' = \frac{1}{1 - \csc^2 y}$

$= \frac{1}{-\cot^2 y} = -\tan^2 y$

15.

$y = \sin(xy)$

$y' = [xy' + y]\cos(xy)$

$y' - x \cos(xy)y' = y \cos(xy)$

$y' = \frac{y \cos(xy)}{1 - x \cos(xy)}$

16. $x = \sec \frac{1}{y}$

$1 = -\frac{y'}{y^2} \sec \frac{1}{y} \tan \frac{1}{y}$

$y' = \frac{-y^2}{\sec(1/y) \tan(1/y)}$

$= -y^2 \cos\left(\frac{1}{y}\right) \cot\left(\frac{1}{y}\right)$

17. (a) $x^2 + y^2 = 16$

$y^2 = 16 - x^2$


$y = \pm \sqrt{16 - x^2}$

(c) Explicitly:

$\frac{dy}{dx} = \pm \frac{1}{2}(16 - x^2)^{-1/2}(-2x)$

$= \frac{\mp x}{\sqrt{16 - x^2}} = \frac{-x}{\pm \sqrt{16 - x^2}} = \frac{-x}{y}$

(b)



(d) Implicitly: $2x + 2yy' = 0$

$y' = -\frac{x}{y}$

18. (a) $(x^2 - 4x + 4) + (y^2 + 6y + 9) = -9 + 4 + 9$

$$(x - 2)^2 + (y + 3)^2 = 4, \text{ Circle}$$

$$(y + 3)^2 = 4 - (x - 2)^2$$

$$y = -3 \pm \sqrt{4 - (x - 2)^2}$$

(c) Explicitly:


$$\frac{dy}{dx} = \pm \frac{1}{2}[4 - (x - 2)^2]^{-1/2}(-2)(x - 2)$$

$$= \frac{\mp(x - 2)}{\sqrt{4 - (x - 2)^2}}$$

$$= \frac{\mp(x - 2)}{\pm\sqrt{4 - (x - 2)^2}}$$

$$= \frac{\mp(x - 2)}{-3 \pm \sqrt{4 - (x - 2)^2} + 3}$$

$$= \frac{\mp(x - 2)}{y + 3}$$



(d) Implicitly:

$$2x + 2yy' - 4 + 6y' = 0$$

$$(2y + 6)y' = -2(x - 2)$$

$$y' = \frac{-(x - 2)}{y + 3}$$

19. (a) $16y^2 = 144 - 9x^2$


$$y^2 = \frac{1}{16}(144 - 9x^2) = \frac{9}{16}(16 - x^2)$$

$$y = \pm \frac{3}{4}\sqrt{16 - x^2}$$

(c) Explicitly:

$$\frac{dy}{dx} = \pm \frac{3}{8}(16 - x^2)^{-1/2}(-2x)$$

$$= \mp \frac{3x}{4\sqrt{16 - x^2}} = \frac{-3x}{4(4/3)y} = \frac{-9x}{16y}$$



(d) Implicitly: $18x + 32yy' = 0$

$$y' = \frac{-9x}{16y}$$

20. (a) $9y^2 = x^2 + 9$


$$y^2 = \frac{x^2}{9} + 1 = \frac{x^2 + 9}{9}$$

$$y = \frac{\pm\sqrt{x^2 + 9}}{3}$$

(c) Explicitly:

$$\frac{dy}{dx} = \frac{\pm\frac{1}{2}(x^2 + 9)^{-1/2}(2x)}{3}$$

$$= \frac{\pm x}{3\sqrt{x^2 + 9}} = \frac{\pm x}{3(\pm 3y)} = \frac{x}{9y}$$



(d) Implicitly: $9y^2 - x^2 = 9$

$$18yy' - 2x = 0$$

$$18yy' = 2x$$

$$y' = \frac{2x}{18y} = \frac{x}{9y}$$

21. $xy = 4$

$$xy' + y(1) = 0$$

$$xy' = -y$$

$$y' = \frac{-y}{x}$$

$$\text{At } (-4, -1): y' = -\frac{1}{4}$$

22. $x^2 - y^3 = 0$

$$2x - 3y^2y' = 0$$

$$y' = \frac{2x}{3y^2}$$

$$\text{At } (1, 1): y' = \frac{2}{3}$$

23. $y^2 = \frac{x^2 - 4}{x^2 + 4}$

$$2yy' = \frac{(x^2 + 4)(2x) - (x^2 - 4)(2x)}{(x^2 + 4)^2}$$

$$2yy' = \frac{16x}{(x^2 + 4)^2}$$

$$y' = \frac{8x}{y(x^2 + 4)^2}$$

At $(2, 0)$: y' is undefined.

24. $(x + y)^3 = x^3 + y^3$

$$x^3 + 3x^2y + 3xy^2 + y^3 = x^3 + y^3$$

$$3x^2y + 3xy^2 = 0$$

$$x^2y + xy^2 = 0$$

$$x^2y' + 2xy + 2xyy' + y^2 = 0$$

$$(x^2 + 2xy)y' = -(y^2 + 2xy)$$

$$y' = -\frac{y(y + 2x)}{x(x + 2y)}$$

At $(-1, 1)$: $y' = -1$

25. $x^{2/3} + y^{2/3} = 5$

$$\frac{2}{3}x^{-1/3} + \frac{2}{3}y^{-1/3}y' = 0$$

$$y' = \frac{-x^{-1/3}}{y^{-1/3}} = -\sqrt[3]{\frac{y}{x}}$$

$$\text{At } (8, 1): y' = -\frac{1}{2}$$

26. $x^3 + y^3 = 4xy + 1$

$$3x^2 + 3y^2y' = 4xy' + 4y$$

$$(3y^2 - 4x)y' = 4y - 3x^2$$

$$y' = \frac{4y - 3x^2}{(3y^2 - 4x)}$$

$$\text{At } (2, 1): y' = \frac{4 - 12}{3 - 8} = \frac{8}{5}$$

27. $\tan(x + y) = x$

$$(1 + y') \sec^2(x + y) = 1$$

$$y' = \frac{1 - \sec^2(x + y)}{\sec^2(x + y)}$$

$$= \frac{-\tan^2(x + y)}{\tan^2(x + y) + 1}$$

$$= -\sin^2(x + y)$$

$$= -\frac{x^2}{x^2 + 1}$$

$$\text{At } (0, 0): y' = 0$$

28. $x \cos y = 1$

$$x[-y' \sin y] + \cos y = 0$$

$$y' = \frac{\cos y}{x \sin y}$$

$$= \frac{1}{x} \cot y$$

$$= \frac{\cot y}{x}$$

$$\text{At } \left(2, \frac{\pi}{3}\right): y' = \frac{1}{2\sqrt{3}}$$

29. $(x^2 + 4)y = 8$

$$(x^2 + 4)y' + y(2x) = 0$$

$$y' = \frac{-2xy}{x^2 + 4}$$

$$= \frac{-2x[8/(x^2 + 4)]}{x^2 + 4}$$

$$= \frac{-16x}{(x^2 + 4)^2}$$

$$\text{At } (2, 1): y' = \frac{-32}{64} = -\frac{1}{2}$$

$$\left(\text{Or, you could just solve for } y: y = \frac{8}{x^2 + 4}\right)$$

30. $(4 - x)y^2 = x^3$

$$(4 - x)(2yy') + y^2(-1) = 3x^2$$

$$y' = \frac{3x^2 + y^2}{2y(4 - x)}$$

$$\text{At } (2, 2): y' = 2$$

31. $(x^2 + y^2)^2 = 4x^2y$

$$2(x^2 + y^2)(2x + 2yy') = 4x^2y' + y(8x)$$

$$4x^3 + 4x^2yy' + 4xy^2 + 4y^3y' = 4x^2y' + 8xy$$

$$4x^2yy' + 4y^3y' - 4x^2y' = 8xy - 4x^3 - 4xy^2$$

$$4y'(x^2y + y^3 - x^2) = 4(2xy - x^3 - xy^2)$$

$$y' = \frac{2xy - x^3 - xy^2}{x^2y + y^3 - x^2}$$

At $(1, 1)$: $y' = 0$

33. $(y - 2)^2 = 4(x - 3), (4, 0)$

$$2(y - 2)y' = 4$$

$$y' = \frac{2}{y - 2}$$

At $(4, 0)$: $y' = -1$

Tangent line: $y - 0 = -1(x - 4)$

$$y = -x + 4$$

35. $xy = 1, (1, 1)$

$$xy' + y = 0$$

$$y' = \frac{-y}{x}$$

At $(1, 1)$: $y' = -1$

Tangent line: $y - 1 = -1(x - 1)$

$$y = -x + 2$$

37. $x^2y^2 - 9x^2 - 4y^2 = 0, (-4, 2\sqrt{3})$

$$x^2 2yy' + 2xy^2 - 18x - 8yy' = 0$$

$$y' = \frac{18x - 2xy^2}{2x^2y - 8y}$$

$$\text{At } (-4, 2\sqrt{3}): y' = \frac{18(-4) - 2(-4)(12)}{2(16)(2\sqrt{3}) - 16\sqrt{3}}$$

$$= \frac{24}{48\sqrt{3}} = \frac{1}{2\sqrt{3}} = \frac{\sqrt{3}}{6}$$

$$\text{Tangent line: } y - 2\sqrt{3} = \frac{\sqrt{3}}{6}(x + 4)$$

$$y = \frac{\sqrt{3}}{6}x + \frac{8}{3}\sqrt{3}$$

32. $x^3 + y^3 - 6xy = 0$

$$3x^2 + 3y^2y' - 6xy' - 6y = 0$$

$$y'(3y^2 - 6x) = 6y - 3x^2$$

$$y' = \frac{6y - 3x^2}{3y^2 - 6x} = \frac{2y - x^2}{y^2 - 2x}$$

$$\text{At } \left(\frac{4}{3}, \frac{8}{3}\right): y' = \frac{(16/3) - (16/9)}{(64/9) - (8/3)} = \frac{32}{40} = \frac{4}{5}$$

34. $(x + 1)^2 + (y - 2)^2 = 20, (3, 4)$

$$2(x + 1) + 2(y - 2)y' = 0$$

$$2(y - 2)y' = -2(x + 1)$$

$$y' = \frac{-(x + 1)}{y - 2}$$

At $(3, 4)$: $y' = -2$

Tangent line: $y - 4 = -2(x - 3)$

$$y = -2x + 10$$

36. $7x^2 - 6\sqrt{3}xy + 13y^2 - 16 = 0, (\sqrt{3}, 1)$

$$14x - 6\sqrt{3}xy' - 6\sqrt{3}y + 26yy' = 0$$

$$y' = \frac{6\sqrt{3}y - 14x}{26y - 6\sqrt{3}x}$$

$$\text{At } (\sqrt{3}, 1): y' = \frac{6\sqrt{3} - 14\sqrt{3}}{26 - 6\sqrt{3}\sqrt{3}} = \frac{-8\sqrt{3}}{8} = -\sqrt{3}$$

Tangent line: $y - 1 = -\sqrt{3}(x - \sqrt{3})$

$$y = -\sqrt{3}x + 4$$

38. $x^{2/3} + y^{2/3} = 5, (8, 1)$

$$\frac{2}{3}x^{-1/3} + \frac{2}{3}y^{-1/3}y' = 0$$

$$y' = \frac{-x^{-1/3}}{y^{-1/3}} = -\left(\frac{y}{x}\right)^{1/3}$$

$$\text{At } (8, 1): y' = -\frac{1}{2}$$

$$\text{Tangent line: } y - 1 = -\frac{1}{2}(x - 8)$$

$$y = -\frac{1}{2}x + 5$$

39. $3(x^2 + y^2)^2 = 100(x^2 - y^2), \quad (4, 2)$

$$6(x^2 + y^2)(2x + 2yy') = 100(2x - 2yy')$$

At $(4, 2)$:

$$6(16 + 4)(8 + 4y') = 100(8 - 4y')$$

$$960 + 480y' = 800 - 400y'$$

$$880y' = -160$$

$$y' = -\frac{2}{11}$$

Tangent line: $y - 2 = -\frac{2}{11}(x - 4)$

$$11y + 2x - 30 = 0$$

$$y = -\frac{2}{11}x + \frac{30}{11}$$

41. (a) $\frac{x^2}{2} + \frac{y^2}{8} = 1, \quad (1, 2)$

$$x + \frac{yy'}{4} = 0$$

$$y' = -\frac{4x}{y}$$

At $(1, 2)$: $y' = -2$

Tangent line: $y - 2 = -2(x - 1)$

$$y = -2x + 4$$

Note: From part (a), $\frac{1(x)}{2} + \frac{2(y)}{8} = 1 \Rightarrow \frac{1}{4}y = -\frac{1}{2}x + 1 \Rightarrow y = -2x + 4$, Tangent line.

42. (a) $\frac{x^2}{6} - \frac{y^2}{8} = 1, \quad (3, -2)$

$$\frac{x}{3} - \frac{y}{4}y' = 0$$

$$\frac{y}{4}y' = \frac{x}{3}$$

$$y' = \frac{4x}{3y}$$

At $(3, -2)$: $y' = \frac{4(3)}{3(-2)} = -2$

Tangent line: $y + 2 = -2(x - 3)$

$$y = -2x + 4$$

Note: From part (a), $\frac{3x}{6} - \frac{(-2)y}{8} = 1 \Rightarrow \frac{1}{2}x + \frac{y}{4} = 1 \Rightarrow y = -2x + 4$, Tangent line.

40. $y^2(x^2 + y^2) = 2x^2, \quad (1, 1)$

$$y^2x^2 + y^4 = 2x^2$$

$$2yy'x^2 + 2xy^2 + 4y^3y' = 4x$$

At $(1, 1)$:

$$2y' + 2 + 4y' = 4$$

$$6y' = 2$$

$$y' = \frac{1}{3}$$

Tangent line: $y - 1 = \frac{1}{3}(x - 1)$

$$y = \frac{1}{3}x + \frac{2}{3}$$

(b) $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \Rightarrow \frac{2x}{a^2} + \frac{2yy'}{b^2} = 0 \Rightarrow y' = -\frac{b^2x}{a^2y}$

$$y - y_0 = \frac{-b^2x_0}{a^2y_0}(x - x_0), \quad \text{Tangent line at } (x_0, y_0)$$

$$\frac{y_0y}{b^2} - \frac{y_0^2}{b^2} = \frac{-x_0x}{a^2} + \frac{x_0^2}{a^2}$$

Since $\frac{x_0^2}{a^2} + \frac{y_0^2}{b^2} = 1$, you have $\frac{y_0y}{b^2} + \frac{x_0x}{a^2} = 1$.

(b) $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \Rightarrow \frac{2x}{a^2} - \frac{2yy'}{b^2} = 0 \Rightarrow y' = \frac{xb^2}{ya^2}$

$$y - y_0 = \frac{x_0b^2}{y_0a^2}(x - x_0), \quad \text{Tangent line at } (x_0, y_0)$$

$$\frac{yy_0}{b^2} - \frac{y_0^2}{b^2} = \frac{x_0x}{a^2} - \frac{x_0^2}{a^2}$$

Since $\frac{x_0^2}{a^2} - \frac{y_0^2}{b^2} = 1$, you have $\frac{x_0x}{a^2} - \frac{yy_0}{b^2} = 1$.

43. $\tan y = x$

$y' \sec^2 y = 1$

$y' = \frac{1}{\sec^2 y} = \cos^2 y, -\frac{\pi}{2} < y < \frac{\pi}{2}$

$\sec^2 y = 1 + \tan^2 y = 1 + x^2$

$y' = \frac{1}{1+x^2}$

44. $\cos y = x$

$-\sin y \cdot y' = 1$

$y' = \frac{-1}{\sin y}, 0 < y < \pi$

$\sin^2 y + \cos^2 y = 1$

$\sin^2 y = 1 - \cos^2 y$

$\sin y = \sqrt{1 - \cos^2 y} = \sqrt{1 - x^2}$

$y' = \frac{-1}{\sqrt{1-x^2}}, -1 < x < 1$

45. $x^2 + y^2 = 36$

$2x + 2yy' = 0$

$y' = \frac{-x}{y}$

$y'' = \frac{y(-1) + xy'}{y^2} = \frac{-y + x(-x/y)}{y^2} = \frac{-y^2 - x^2}{y^3} = \frac{-36}{y^3}$

46. $x^2y^2 - 2x = 3$

$2x^2yy' + 2xy^2 - 2 = 0$

$x^2yy' + xy^2 - 1 = 0$

$y' = \frac{1 - xy^2}{x^2y}$

$2xyy' + x^2(y')^2 + x^2yy'' + 2xyy' + y^2 = 0$

$4xyy' + x^2(y')^2 + x^2yy'' + y^2 = 0$

$\frac{4 - 4xy^2}{x} + \frac{(1 - xy^2)^2}{x^2y^2} + x^2yy'' + y^2 = 0$

$4xy^2 - 4x^2y^4 + 1 - 2xy^2 + x^2y^4 + x^4y^3y'' + x^2y^4 = 0$

$x^4y^3y'' = 2x^2y^4 - 2xy^2 - 1$

$y'' = \frac{2x^2y^4 - 2xy^2 - 1}{x^4y^3}$

47. $x^2 - y^2 = 16$

$2x - 2yy' = 0$

$y' = \frac{x}{y}$

$x - yy' = 0$

$1 - yy'' - (y')^2 = 0$

$1 - yy'' - \left(\frac{x}{y}\right)^2 = 0$

$y^2 - y^3y'' = x^2$

$y'' = \frac{y^2 - x^2}{y^3} = \frac{-16}{y^3}$

48. $1 - xy = x - y$

$y - xy = x - 1$

$y = \frac{x-1}{1-x} = -1$

$y' = 0$

$y'' = 0$

49. $y^2 = x^3$

$2yy' = 3x^2$

$y' = \frac{3x^2}{2y} = \frac{3x^2}{2y} \cdot \frac{xy}{xy}$

$= \frac{3y}{2x} \cdot \frac{x^3}{y^2} = \frac{3y}{2x}$

$y'' = \frac{2x(3y') - 3y(2)}{4x^2}$

$= \frac{2x[3 \cdot (3y/2x)] - 6y}{4x^2}$

$= \frac{3y}{4x^2} = \frac{3x}{4y}$

50. $y^2 = 4x$

$2yy' = 4$

$y' = \frac{2}{y}$

$y'' = -2y^{-2}y' = \left[\frac{-2}{y^2} \right] \cdot \frac{2}{y} = \frac{-4}{y^3}$

51. $\sqrt{x} + \sqrt{y} = 4$

$\frac{1}{2}x^{-1/2} + \frac{1}{2}y^{-1/2}y' = 0$


$y' = \frac{-\sqrt{y}}{\sqrt{x}}$

At $(9, 1)$: $y' = -\frac{1}{3}$

Tangent line: $y - 1 = -\frac{1}{3}(x - 9)$

$y = -\frac{1}{3}x + 4$

$x + 3y - 12 = 0$



52. $y^2 = \frac{x-1}{x^2+1}$

$2yy' = \frac{(x^2+1)(1) - (x-1)(2x)}{(x^2+1)^2}$

$= \frac{x^2+1-2x^2+2x}{(x^2+1)^2}$


$y' = \frac{1+2x-x^2}{2y(x^2+1)^2}$

At $\left(2, \frac{\sqrt{5}}{5}\right)$: $y' = \frac{1+4-4}{[(2\sqrt{5})/5](4+1)^2} = \frac{1}{10\sqrt{5}}$

Tangent line: $y - \frac{\sqrt{5}}{5} = \frac{1}{10\sqrt{5}}(x - 2)$

$10\sqrt{5}y - 10 = x - 2$

$x - 10\sqrt{5}y + 8 = 0$



53. $x^2 + y^2 = 25$

$2x + 2yy' = 0$

$y' = \frac{-x}{y}$

At $(4, 3)$:



Tangent line: $y - 3 = \frac{-4}{3}(x - 4) \Rightarrow 4x + 3y - 25 = 0$

Normal line: $y - 3 = \frac{3}{4}(x - 4) \Rightarrow 3x - 4y = 0$

At $(-3, 4)$:

Tangent line: $y - 4 = \frac{3}{4}(x + 3) \Rightarrow 3x - 4y + 25 = 0$

Normal line: $y - 4 = \frac{-4}{3}(x + 3) \Rightarrow 4x + 3y = 0$



54. $x^2 + y^2 = 9$

$$y' = \frac{-x}{y}$$

At $(0, 3)$:



Tangent line: $y = 3$

Normal line: $x = 0$

At $(2, \sqrt{5})$:

$$\text{Tangent line: } y - \sqrt{5} = \frac{-2}{\sqrt{5}}(x - 2) \Rightarrow 2x + \sqrt{5}y - 9 = 0$$

$$\text{Normal line: } y - \sqrt{5} = \frac{\sqrt{5}}{2}(x - 2) \Rightarrow \sqrt{5}x - 2y = 0$$



55. $x^2 + y^2 = r^2$

$$2x + 2yy' = 0$$

$$y' = \frac{-x}{y} = \text{slope of tangent line}$$

$$\frac{y}{x} = \text{slope of normal line}$$

Let (x_0, y_0) be a point on the circle. If $x_0 = 0$, then the tangent line is horizontal, the normal line is vertical and, hence, passes through the origin. If $x_0 \neq 0$, then the equation of the normal line is

$$y - y_0 = \frac{y_0}{x_0}(x - x_0)$$

$$y = \frac{y_0}{x_0}x$$

which passes through the origin.

56. $y^2 = 4x$

$$2yy' = 4$$

$$y' = \frac{2}{y} = 1 \text{ at } (1, 2)$$

Equation of normal line at $(1, 2)$ is $y - 2 = -1(x - 1)$, $y = 3 - x$. The centers of the circles must be on the normal line and at a distance of 4 units from $(1, 2)$. Therefore,

$$(x - 1)^2 + [(3 - x) - 2]^2 = 16$$

$$2(x - 1)^2 = 16$$

$$x = 1 \pm 2\sqrt{2}.$$

Centers of the circles: $(1 + 2\sqrt{2}, 2 - 2\sqrt{2})$ and $(1 - 2\sqrt{2}, 2 + 2\sqrt{2})$

$$(x - 1 - 2\sqrt{2})^2 + (y - 2 + 2\sqrt{2})^2 = 16$$

$$(x - 1 + 2\sqrt{2})^2 + (y - 2 - 2\sqrt{2})^2 = 16$$

57. $25x^2 + 16y^2 + 200x - 160y + 400 = 0$

$$50x + 32yy' + 200 - 160y' = 0$$

$$y' = \frac{200 + 50x}{160 - 32y}$$

Horizontal tangents occur when $x = -4$:

$$25(16) + 16y^2 + 200(-4) - 160y + 400 = 0$$

$$y(y - 10) = 0 \Rightarrow y = 0, 10$$


Horizontal tangents: $(-4, 0), (-4, 10)$

Vertical tangents occur when $y = 5$:

$$25x^2 + 400 + 200x - 800 + 400 = 0$$

$$25x(x + 8) = 0 \Rightarrow x = 0, -8$$

Vertical tangents: $(0, 5), (-8, 5)$



58. $4x^2 + y^2 - 8x + 4y + 4 = 0$

$$8x + 2yy' - 8 + 4y' = 0$$

$$y' = \frac{8 - 8x}{2y + 4} = \frac{4 - 4x}{y + 2}$$

Horizontal tangents occur when $x = 1$:

$$4(1)^2 + y^2 - 8(1) + 4y + 4 = 0$$

$$y^2 + 4y = y(y + 4) = 0 \Rightarrow y = 0, -4$$


Horizontal tangents: $(1, 0), (1, -4)$

Vertical tangents occur when $y = -2$:

$$4x^2 + (-2)^2 - 8x + 4(-2) + 4 = 0$$

$$4x^2 - 8x = 4x(x - 2) = 0 \Rightarrow x = 0, 2$$

Vertical tangents: $(0, -2), (2, -2)$



59. Find the points of intersection by letting $y^2 = 4x$ in the equation $2x^2 + y^2 = 6$.

$$2x^2 + 4x = 6 \quad \text{and} \quad (x + 3)(x - 1) = 0$$

The curves intersect at $(1, \pm 2)$.

Ellipse:


$$4x + 2yy' = 0$$

$$y' = -\frac{2x}{y}$$

Parabola:

$$2yy' = 4$$

$$y' = \frac{2}{y}$$



At $(1, 2)$, the slopes are:

$$y' = -1$$

$$y' = 1$$

At $(1, -2)$, the slopes are:

$$y' = 1$$

$$y' = -1$$

Tangents are perpendicular.

60. Find the points of intersection by letting $y^2 = x^3$ in the equation $2x^2 + 3y^2 = 5$.

$$2x^2 + 3x^3 = 5 \quad \text{and} \quad 3x^3 + 2x^2 - 5 = 0$$

Intersect when $x = 1$.

Points of intersection: $(1, \pm 1)$

$$y^2 = x^3:$$

$$2yy' = 3x^2$$

$$y' = \frac{3x^2}{2y}$$

$$2x^2 + 3y^2 = 5:$$

$$4x + 6yy' = 0$$

$$y' = -\frac{2x}{3y}$$

At $(1, 1)$, the slopes are:

$$y' = \frac{3}{2}$$


$$y' = -\frac{2}{3}$$

At $(1, -1)$, the slopes are:

$$y' = -\frac{3}{2}$$

$$y' = \frac{2}{3}$$


Tangents are perpendicular.



61. $y = -x$ and $x = \sin y$

Point of intersection: $(0, 0)$

$$\begin{aligned}y &= -x \\y' &= -1 \\x &= \sin y \\1 &= y' \cos y \\y' &= \sec y\end{aligned}$$




At $(0, 0)$, the slopes are:

$$y' = -1 \quad y' = 1$$

Tangents are perpendicular.



62. Rewriting each equation and differentiating:

$$\begin{aligned}x^3 &= 3(y - 1) & x(3y - 29) &= 3 \\y &= \frac{x^3}{3} + 1 & y &= \frac{1}{3}(x + 29) \\y' &= x^2 & y' &= -\frac{1}{x^2}\end{aligned}$$





For each value of x , the derivatives are negative reciprocals of each other. Thus, the tangent lines are orthogonal at both points of intersection.

63. $xy = C$ $x^2 - y^2 = K$
 $xy' + y = 0$ $2x - 2yy' = 0$
 $y' = -\frac{y}{x}$ $y' = \frac{x}{y}$



At any point of intersection (x, y) the product of the slopes is $(-y/x)(x/y) = -1$. The curves are orthogonal.

64. $x^2 + y^2 = C^2$ $y = Kx$
 $2x + 2yy' = 0$ $y' = K$
 $y' = -\frac{x}{y}$



At the point of intersection (x, y) , the product of the slopes is $(-x/y)(K) = (-x/Kx)(K) = -1$. The curves are orthogonal.

65. $2y^2 - 3x^4 = 0$

(a) $4yy' - 12x^3 = 0$

$$4yy' = 12x^3$$

$$y' = \frac{12x^3}{4y} = \frac{3x^3}{y}$$

(b) $4y \frac{dy}{dt} - 12x^3 \frac{dx}{dt} = 0$

$$y \frac{dy}{dt} = 3x^3 \frac{dx}{dt}$$

66. $x^2 - 3xy^2 + y^3 = 10$

(a) $2x - 3y^2 - 6xyy' + 3y^2y' = 0$

$$(-6xy + 3y^2)y' = 3y^2 - 2x$$

$$y' = \frac{3y^2 - 2x}{3y^2 - 6xy}$$

(b) $2x \frac{dx}{dt} - 3y^2 \frac{dx}{dt} - 6xy \frac{dy}{dt} + 3y^2 \frac{dy}{dt} = 0$

$$(2x - 3y^2) \frac{dx}{dt} = (6xy - 3y^2) \frac{dy}{dt}$$

67. $\cos \pi y - 3 \sin \pi x = 1$

(a) $-\pi \sin(\pi y)y' - 3\pi \cos(\pi x) = 0$


$$y' = \frac{-3\cos(\pi x)}{\sin(\pi y)}$$

(b) $-\pi \sin(\pi y)\frac{dy}{dt} - 3\pi \cos(\pi x)\frac{dx}{dt} = 0$

$$-\sin(\pi y)\frac{dy}{dt} = 3\cos(\pi x)\frac{dx}{dt}$$

69. A function is in explicit form if y is written as a function of x : $y = f(x)$. For example, $y = x^3$. An implicit equation is not in the form $y = f(x)$. For example, $x^2 + y^2 = 5$.

71.

Use starting point B .

73. (a) $x^4 = 4(4x^2 - y^2)$

$$4y^2 = 16x^2 - x^4$$

$$y^2 = 4x^2 - \frac{1}{4}x^4$$

$$y = \pm \sqrt{4x^2 - \frac{1}{4}x^4}$$

(b) $y = 3 \Rightarrow 9 = 4x^2 - \frac{1}{4}x^4$

$$36 = 16x^2 - x^4$$

$$x^4 - 16x^2 + 36 = 0$$

$$x^2 = \frac{16 \pm \sqrt{256 - 144}}{2} = 8 \pm \sqrt{28}$$

Note that $x^2 = 8 \pm \sqrt{28} = 8 \pm 2\sqrt{7} = (1 \pm \sqrt{7})^2$. Hence, there are four values of x :

$$-1 - \sqrt{7}, 1 - \sqrt{7}, -1 + \sqrt{7}, 1 + \sqrt{7}$$

To find the slope, $2yy' = 8x - x^3 \Rightarrow y' = \frac{x(8 - x^2)}{2(3)}$.

68. $4 \sin x \cos y = 1$



(a) $4 \sin x(-\sin y)y' + 4 \cos x \cos y = 0$

$$y' = \frac{\cos x \cos y}{\sin x \sin y}$$

(b) $4 \sin x(-\sin y)\frac{dy}{dt} + 4 \cos x \frac{dx}{dt} \cos y = 0$

$$\cos x \cos y \frac{dx}{dt} = \sin x \sin y \frac{dy}{dt}$$

70. Given an implicit equation, first differentiate both sides with respect to x . Collect all terms involving y' on the left, and all other terms to the right. Factor out y' on the left side. Finally, divide both sides by the left-hand factor that does not contain y' .

72. Highest wind speed near L 

—CONTINUED—

73. —CONTINUED—

For $x = -1 - \sqrt{7}$, $y' = \frac{1}{3}(\sqrt{7} + 7)$, and the line is

$$y_1 = \frac{1}{3}(\sqrt{7} + 7)(x + 1 + \sqrt{7}) + 3 = \frac{1}{3}[(\sqrt{7} + 7)x + 8\sqrt{7} + 23].$$

For $x = 1 - \sqrt{7}$, $y' = \frac{1}{3}(\sqrt{7} - 7)$, and the line is

$$y_2 = \frac{1}{3}(\sqrt{7} - 7)(x - 1 + \sqrt{7}) + 3 = \frac{1}{3}[(\sqrt{7} - 7)x + 23 - 8\sqrt{7}].$$

For $x = -1 + \sqrt{7}$, $y' = -\frac{1}{3}(\sqrt{7} - 7)$, and the line is

$$y_3 = -\frac{1}{3}(\sqrt{7} - 7)(x + 1 - \sqrt{7}) + 3 = -\frac{1}{3}[(\sqrt{7} - 7)x - (23 - 8\sqrt{7})].$$

For $x = 1 + \sqrt{7}$, $y' = -\frac{1}{3}(\sqrt{7} + 7)$, and the line is

$$y_4 = -\frac{1}{3}(\sqrt{7} + 7)(x - 1 - \sqrt{7}) + 3 = -\frac{1}{3}[(\sqrt{7} + 7)x - (8\sqrt{7} + 23)].$$

(c) Equating y_3 and y_4 :

$$-\frac{1}{3}(\sqrt{7} - 7)(x + 1 - \sqrt{7}) + 3 = -\frac{1}{3}(\sqrt{7} + 7)(x - 1 - \sqrt{7}) + 3$$

$$(\sqrt{7} - 7)(x + 1 - \sqrt{7}) = (\sqrt{7} + 7)(x - 1 - \sqrt{7})$$

$$\sqrt{7}x + \sqrt{7} - 7 - 7x - 7 + 7\sqrt{7} = \sqrt{7}x - \sqrt{7} - 7 + 7x - 7 - 7\sqrt{7}$$

$$16\sqrt{7} = 14x$$

$$x = \frac{8\sqrt{7}}{7}$$

If $x = \frac{8\sqrt{7}}{7}$, then $y = 5$ and the lines intersect at $\left(\frac{8\sqrt{7}}{7}, 5\right)$.

74. $\sqrt{x} + \sqrt{y} = \sqrt{c}$

$$\frac{1}{2\sqrt{x}} + \frac{1}{2\sqrt{y}} \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = -\frac{\sqrt{y}}{\sqrt{x}}$$

$$\text{Tangent line at } (x_0, y_0): y - y_0 = -\frac{\sqrt{y_0}}{\sqrt{x_0}}(x - x_0)$$

$$x\text{-intercept: } (x_0 + \sqrt{x_0}\sqrt{y_0}, 0)$$

$$y\text{-intercept: } (0, y_0 + \sqrt{x_0}\sqrt{y_0})$$

Sum of intercepts:

$$\begin{aligned} (x_0 + \sqrt{x_0}\sqrt{y_0}) + (y_0 + \sqrt{x_0}\sqrt{y_0}) &= x_0 + 2\sqrt{x_0}\sqrt{y_0} + y_0 \\ &= (\sqrt{x_0} + \sqrt{y_0})^2 \\ &= (\sqrt{c})^2 = c \end{aligned}$$

75. $y = x^{p/q}; p, q \text{ integers and } q > 0$

$$y^q = x^p$$

$$qy^{q-1}y' = px^{p-1}$$

$$y' = \frac{p}{q} \cdot \frac{x^{p-1}}{y^{q-1}} = \frac{p}{q} \cdot \frac{x^{p-1}y}{y^q}$$

$$= \frac{p}{q} \cdot \frac{x^{p-1}}{x^p} x^{p/q} = \frac{p}{q} x^{p/q-1}$$

Thus, if $y = x^n$, $n = p/q$, then $y' = nx^{n-1}$.

76. $x^2 + y^2 = 25$, slope = $\frac{3}{4}$

$$2x + 2yy' = 0$$

$$y' = -\frac{x}{y} = \frac{3}{4} \Rightarrow y = -\frac{4}{3}x$$

$$x^2 + \left(\frac{16}{9}x^2\right) = 25$$

$$\frac{25}{9}x^2 = 25$$

$$x = \pm 3$$

Points: $(3, -4)$ and $(-3, 4)$

78. $\frac{x^2}{4} + \frac{y^2}{9} = 1$, $(4, 0)$

$$\frac{2x}{4} + \frac{2yy'}{9} = 0$$

$$y' = \frac{-9x}{4y}$$

$$\frac{-9x}{4y} = \frac{y - 0}{x - 4}$$

$$-9x(x - 4) = 4y^2$$

But, $9x^2 + 4y^2 = 36 \Rightarrow 4y^2 = 36 - 9x^2$. Hence, $-9x^2 + 36x = 4y^2 = 36 - 9x^2 \Rightarrow x = 1$.

Points on ellipse: $\left(1, \pm \frac{3}{2}\sqrt{3}\right)$

$$\text{At } \left(1, \frac{3}{2}\sqrt{3}\right): y' = \frac{-9x}{4y} = \frac{-9}{4[(3/2)\sqrt{3}]} = -\frac{\sqrt{3}}{2}$$

$$\text{At } \left(1, -\frac{3}{2}\sqrt{3}\right): y' = \frac{\sqrt{3}}{2}$$

$$\text{Tangent lines: } y = -\frac{\sqrt{3}}{2}(x - 4) = -\frac{\sqrt{3}}{2}x + 2\sqrt{3}$$

$$y = \frac{\sqrt{3}}{2}(x - 4) = \frac{\sqrt{3}}{2}x - 2\sqrt{3}$$

79. $x = y^2$

$$1 = 2yy'$$

$$y' = \frac{1}{2y}, \text{ slope of tangent line}$$

Consider the slope of the normal line joining $(x_0, 0)$ and $(x, y) = (y^2, y)$ on the parabola.

$$-2y = \frac{y - 0}{y^2 - x_0}$$

$$y^2 - x_0 = -\frac{1}{2}$$

$$y^2 = x_0 - \frac{1}{2}$$

77. $y^4 = y^2 - x^2$

$$4y^3y' = 2yy' - 2x$$

$$2x = (2y - 4y^3)y'$$

$$y' = \frac{2x}{2y - 4y^3} = 0 \Rightarrow x = 0$$

Horizontal tangents at $(0, 1)$ and $(0, -1)$

Note: $y^4 - y^2 + x^2 = 0$

$$y^2 = \frac{1 \pm \sqrt{1 - 4x^2}}{2}$$

If you graph these four equations, you will see that these are horizontal tangents at $(0, \pm 1)$, but not at $(0, 0)$.

79. —CONTINUED—

- (a) If $x_0 = \frac{1}{4}$, then $y^2 = \frac{1}{4} - \frac{1}{2} = -\frac{1}{4}$, which is impossible. Thus, the only normal line is the x -axis ($y = 0$).
 (b) If $x_0 = \frac{1}{2}$, then $y^2 = 0 \Rightarrow y = 0$. Same as part (a).
 (c) If $x_0 = 1$, then $y^2 = \frac{1}{2} = x$ and there are three normal lines:

The x -axis, the line joining $(x_0, 0)$ and $\left(\frac{1}{2}, \frac{1}{\sqrt{2}}\right)$, and the line joining $(x_0, 0)$ and $\left(\frac{1}{2}, -\frac{1}{\sqrt{2}}\right)$

If two normals are perpendicular, then their slopes are -1 and 1 . Thus,

$$-2y = -1 = \frac{y - 0}{y^2 - x_0} \Rightarrow y = \frac{1}{2} \quad \text{and} \quad \frac{1/2}{(1/4) - x_0} = -1 \Rightarrow \frac{1}{4} - x_0 = -\frac{1}{2} \Rightarrow x_0 = \frac{3}{4}.$$

The perpendicular normal lines are $y = -x + \frac{3}{4}$ and $y = x - \frac{3}{4}$.

80. (a) $\frac{x^2}{32} + \frac{y^2}{8} = 1$

$$\frac{2x}{32} + \frac{2yy'}{8} = 0 \Rightarrow y' = \frac{-x}{4y}$$


$$\text{At } (4, 2): y' = \frac{-4}{4(2)} = -\frac{1}{2}$$

Slope of normal line is 2.

$$y - 2 = 2(x - 4)$$

$$y = 2x - 6$$

(b)



(c) $\frac{x^2}{32} + \frac{(2x - 6)^2}{8} = 1$

$$x^2 + 4(4x^2 - 24x + 36) = 32$$

$$17x^2 - 96x + 112 = 0$$

$$(17x - 28)(x - 4) = 0 \Rightarrow x = 4, \frac{28}{17}$$

Second point: $\left(\frac{28}{17}, -\frac{46}{17}\right)$

Section 2.6 Related Rates

1. $y = \sqrt{x}$

$$\frac{dy}{dt} = \left(\frac{1}{2\sqrt{x}}\right) \frac{dx}{dt}$$

$$\frac{dx}{dt} = 2\sqrt{x} \frac{dy}{dt}$$

(a) When $x = 4$ and $dx/dt = 3$,

$$\frac{dy}{dt} = \frac{1}{2\sqrt{4}}(3) = \frac{3}{4}.$$

(b) When $x = 25$ and $dy/dt = 2$,

$$\frac{dx}{dt} = 2\sqrt{25}(2) = 20.$$

2. $y = 2(x^2 - 3x)$

$$\frac{dy}{dt} = (4x - 6) \frac{dx}{dt}$$

$$\frac{dx}{dt} = \frac{1}{4x - 6} \frac{dy}{dt}$$

(a) When $x = 3$ and

$$\frac{dx}{dt} = 2, \frac{dy}{dt} = [4(3) - 6](2) = 12.$$

(b) When $x = 1$ and

$$\frac{dy}{dt} = 5, \frac{dx}{dt} = \frac{1}{4(1) - 6}(5) = -\frac{5}{2}.$$

3. $xy = 4$

$$x \frac{dy}{dt} + y \frac{dx}{dt} = 0$$

$$\frac{dy}{dt} = \left(-\frac{y}{x} \right) \frac{dx}{dt}$$

$$\frac{dx}{dt} = \left(-\frac{x}{y} \right) \frac{dy}{dt}$$

(a) When $x = 8$, $y = 1/2$, and $dx/dt = 10$,

$$\frac{dy}{dt} = -\frac{1/2}{8}(10) = -\frac{5}{8}$$

(b) When $x = 1$, $y = 4$, and $dy/dt = -6$,

$$\frac{dx}{dt} = -\frac{1}{4}(-6) = \frac{3}{2}$$

5. $y = x^2 + 1$

$$\frac{dx}{dt} = 2$$

$$\frac{dy}{dt} = 2x \frac{dx}{dt}$$

(a) When $x = -1$,

$$\frac{dy}{dt} = 2(-1)(2) = -4 \text{ cm/sec.}$$

(b) When $x = 0$,

$$\frac{dy}{dt} = 2(0)(2) = 0 \text{ cm/sec.}$$

(c) When $x = 1$,

$$\frac{dy}{dt} = 2(1)(2) = 4 \text{ cm/sec.}$$

7. $y = \tan x$

$$\frac{dx}{dt} = 2$$

$$\frac{dy}{dt} = \sec^2 x \frac{dx}{dt}$$

(a) When $x = -\pi/3$,

$$\frac{dy}{dt} = (2)^2(2) = 8 \text{ cm/sec.}$$

(b) When $x = -\pi/4$,

$$\frac{dy}{dt} = (\sqrt{2})^2(2) = 4 \text{ cm/sec.}$$

(c) When $x = 0$,

$$\frac{dy}{dt} = (1)^2(2) = 2 \text{ cm/sec.}$$

4. $x^2 + y^2 = 25$

$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

$$\frac{dy}{dt} = \left(-\frac{x}{y} \right) \frac{dx}{dt}$$

$$\frac{dx}{dt} = \left(-\frac{y}{x} \right) \frac{dy}{dt}$$

(a) When $x = 3$, $y = 4$, and $dx/dt = 8$,

$$\frac{dy}{dt} = -\frac{3}{4}(8) = -6.$$

(b) When $x = 4$, $y = 3$, and $dy/dt = -2$,

$$\frac{dx}{dt} = -\frac{3}{4}(-2) = \frac{3}{2}.$$

6. $y = \frac{1}{1+x^2}$

$$\frac{dx}{dt} = 2$$

$$\frac{dy}{dt} = \left[\frac{-2x}{(1+x^2)^2} \right] \frac{dx}{dt}$$

(a) When $x = -2$,

$$\frac{dy}{dt} = \frac{-2(-2)(2)}{25} = \frac{8}{25} \text{ cm/sec.}$$

(b) When $x = 0$,

$$\frac{dy}{dt} = 0 \text{ cm/sec.}$$

(c) When $x = 2$,

$$\frac{dy}{dt} = \frac{-2(2)(2)}{25} = \frac{-8}{25} \text{ cm/sec.}$$

8. $y = \sin x$

$$\frac{dx}{dt} = 2$$

$$\frac{dy}{dt} = \cos x \frac{dx}{dt}$$

(a) When $x = \pi/6$,

$$\frac{dy}{dt} = \left(\cos \frac{\pi}{6} \right)(2) = \sqrt{3} \text{ cm/sec.}$$

(b) When $x = \pi/4$,

$$\frac{dy}{dt} = \left(\cos \frac{\pi}{4} \right)(2) = \sqrt{2} \text{ cm/sec.}$$

(c) When $x = \pi/3$,

$$\frac{dy}{dt} = \left(\cos \frac{\pi}{3} \right)(2) = 1 \text{ cm/sec.}$$

9. (a) $\frac{dx}{dt}$ negative $\Rightarrow \frac{dy}{dt}$ positive

(b) $\frac{dy}{dt}$ positive $\Rightarrow \frac{dx}{dt}$ negative

10. (a) $\frac{dx}{dt}$ negative $\Rightarrow \frac{dy}{dt}$ negative

(b) $\frac{dy}{dt}$ positive $\Rightarrow \frac{dx}{dt}$ positive

11. Yes, y changes at a constant rate.

$$\frac{dy}{dt} = a \cdot \frac{dx}{dt}$$

No, the rate dy/dt is a multiple of dx/dt .

13. $D = \sqrt{x^2 + y^2} = \sqrt{x^2 + (x^2 + 1)^2} = \sqrt{x^4 + 3x^2 + 1}$

$$\frac{dx}{dt} = 2$$

$$\begin{aligned}\frac{dD}{dt} &= \frac{1}{2}(x^4 + 3x^2 + 1)^{-1/2}(4x^3 + 6x)\frac{dx}{dt} \\ &= \frac{2x^3 + 3x}{\sqrt{x^4 + 3x^2 + 1}} \frac{dx}{dt} \\ &= \frac{4x^3 + 6x}{\sqrt{x^4 + 3x^2 + 1}}.\end{aligned}$$

15. $A = \pi r^2$

$$\frac{dr}{dt} = 3$$

$$\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$$

(a) When $r = 6$, $\frac{dA}{dt} = 2\pi(6)(3) = 36\pi \text{ cm}^2/\text{min.}$

(b) When $r = 24$, $\frac{dA}{dt} = 2\pi(24)(3) = 144\pi \text{ cm}^2/\text{min.}$

17. (a) $\sin \frac{\theta}{2} = \frac{(1/2)b}{s} \Rightarrow b = 2s \sin \frac{\theta}{2}$

$$\cos \frac{\theta}{2} = \frac{h}{s} \Rightarrow h = s \cos \frac{\theta}{2}$$

$$A = \frac{1}{2}bh = \frac{1}{2}\left(2s \sin \frac{\theta}{2}\right)\left(s \cos \frac{\theta}{2}\right)$$

$$= \frac{s^2}{2}\left(2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}\right) = \frac{s^2}{2} \sin \theta$$

(b) $\frac{dA}{dt} = \frac{s^2}{2} \cos \theta \frac{d\theta}{dt}$ where $\frac{d\theta}{dt} = \frac{1}{2} \text{ rad/min.}$

When $\theta = \frac{\pi}{6}$, $\frac{dA}{dt} = \frac{s^2}{2}\left(\frac{\sqrt{3}}{2}\right)\left(\frac{1}{2}\right) = \frac{\sqrt{3}s^2}{8}$.

When $\theta = \frac{\pi}{3}$, $\frac{dA}{dt} = \frac{s^2}{2}\left(\frac{1}{2}\right)\left(\frac{1}{2}\right) = \frac{s^2}{8}$.

12. Answers will vary. See page 149.

14. $D = \sqrt{x^2 + y^2} = \sqrt{x^2 + \sin^2 x}$

$$\frac{dx}{dt} = 2$$


$$\begin{aligned}\frac{dD}{dt} &= \frac{1}{2}(x^2 + \sin^2 x)^{-1/2}(2x + 2 \sin x \cos x)\frac{dx}{dt} \\ &= \frac{x + \sin x \cos x}{\sqrt{x^2 + \sin^2 x}} \frac{dx}{dt} \\ &= \frac{2x + 2 \sin x \cos x}{\sqrt{x^2 + \sin^2 x}}.\end{aligned}$$

16. $A = \pi r^2$

$$\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$$

If dr/dt is constant, dA/dt is not constant.

dA/dt depends on r and dr/dt .



(c) If $\frac{d\theta}{dt}$ is constant, $\frac{dA}{dt}$ is proportional to $\cos \theta$.

18. $V = \frac{4}{3}\pi r^3$

$$\frac{dr}{dt} = 2$$

$$\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

(a) When $r = 6$,

$$\frac{dV}{dt} = 4\pi(6)^2(2) = 288\pi \text{ in}^3/\text{min.}$$

When $r = 24$,

$$\frac{dV}{dt} = 4\pi(24)^2(2) = 4608\pi \text{ in}^3/\text{min.}$$

(b) If dr/dt is constant, dV/dt is proportional to r^2 .

20. $V = x^3$

$$\frac{dx}{dt} = 3$$

$$\frac{dV}{dt} = 3x^2 \frac{dx}{dt}$$

(a) When $x = 1$,

$$\frac{dV}{dt} = 3(1)^2(3) = 9 \text{ cm}^3/\text{sec.}$$

(b) When $x = 10$,

$$\frac{dV}{dt} = 3(10)^2(3) = 900 \text{ cm}^3/\text{sec.}$$

22. $V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi r^2(3r) = \pi r^3$

$$\frac{dr}{dt} = 2$$

$$\frac{dV}{dt} = 3\pi r^2 \frac{dr}{dt}$$

(a) When $r = 6$,

$$\frac{dV}{dt} = 3\pi(6)^2(2) = 216\pi \text{ in}^3/\text{min.}$$

(b) When $r = 24$,

$$\frac{dV}{dt} = 3\pi(24)^2(2) = 3456\pi \text{ in}^3/\text{min.}$$

19. $V = \frac{4}{3}\pi r^3, \frac{dV}{dt} = 800$

$$\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

$$\frac{dr}{dt} = \frac{1}{4\pi r^2} \left(\frac{dV}{dt} \right) = \frac{1}{4\pi r^2}(800)$$

(a) When $r = 30$,

$$\frac{dr}{dt} = \frac{1}{4\pi(30)^2}(800) = \frac{2}{9\pi} \text{ cm/min.}$$

(b) When $r = 60$,

$$\frac{dr}{dt} = \frac{1}{4\pi(60)^2}(800) = \frac{1}{18\pi} \text{ cm/min.}$$

21. $s = 6x^2$

$$\frac{dx}{dt} = 3$$

$$\frac{ds}{dt} = 12x \frac{dx}{dt}$$

(a) When $x = 1$,

$$\frac{ds}{dt} = 12(1)(3) = 36 \text{ cm}^2/\text{sec.}$$

(b) When $x = 10$,

$$\frac{ds}{dt} = 12(10)(3) = 360 \text{ cm}^2/\text{sec.}$$

23. $V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi \left(\frac{9}{4}h^2\right)h$ [since $2r = 3h$]


$$= \frac{3\pi}{4}h^3$$

$$\frac{dV}{dt} = 10$$

$$\frac{dV}{dt} = \frac{9\pi}{4}h^2 \frac{dh}{dt} \Rightarrow \frac{dh}{dt} = \frac{4(dV/dt)}{9\pi h^2}$$

When $h = 15$,

$$\frac{dh}{dt} = \frac{4(10)}{9\pi(15)^2} = \frac{8}{405\pi} \text{ ft/min.}$$




24. $V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi \frac{25}{144} h^3 = \frac{25\pi}{3(144)} h^3$ (By similar triangles, $\frac{r}{5} = \frac{h}{12} \Rightarrow r = \frac{5}{12}h$)


$$\frac{dV}{dt} = 10$$

$$\frac{dV}{dt} = \frac{25\pi}{144} h^2 \frac{dh}{dt} \Rightarrow \frac{dh}{dt} = \left(\frac{144}{25\pi h^2}\right) \frac{dV}{dt}$$

When $h = 8$, $\frac{dh}{dt} = \frac{144}{25\pi(64)}(10) = \frac{9}{10\pi}$ ft/min.



25.



(a) Total volume of pool = $\frac{1}{2}(2)(12)(6) + (1)(6)(12) = 144 \text{ m}^3$


Volume of 1 m of water = $\frac{1}{2}(1)(6)(6) = 18 \text{ m}^3$ (see similar triangle diagram)

% pool filled = $\frac{18}{144}(100\%) = 12.5\%$

(b) Since for $0 \leq h \leq 2$, $b = 6h$, you have

$$V = \frac{1}{2}bh(6) = 3bh = 3(6h)h = 18h^2$$


$$\frac{dV}{dt} = 36h \frac{dh}{dt} = \frac{1}{4} \Rightarrow \frac{dh}{dt} = \frac{1}{144h} = \frac{1}{144(1)} = \frac{1}{144} \text{ m/min.}$$



26. $V = \frac{1}{2}bh(12) = 6bh = 6h^2$ (since $b = h$)

(a) $\frac{dV}{dt} = 12h \frac{dh}{dt} \Rightarrow \frac{dh}{dt} = \frac{1}{12h} \frac{dV}{dt}$

When $h = 1$ and $\frac{dV}{dt} = 2$, $\frac{dh}{dt} = \frac{1}{12(1)}(2) = \frac{1}{6}$ ft/min.



(b) If $\frac{dh}{dt} = \frac{3}{8}$ in./min = $\frac{1}{32}$ ft/min and $h = 2$ feet, then $\frac{dV}{dt} = (12)(2)\left(\frac{1}{32}\right) = \frac{3}{4} \text{ ft}^3/\text{min.}$

27. $x^2 + y^2 = 25^2$


$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

$$\frac{dy}{dt} = \frac{-x}{y} \cdot \frac{dx}{dt} = \frac{-2x}{y} \text{ since } \frac{dx}{dt} = 2.$$

(a) When $x = 7$, $y = \sqrt{576} = 24$, $\frac{dy}{dt} = \frac{-2(7)}{24} = \frac{-7}{12}$ ft/sec.

When $x = 15$, $y = \sqrt{400} = 20$, $\frac{dy}{dt} = \frac{-2(15)}{20} = \frac{-3}{2}$ ft/sec.

When $x = 24$, $y = 7$, $\frac{dy}{dt} = \frac{-2(24)}{7} = \frac{-48}{7}$ ft/sec.



—CONTINUED—

27. —CONTINUED—

(b) $A = \frac{1}{2}xy$

$$\frac{dA}{dt} = \frac{1}{2}\left(x \frac{dy}{dt} + y \frac{dx}{dt}\right)$$


From part (a) we have $x = 7$, $y = 24$, $\frac{dx}{dt} = 2$, and $\frac{dy}{dt} = -\frac{7}{12}$. Thus,

$$\frac{dA}{dt} = \frac{1}{2}\left[7\left(-\frac{7}{12}\right) + 24(2)\right] = \frac{527}{24} \approx 21.96 \text{ ft}^2/\text{sec.}$$

(c) $\tan \theta = \frac{x}{y}$

$$\sec^2 \theta \frac{d\theta}{dt} = \frac{1}{y} \cdot \frac{dx}{dt} - \frac{x}{y^2} \cdot \frac{dy}{dt}$$

$$\frac{d\theta}{dt} = \cos^2 \theta \left[\frac{1}{y} \cdot \frac{dx}{dt} - \frac{x}{y^2} \cdot \frac{dy}{dt} \right]$$



Using $x = 7$, $y = 24$, $\frac{dx}{dt} = 2$, $\frac{dy}{dt} = -\frac{7}{12}$ and $\cos \theta = \frac{24}{25}$, we have


$$\frac{d\theta}{dt} = \left(\frac{24}{25}\right)^2 \left[\frac{1}{24}(2) - \frac{7}{(24)^2} \left(-\frac{7}{12}\right) \right] = \frac{1}{12} \text{ rad/sec.}$$

28. $x^2 + y^2 = 25$

$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

$$\frac{dx}{dt} = -\frac{y}{x} \cdot \frac{dy}{dt} = -\frac{0.15y}{x} \quad \left(\text{since } \frac{dy}{dt} = 0.15\right)$$

When $x = 2.5$, $y = \sqrt{18.75}$, $\frac{dx}{dt} = -\frac{\sqrt{18.75}}{2.5} 0.15 \approx -0.26 \text{ m/sec.}$



29. When $y = 6$, $x = \sqrt{12^2 - 6^2} = 6\sqrt{3}$, and $s = \sqrt{x^2 + (12-y)^2}$
 $= \sqrt{108 + 36} = 12$.

$$x^2 + (12-y)^2 = s^2$$

$$2x \frac{dx}{dt} + 2(12-y)(-1) \frac{dy}{dt} = 2s \frac{ds}{dt}$$

$$x \frac{dx}{dt} + (y-12) \frac{dy}{dt} = s \frac{ds}{dt}$$


Also, $x^2 + y^2 = 12^2$.

$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0 \Rightarrow \frac{dy}{dt} = -\frac{x}{y} \frac{dx}{dt}$$

$$\text{Thus, } x \frac{dx}{dt} + (y-12) \left(\frac{-x}{y} \frac{dx}{dt} \right) = s \frac{ds}{dt}$$

$$\frac{dx}{dt} \left[x - x + \frac{12x}{y} \right] = s \frac{ds}{dt} \Rightarrow \frac{dx}{dt} = \frac{sy}{12x} \cdot \frac{ds}{dt} = \frac{(12)(6)}{(12)(6\sqrt{3})} (-0.2) = \frac{-1}{5\sqrt{3}} = \frac{-\sqrt{3}}{15} \text{ m/sec (horizontal)}$$

$$\frac{dy}{dt} = \frac{-x}{y} \frac{dx}{dt} = \frac{-6\sqrt{3}}{6} \cdot \frac{(-\sqrt{3})}{15} = \frac{1}{5} \text{ m/sec (vertical)}$$



30. Let L be the length of the rope.

$$(a) L^2 = 144 + x^2$$

$$2L \frac{dL}{dt} = 2x \frac{dx}{dt}$$


$$\frac{dx}{dt} = \frac{L}{x} \cdot \frac{dL}{dt} = -\frac{4L}{x} \quad \left(\text{since } \frac{dL}{dt} = -4 \text{ ft/sec} \right)$$

When $L = 13$:

$$x = \sqrt{L^2 - 144} = \sqrt{169 - 144} = 5$$

$$\frac{dx}{dt} = -\frac{4(13)}{5} = -\frac{52}{5} = -10.4 \text{ ft/sec}$$

Speed of the boat increases as it approaches the dock.



(b) If $\frac{dx}{dt} = -4$, and $L = 13$:

$$\frac{dL}{dt} = \frac{x}{L} \frac{dx}{dt} = \frac{5}{13}(-4) = -\frac{20}{13} \text{ ft/sec}$$

$$\frac{dL}{dt} = \frac{x}{L} \frac{dx}{dt} = \frac{\sqrt{L^2 - 144}}{L}(-4)$$

$$\lim_{L \rightarrow 12^+} \frac{dL}{dt} = \lim_{L \rightarrow 12^+} \frac{-4}{L} \sqrt{L^2 - 144} = 0$$

$$31. (a) s^2 = x^2 + y^2$$

$$\frac{dx}{dt} = -450$$


$$\frac{dy}{dt} = -600$$

$$2s \frac{ds}{dt} = 2x \frac{dx}{dt} + 2y \frac{dy}{dt}$$

$$\frac{ds}{dt} = \frac{x(dx/dt) + y(dy/dt)}{s}$$

$$\text{When } x = 150 \text{ and } y = 200, s = 250 \text{ and } \frac{ds}{dt} = \frac{150(-450) + 200(-600)}{250} = -750 \text{ mph.}$$

$$(b) t = \frac{250}{750} = \frac{1}{3} \text{ hr} = 20 \text{ min}$$




$$32. x^2 + y^2 = s^2$$

$$2x \frac{dx}{dt} + 0 = 2s \frac{ds}{dt} \quad \left(\text{since } \frac{dy}{dt} = 0 \right)$$

$$\frac{dx}{dt} = \frac{s}{x} \frac{ds}{dt}$$

$$\text{When } s = 10, x = \sqrt{100 - 25} = \sqrt{75} = 5\sqrt{3},$$

$$\frac{dx}{dt} = \frac{10}{5\sqrt{3}}(240) = \frac{480}{\sqrt{3}} = 160\sqrt{3} \approx 277.13 \text{ mph.}$$



$$33. s^2 = 90^2 + x^2$$


$$x = 30$$

$$\frac{dx}{dt} = -28$$

$$2s \frac{ds}{dt} = 2x \frac{dx}{dt} \Rightarrow \frac{ds}{dt} = \frac{x}{s} \cdot \frac{dx}{dt}$$

$$\text{When } x = 30, s = \sqrt{90^2 + 30^2} = 30\sqrt{10},$$

$$\frac{ds}{dt} = \frac{30}{30\sqrt{10}}(-28) = \frac{-28}{\sqrt{10}} \approx -8.85 \text{ ft/sec.}$$



34. $s^2 = 90^2 + x^2$


$x = 60$

$\frac{dx}{dt} = 28$

$\frac{ds}{dt} = \frac{x}{s} \cdot \frac{dx}{dt}$

When $x = 60$, $s = \sqrt{90^2 + 60^2} = 30\sqrt{13}$,

$\frac{ds}{dt} = \frac{60}{30\sqrt{13}}(28) = \frac{56}{\sqrt{13}} \approx 15.53 \text{ ft/sec.}$



36. (a) $\frac{20}{6} = \frac{y}{y-x}$

$20y - 20x = 6y$

$14y = 20x$

$y = \frac{10}{7}x$

$\frac{dx}{dt} = -5$

$\frac{dy}{dt} = \frac{10}{7} \frac{dx}{dt} = \frac{10}{7}(-5) = -\frac{50}{7} \text{ ft/sec}$

(b) $\frac{d(y-x)}{dt} = \frac{dy}{dt} - \frac{dx}{dt} = -\frac{50}{7} - (-5) = -\frac{50}{7} + \frac{35}{7} = -\frac{15}{7} \text{ ft/sec}$

37. $x(t) = \frac{1}{2} \sin \frac{\pi t}{6}, x^2 + y^2 = 1$

(a) Period: $\frac{2\pi}{\pi/6} = 12 \text{ seconds}$

(b) When $x = \frac{1}{2}$, $y = \sqrt{1^2 - \left(\frac{1}{2}\right)^2} = \frac{\sqrt{3}}{2} \text{ m.}$


Lowest point: $\left(0, \frac{\sqrt{3}}{2}\right)$

35. (a) $\frac{15}{6} = \frac{y}{y-x} \Rightarrow 15y - 15x = 6y$


$y = \frac{5}{3}x$

$\frac{dx}{dt} = 5$

$\frac{dy}{dt} = \frac{5}{3} \cdot \frac{dx}{dt} = \frac{5}{3}(5) = \frac{25}{3} \text{ ft/sec}$



(b) $\frac{d(y-x)}{dt} = \frac{dy}{dt} - \frac{dx}{dt} = \frac{25}{3} - 5 = \frac{10}{3} \text{ ft/sec}$



(c) When $x = \frac{1}{4}$, $y = \sqrt{1 - \left(\frac{1}{4}\right)^2} = \frac{\sqrt{15}}{4}$ and $t = 1$:

$\frac{dx}{dt} = \frac{1}{2} \left(\frac{\pi}{6}\right) \cos \frac{\pi t}{6} = \frac{\pi}{12} \cos \frac{\pi t}{6}$

$x^2 + y^2 = 1$

$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0 \Rightarrow \frac{dy}{dt} = -\frac{x}{y} \frac{dx}{dt}$

Thus, $\frac{dy}{dt} = -\frac{1/4}{\sqrt{15}/4} \cdot \frac{\pi}{12} \cos\left(\frac{\pi}{6}\right)$

$= -\frac{\pi}{\sqrt{15}} \left(\frac{1}{12}\right) \frac{\sqrt{3}}{2} = \frac{-\pi}{24} \frac{1}{\sqrt{5}} = -\frac{\sqrt{5}\pi}{120}$

Speed = $\left| \frac{-\sqrt{5}\pi}{120} \right| = \frac{\sqrt{5}\pi}{120} \text{ m/sec}$

38. $x(t) = \frac{3}{5} \sin \pi t, x^2 + y^2 = 1$

(a) Period: $\frac{2\pi}{\pi} = 2$ seconds

(b) When $x = \frac{3}{5}$, $y = \sqrt{1 - \left(\frac{3}{5}\right)^2} = \frac{4}{5}$ m.

Lowest point: $\left(0, -\frac{4}{5}\right)$

(c) When $x = \frac{3}{10}$, $y = \sqrt{1 - \left(\frac{1}{4}\right)^2} = \frac{\sqrt{15}}{4}$ and $\frac{3}{10} = \frac{3}{5} \sin \pi t \Rightarrow \sin \pi t = \frac{1}{2} \Rightarrow t = \frac{1}{6}$.

$$\frac{dx}{dt} = \frac{3}{5} \pi \cos \pi t$$

$$x^2 + y^2 = 1$$

$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0 \Rightarrow \frac{dy}{dt} = \frac{-x}{y} \frac{dx}{dt}$$

$$\begin{aligned} \text{Thus, } \frac{dy}{dt} &= \frac{-3/10}{\sqrt{15}/4} \cdot \frac{3}{5} \pi \cos\left(\frac{\pi}{6}\right) \\ &= \frac{-9\pi}{25\sqrt{5}} = \frac{-9\sqrt{5}\pi}{125}. \end{aligned}$$

$$\text{Speed} = \left| \frac{-9\sqrt{5}\pi}{125} \right| \approx 0.5058 \text{ m/sec}$$

39. Since the evaporation rate is proportional to the surface area, $dV/dt = k(4\pi r^2)$. However, since $V = (4/3)\pi r^3$, we have

$$\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

$$\text{Therefore, } k(4\pi r^2) = 4\pi r^2 \frac{dr}{dt} \Rightarrow k = \frac{dr}{dt}$$

40. $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2}$

$$\frac{dR_1}{dt} = 1$$

$$\frac{dR_2}{dt} = 1.5$$

$$\frac{1}{R^2} \cdot \frac{dR}{dt} = \frac{1}{R_1^2} \cdot \frac{dR_1}{dt} + \frac{1}{R_2^2} \cdot \frac{dR_2}{dt}$$

When $R_1 = 50$ and $R_2 = 75$:

$$R = 30$$

$$\frac{dR}{dt} = (30)^2 \left[\frac{1}{(50)^2}(1) + \frac{1}{(75)^2}(1.5) \right] = 0.6 \text{ ohms/sec}$$

41. $pV^{1.3} = k$

$$1.3pV^{0.3} \frac{dV}{dt} + V^{1.3} \frac{dp}{dt} = 0$$

$$V^{0.3} \left(1.3p \frac{dV}{dt} + V \frac{dp}{dt} \right) = 0$$

$$1.3p \frac{dV}{dt} = -V \frac{dp}{dt}$$

42. $rg \tan \theta = v^2$

$$32r \tan \theta = v^2, \quad r \text{ is a constant.}$$

$$32r \sec^2 \theta \frac{d\theta}{dt} = 2v \frac{dv}{dt}$$

$$\frac{dv}{dt} = \frac{16r}{v} \sec^2 \theta \frac{d\theta}{dt}$$

$$\text{Likewise, } \frac{d\theta}{dt} = \frac{v}{16r} \cos^2 \theta \frac{dv}{dt}$$


43. $\tan \theta = \frac{y}{30}$

$$\frac{dy}{dt} = 3 \text{ m/sec}$$

$$\sec^2 \theta \cdot \frac{d\theta}{dt} = \frac{1}{30} \frac{dy}{dt}$$

$$\frac{d\theta}{dt} = \frac{1}{30} \cos^2 \theta \cdot \frac{dy}{dt}$$

When $y = 30$, $\theta = \frac{\pi}{4}$, and $\cos \theta = \frac{\sqrt{2}}{2}$. Thus, $\frac{d\theta}{dt} = \frac{1}{30} \left(\frac{1}{2}\right)(3) = \frac{1}{20} \text{ rad/sec.}$




44. $\sin \theta = \frac{10}{x}$

$$\frac{dx}{dt} = (-1) \text{ ft/sec}$$

$$\cos \theta \left(\frac{d\theta}{dt} \right) = \frac{-10}{x^2} \cdot \frac{dx}{dt}$$

$$\frac{d\theta}{dt} = \frac{-10}{x^2} \frac{dx}{dt} (\sec \theta)$$

$$= \frac{-10}{25^2} (-1) \frac{25}{\sqrt{25^2 - 10^2}} = \frac{10}{25} \frac{1}{5\sqrt{21}} = \frac{2}{25\sqrt{21}} = \frac{2\sqrt{21}}{525} \approx 0.017 \text{ rad/sec}$$




45. $\tan \theta = \frac{y}{x}, y = 5$

$$\frac{dx}{dt} = -600 \text{ mi/hr}$$

$$(\sec^2 \theta) \frac{d\theta}{dt} = -\frac{5}{x^2} \cdot \frac{dx}{dt}$$

$$\frac{d\theta}{dt} = \cos^2 \theta \left(-\frac{5}{x^2} \right) \frac{dx}{dt} = \frac{x^2}{L^2} \left(-\frac{5}{x^2} \right) \frac{dx}{dt}$$

$$= \left(-\frac{5^2}{L^2} \right) \left(\frac{1}{5} \right) \frac{dx}{dt} = (-\sin^2 \theta) \left(\frac{1}{5} \right) (-600) = 120 \sin^2 \theta$$



(a) When $\theta = 30^\circ$,

$$\frac{d\theta}{dt} = \frac{120}{4}$$

$$= 30 \text{ rad/hr} = \frac{1}{2} \text{ rad/min.}$$

(b) When $\theta = 60^\circ$,

$$\frac{d\theta}{dt} = 120 \left(\frac{3}{4} \right)$$

$$= 90 \text{ rad/hr} = \frac{3}{2} \text{ rad/min.}$$

(c) When $\theta = 75^\circ$,

$$\frac{d\theta}{dt} = 120 \sin^2 75^\circ$$

$$\approx 111.96 \text{ rad/hr} \approx 1.87 \text{ rad/min.}$$


46. $\tan \theta = \frac{x}{50}$

$$\frac{d\theta}{dt} = 30(2\pi) = 60\pi \text{ rad/min} = \pi \text{ rad/sec}$$

$$\sec^2 \theta \left(\frac{d\theta}{dt} \right) = \frac{1}{50} \left(\frac{dx}{dt} \right)$$

$$\frac{dx}{dt} = 50 \sec^2 \theta \left(\frac{d\theta}{dt} \right)$$

(a) When $\theta = 30^\circ$, $\frac{dx}{dt} = \frac{200\pi}{3}$ ft/sec. (b) When $\theta = 60^\circ$, $\frac{dx}{dt} = 200\pi$ ft/sec. (c) When $\theta = 70^\circ$, $\frac{dx}{dt} \approx 427.43\pi$ ft/sec.



47. $\frac{d\theta}{dt} = (10 \text{ rev/sec})(2\pi \text{ rad/rev}) = 20\pi \text{ rad/sec}$

(a) $\cos \theta = \frac{x}{30}$

$$-\sin \theta \frac{d\theta}{dt} = \frac{1}{30} \frac{dx}{dt}$$

$$\frac{dx}{dt} = -30 \sin \theta \frac{d\theta}{dt}$$



$$= -30 \sin \theta (20\pi)$$

$$= -600\pi \sin \theta$$

(c) $|dx/dt| = |-600\pi \sin \theta|$ is greatest when

$$|\sin \theta| = 1 \Rightarrow \theta = \frac{\pi}{2} + n\pi \quad (\text{or } 90^\circ + n \cdot 180^\circ).$$

$|dx/dt|$ is least when $\theta = n\pi$ (or $n \cdot 180^\circ$).



(d) For $\theta = 30^\circ$,

$$\frac{dx}{dt} = -600\pi \sin(30^\circ)$$

$$= -600\pi \frac{1}{2} = -300\pi \text{ cm/sec.}$$

For $\theta = 60^\circ$,


$$\frac{dx}{dt} = -600\pi \sin(60^\circ)$$

$$= -600\pi \frac{\sqrt{3}}{2} = -300\sqrt{3}\pi \text{ cm/sec.}$$

48. $\sin 22^\circ = \frac{x}{y}$

$$0 = -\frac{x}{y^2} \cdot \frac{dy}{dt} + \frac{1}{y} \cdot \frac{dx}{dt}$$

$$\frac{dx}{dt} = \frac{x}{y} \cdot \frac{dy}{dt} = (\sin 22^\circ)(240) \approx 89.9056 \text{ mi/hr}$$



49. $\tan \theta = \frac{x}{50} \Rightarrow x = 50 \tan \theta$

$$\frac{dx}{dt} = 50 \sec^2 \theta \frac{d\theta}{dt}$$

$$2 = 50 \sec^2 \theta \frac{d\theta}{dt}$$

$$\frac{d\theta}{dt} = \frac{1}{25} \cos^2 \theta, \quad -\frac{\pi}{4} \leq \theta \leq \frac{\pi}{4}$$

50. (a) $dy/dt = 3(dx/dt)$ means that y changes three times as fast as x changes.

(b) y changes slowly when $x \approx 0$ or $x \approx L$. y changes more rapidly when x is near the middle of the interval.

51. $x^2 + y^2 = 25$; acceleration of the top of the ladder $= \frac{d^2y}{dt^2}$

First derivative: $2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$

$$x \frac{dx}{dt} + y \frac{dy}{dt} = 0$$

Second derivative: $x \frac{d^2x}{dt^2} + \frac{dx}{dt} \cdot \frac{dx}{dt} + y \frac{d^2y}{dt^2} + \frac{dy}{dt} \cdot \frac{dy}{dt} = 0$

$$\frac{d^2y}{dt^2} = \left(\frac{1}{y}\right) \left[-x \frac{d^2x}{dt^2} - \left(\frac{dx}{dt}\right)^2 - \left(\frac{dy}{dt}\right)^2 \right]$$

When $x = 7$, $y = 24$, $\frac{dy}{dt} = -\frac{7}{12}$, and $\frac{dx}{dt} = 2$ (see Exercise 27). Since $\frac{dx}{dt}$ is constant, $\frac{d^2x}{dt^2} = 0$.

$$\frac{d^2y}{dt^2} = \frac{1}{24} \left[-7(0) - (2)^2 - \left(-\frac{7}{12}\right)^2 \right] = \frac{1}{24} \left[-4 - \frac{49}{144} \right] = \frac{1}{24} \left[-\frac{625}{144} \right] \approx -0.1808 \text{ ft/sec}^2$$

52. $L^2 = 144 + x^2$; acceleration of the boat $= \frac{d^2x}{dt^2}$

First derivative: $2L \frac{dL}{dt} = 2x \frac{dx}{dt}$

$$L \frac{dL}{dt} = x \frac{dx}{dt}$$

Second derivative: $L \frac{d^2L}{dt^2} + \frac{dL}{dt} \cdot \frac{dL}{dt} = x \frac{d^2x}{dt^2} + \frac{dx}{dt} \cdot \frac{dx}{dt}$

$$\frac{d^2x}{dt^2} = \left(\frac{1}{x}\right) \left[L \frac{d^2L}{dt^2} + \left(\frac{dL}{dt}\right)^2 - \left(\frac{dx}{dt}\right)^2 \right]$$

When $L = 13$, $x = 5$, $\frac{dx}{dt} = -10.4$, and $\frac{dL}{dt} = -4$ (see Exercise 30). Since $\frac{dL}{dt}$ is constant, $\frac{d^2L}{dt^2} = 0$.

$$\frac{d^2x}{dt^2} = \frac{1}{5} [13(0) + (-4)^2 - (-10.4)^2]$$

$$= \frac{1}{5}[16 - 108.16] = \frac{1}{5}[-92.16] = -18.432 \text{ ft/sec}^2$$

53. (a) $m(s) = 0.3754s^3 - 18.780s^2 + 313.23s - 1707.8$

(b) $\frac{dm}{dt} = (1.1262s^2 - 37.560s + 313.23) \frac{ds}{dt}$

If $t = 10$ and $\frac{ds}{dt} = 0.75$, then $s = 17.8$ and $\frac{dm}{dt} \approx 1.1154$ million/year.

54. $y(t) = -4.9t^2 + 20$

$$\frac{dy}{dt} = -9.8t$$

$$y(1) = -4.9 + 20 = 15.1$$

$$y'(1) = -9.8$$

By similar triangles: $\frac{20}{x} = \frac{y}{x - 12}$

$$20x - 240 = xy$$

When $y = 15.1$: $20x - 240 = x(15.1)$

$$(20 - 15.1)x = 240$$


$$x = \frac{240}{4.9}$$

$$20x - 240 = xy$$

$$20 \frac{dx}{dt} = x \frac{dy}{dt} + y \frac{dx}{dt}$$

$$\frac{dx}{dt} = \frac{x}{20-y} \frac{dy}{dt}$$

At $t = 1$, $\frac{dx}{dt} = \frac{240/4.9}{20 - 15.1}(-9.8) \approx -97.96 \text{ m/sec.}$



Review Exercises for Chapter 2

1. $f(x) = x^2 - 2x + 3$

$$\begin{aligned}
 f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{[(x + \Delta x)^2 - 2(x + \Delta x) + 3] - [x^2 - 2x + 3]}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{(x^2 + 2x(\Delta x) + (\Delta x)^2 - 2x - 2(\Delta x) + 3) - (x^2 - 2x + 3)}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{2x(\Delta x) + (\Delta x)^2 - 2(\Delta x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} (2x + \Delta x - 2) = 2x - 2
 \end{aligned}$$

2. $f(x) = \sqrt{x} + 1$

$$\begin{aligned}
 f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{(\sqrt{x + \Delta x} + 1) - (\sqrt{x} + 1)}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{\sqrt{x + \Delta x} - \sqrt{x}}{\Delta x} \cdot \frac{\sqrt{x + \Delta x} + \sqrt{x}}{\sqrt{x + \Delta x} + \sqrt{x}} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x) - x}{\Delta x(\sqrt{x + \Delta x} + \sqrt{x})} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{1}{\sqrt{x + \Delta x} + \sqrt{x}} = \frac{1}{2\sqrt{x}}
 \end{aligned}$$

3. $f(x) = \frac{x+1}{x-1}$

$$\begin{aligned}
 f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\frac{x + \Delta x + 1}{x + \Delta x - 1} - \frac{x+1}{x-1}}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x + 1)(x - 1) - (x + \Delta x - 1)(x + 1)}{\Delta x(x + \Delta x - 1)(x - 1)} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{(x^2 + x \Delta x + x - x - \Delta x - 1) - (x^2 + x \Delta x - x + x + \Delta x - 1)}{\Delta x(x + \Delta x - 1)(x - 1)} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{-2 \Delta x}{\Delta x(x + \Delta x - 1)(x - 1)} = \lim_{\Delta x \rightarrow 0} \frac{-2}{(x + \Delta x - 1)(x - 1)} = \frac{-2}{(x - 1)^2}
 \end{aligned}$$

4. $f(x) = \frac{2}{x}$


$$\begin{aligned}
 f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\frac{2}{x + \Delta x} - \frac{2}{x}}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{2x - (2x + 2 \Delta x)}{\Delta x(x + \Delta x)x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{-2 \Delta x}{\Delta x(x + \Delta x)x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{-2}{(x + \Delta x)x} = \frac{-2}{x^2}
 \end{aligned}$$

5. f is differentiable for all $x \neq -1$.

7. $f(x) = 4 - |x - 2|$

(a) Continuous at $x = 2$

(b) Not differentiable at $x = 2$ because of the sharp turn in the graph.




9. Using the limit definition, you obtain $g'(x) = \frac{4}{3}x - \frac{1}{6}$. At $x = -1$,

$$g'(-1) = -\frac{4}{3} - \frac{1}{6} = -\frac{3}{2}.$$

11. (a) Using the limit definition, $f'(x) = 3x^2$. At $x = -1$, $f'(-1) = 3$. The tangent line is

$$y - (-2) = 3(x - (-1))$$

$$y = 3x + 1.$$



$$\begin{aligned} 13. g'(2) &= \lim_{x \rightarrow 2} \frac{g(x) - g(2)}{x - 2} \\ &= \lim_{x \rightarrow 2} \frac{x^2(x - 1) - 4}{x - 2} \\ &= \lim_{x \rightarrow 2} \frac{x^3 - x^2 - 4}{x - 2} \\ &= \lim_{x \rightarrow 2} \frac{(x - 2)(x^2 + x + 2)}{x - 2} \\ &= \lim_{x \rightarrow 2} (x^2 + x + 2) = 8 \end{aligned}$$

15. $y = 25$

$$y' = 0$$

16. $y = -12$


$$y' = 0$$

6. f is differentiable for all $x \neq -3$.

8. $f(x) = \begin{cases} x^2 + 4x + 2, & \text{if } x < -2 \\ 1 - 4x - x^2, & \text{if } x \geq -2 \end{cases}$

(a) Nonremovable discontinuity at $x = -2$.

(b) Not differentiable at $x = -2$ because the function is discontinuous there.



10. Using the limit definition, you obtain $h'(x) = \frac{3}{8} - 4x$. At $x = -2$,

$$h'(-2) = \frac{3}{8} - 4(-2) = \frac{67}{8}.$$

12. (a) Using the limit definition,


$$f'(x) = \frac{-2}{(x + 1)^2}.$$

At $x = 0$, $f'(0) = -2$. The tangent line is

$$y - 2 = -2(x - 0)$$

$$y = -2x + 2.$$

(b)



$$\begin{aligned} 14. f'(2) &= \lim_{x \rightarrow 2} \frac{f(x) - f(2)}{x - 2} \\ &= \lim_{x \rightarrow 2} \frac{\frac{1}{x+1} - \frac{1}{3}}{x - 2} \\ &= \lim_{x \rightarrow 2} \frac{3 - x - 1}{(x - 2)(x + 1)3} \\ &= \lim_{x \rightarrow 2} \frac{-1}{(x + 1)3} = -\frac{1}{9} \end{aligned}$$

17. $f(x) = x^8$

$$f'(x) = 8x^7$$

18. $g(x) = x^{12}$

$$g'(x) = 12x^{11}$$

19. $h(t) = 3t^4$

$$h'(t) = 12t^3$$

20. $f(t) = -8t^5$

$$f'(t) = -40t^4$$

21. $f(x) = x^3 - 3x^2$

$$\begin{aligned}f'(x) &= 3x^2 - 6x \\&= 3x(x - 2)\end{aligned}$$

22. $g(s) = 4s^4 - 5s^2$

$$g'(s) = 16s^3 - 10s$$

23. $h(x) = 6\sqrt{x} + 3\sqrt[3]{x} = 6x^{1/2} + 3x^{1/3}$

$$h'(x) = 3x^{-1/2} + x^{-2/3} = \frac{3}{\sqrt{x}} + \frac{1}{\sqrt[3]{x^2}}$$

24. $f(x) = x^{1/2} - x^{-1/2}$

$$f'(x) = \frac{1}{2}x^{-1/2} + \frac{1}{2}x^{-3/2} = \frac{x+1}{2x^{3/2}}$$

25. $g(t) = \frac{2}{3}t^{-2}$

$$g'(t) = \frac{-4}{3}t^{-3} = \frac{-4}{3t^3}$$

26. $h(x) = \frac{2}{9}x^{-2}$

$$h'(x) = \frac{-4}{9}x^{-3} = \frac{-4}{9x^3}$$

27. $f(\theta) = 2\theta - 3 \sin \theta$

$$f'(\theta) = 2 - 3 \cos \theta$$

28. $g(\alpha) = 4 \cos \alpha + 6$

$$g'(\alpha) = -4 \sin \alpha$$


29. $f(\theta) = 3 \cos \theta - \frac{\sin \theta}{4}$

$$f'(\theta) = -3 \sin \theta - \frac{\cos \theta}{4}$$


30. $g(\alpha) = \frac{5 \sin \alpha}{3} - 2\alpha$

$$g'(\alpha) = \frac{5 \cos \alpha}{3} - 2$$

31.



32.

33. $F = 200\sqrt{T}$

$$F'(t) = \frac{100}{\sqrt{T}}$$

(a) When $T = 4$,
 $F'(4) = 50$ vibrations/sec/lb.(b) When $T = 9$,
 $F'(9) = 33\frac{1}{3}$ vibrations/sec/lb.

34. $s = -16t^2 + s_0$

First ball:

$$-16t^2 + 100 = 0$$

$$t = \sqrt{\frac{100}{16}} = \frac{10}{4} = 2.5 \text{ seconds to hit ground}$$

Second ball:

$$-16t^2 + 75 = 0$$

$$t^2 = \sqrt{\frac{75}{16}} = \frac{5\sqrt{3}}{4} \approx 2.165 \text{ seconds to hit ground}$$

Since the second ball was released one second after the first ball, the first ball will hit the ground first.

The second ball will hit the ground $3.165 - 2.5 = 0.665$ second later.

35. $s(t) = -16t^2 + s_0$

$$s(9.2) = -16(9.2)^2 + s_0 = 0$$

$$s_0 = 1354.24$$

The building is approximately 1354 feet high (or 415 m).


36. $s(t) = -16t^2 + 14,400 = 0$

$$16t^2 = 14,400$$

$$t = 30 \text{ sec}$$

Since $600 \text{ mph} = \frac{1}{6} \text{ mi/sec}$, in 30 seconds the bomb will move horizontally $(\frac{1}{6})(30) = 5$ miles.

37. (a)



Total horizontal distance: 50

(b) $0 = x - 0.02x^2$

$0 = x\left(1 - \frac{x}{50}\right)$ implies $x = 50$.

(c) Ball reaches maximum height when $x = 25$.

(d) $y = x - 0.02x^2$

$y' = 1 - 0.04x$

$y'(0) = 1$

$y'(10) = 0.6$


$y'(25) = 0$

$y'(30) = -0.2$

$y'(50) = -1$

(e) $y'(25) = 0$

38.



(a) $y = x - \frac{32}{v_0^2}x^2 = x\left(1 - \frac{32}{v_0^2}x\right)$

= 0 if $x = 0$ or $x = \frac{v_0^2}{32}$

Projectile strikes the ground when $x = v_0^2/32$.Projectile reaches its maximum height at $x = v_0^2/64$ (one-half the distance).

(c) $y = x - \frac{32}{v_0^2}x^2 = x\left(1 - \frac{32}{v_0^2}x\right) = 0$

when $x = 0$ and $x = s_0^2/32$. Therefore, the range is $x = v_0^2/32$. When the initial velocity is doubled the range is

$x = \frac{(2v_0)^2}{32} = \frac{4v_0^2}{32}$

or four times the initial range. From part (a), the maximum height occurs when $x = v_0^2/64$. The maximum height is

$y\left(\frac{v_0^2}{64}\right) = \frac{v_0^2}{64} - \frac{32}{v_0^2}\left(\frac{v_0^2}{64}\right)^2 = \frac{v_0^2}{64} - \frac{v_0^2}{128} = \frac{v_0^2}{128}$

If the initial velocity is doubled, the maximum height is

$y\left[\frac{(2v_0)^2}{64}\right] = \frac{(2v_0)^2}{128} = 4\left(\frac{v_0^2}{128}\right)$

or four times the original maximum height.


(b) $y' = 1 - \frac{64}{v_0^2}x$

When $x = \frac{v_0^2}{64}$, $y' = 1 - \frac{64}{v_0^2}\left(\frac{v_0^2}{64}\right) = 0$.

(d) $v_0 = 70$ ft/sec

Range: $x = \frac{v_0^2}{32} = \frac{(70)^2}{32} = 153.125$ ft

Maximum height: $y = \frac{v_0^2}{128} = \frac{(70)^2}{128} \approx 38.28$ ft



39. $x(t) = t^2 - 3t + 2 = (t-2)(t-1)$

(a) $v(t) = x'(t) = 2t - 3$

(c) $v(t) = 0$ for $t = \frac{3}{2}$

$$x = \left(\frac{3}{2} - 2\right)\left(\frac{3}{2} - 1\right) = \left(-\frac{1}{2}\right)\left(\frac{1}{2}\right) = -\frac{1}{4}$$

(b) $v(t) < 0$ for $t < \frac{3}{2}$

(d) $x(t) = 0$ for $t = 1, 2$


$$|v(1)| = |2(1) - 3| = 1$$

$$|v(2)| = |2(2) - 3| = 1$$


The speed is 1 when the position is 0.

40. (a) $y = 0.14x^2 - 4.43x + 58.4$

(c)



(b)



(d) If $x = 65$, $y \approx 362$ feet.

- (e) As the speed increases, the stopping distance increases at an increasing rate.

41. $f(x) = (3x^2 + 7)(x^2 - 2x + 3)$

$$\begin{aligned} f'(x) &= (3x^2 + 7)(2x - 2) + (x^2 - 2x + 3)(6x) \\ &= 2(6x^3 - 9x^2 + 16x - 7) \end{aligned}$$

42. $g(x) = (x^3 - 3x)(x + 2)$

$$\begin{aligned} g'(x) &= (x^3 - 3x)(1) + (x + 2)(3x^2 - 3) \\ &= x^3 - 3x + 3x^3 + 6x^2 - 3x - 6 \\ &= 4x^3 + 6x^2 - 6x - 6 \end{aligned}$$

43. $h(x) = \sqrt{x} \sin x = x^{1/2} \sin x$

$$h'(x) = \frac{1}{2\sqrt{x}} \sin x + \sqrt{x} \cos x$$

44. $f(t) = t^3 \cos t$

$$\begin{aligned} f'(t) &= t^3(-\sin t) + \cos t(3t^2) \\ &= -t^3 \sin t + 3t^2 \cos t \end{aligned}$$

45. $f(x) = \frac{x^2 + x - 1}{x^2 - 1}$

$$\begin{aligned} f'(x) &= \frac{(x^2 - 1)(2x + 1) - (x^2 + x - 1)(2x)}{(x^2 - 1)^2} \\ &= \frac{-(x^2 + 1)}{(x^2 - 1)^2} \end{aligned}$$

46. $f(x) = \frac{6x - 5}{x^2 + 1}$

$$\begin{aligned} f'(x) &= \frac{(x^2 + 1)(6) - (6x - 5)(2x)}{(x^2 + 1)^2} \\ &= \frac{2(3 + 5x - 3x^2)}{(x^2 + 1)^2} \end{aligned}$$

47. $f(x) = (4 - 3x^2)^{-1}$

$$f'(x) = -(4 - 3x^2)^{-2}(-6x) = \frac{6x}{(4 - 3x^2)^2}$$

48. $f(x) = 9(3x^2 - 2x)^{-1}$

$$f'(x) = -9(3x^2 - 2x)^{-2}(6x - 2) = \frac{18(1 - 3x)}{(3x^2 - 2x)^2}$$

49. $y = \frac{x^2}{\cos x}$

$$y' = \frac{\cos x(2x) - x^2(-\sin x)}{\cos^2 x} = \frac{2x \cos x + x^2 \sin x}{\cos^2 x}$$

50. $y = \frac{\sin x}{x^2}$

$$y' = \frac{(x^2) \cos x - (\sin x)(2x)}{x^4} = \frac{x \cos x - 2 \sin x}{x^3}$$

51. $y = 3x^2 \sec x$

$$y' = 3x^2 \sec x \tan x + 6x \sec x$$

52. $y = 2x - x^2 \tan x$

$$y' = 2 - x^2 \sec^2 x - 2x \tan x$$

53. $y = x \cos x - \sin x$

$y' = -x \sin x + \cos x - \cos x = -x \sin x$

55. $f(x) = \frac{2x^3 - 1}{x^2} = 2x - x^{-2}$, $(1, 1)$

$f'(x) = 2 + 2x^{-3}$

$f'(1) = 4$

Tangent line: $y - 1 = 4(x - 1)$

$y = 4x - 3$

54. $g(x) = 3x \sin x + x^2 \cos x$

$$\begin{aligned}g'(x) &= 3x \cos x + 3 \sin x - x^2 \sin x + 2x \cos x \\&= 5x \cos x + (3 - x^2) \sin x\end{aligned}$$

56. $f(x) = \frac{x+1}{x-1}$, $\left(\frac{1}{2}, -3\right)$

$f'(x) = \frac{(x-1)-(x+1)}{(x-1)^2} = \frac{-2}{(x-1)^2}$

$f'\left(\frac{1}{2}\right) = \frac{-2}{(1/4)} = -8$

Tangent line: $y + 3 = -8\left(x - \frac{1}{2}\right)$

$y = -8x + 1$

57. $f(x) = -x \tan x$, $(0, 0)$

$f'(x) = -x \sec^2 x - \tan x$

$f'(0) = 0$

Tangent line: $y - 0 = 0(x - 0)$

$y = 0$

58. $f(x) = \frac{1 + \sin x}{1 - \sin x}$, $(\pi, 1)$

$f'(x) = \frac{(1 - \sin x)\cos x - (1 + \sin x)(-\cos x)}{(1 - \sin x)^2}$

$f'(\pi) = \frac{-1 - 1}{1} = -2$

Tangent line: $y - 1 = -2(x - \pi)$

$y = -2x + 2\pi + 1$

59. $v(t) = 36 - t^2$, $0 \leq t \leq 6$

$a(t) = v'(t) = -2t$

$v(4) = 36 - 16 = 20 \text{ m/sec}$

$a(4) = -8 \text{ m/sec}^2$

60. $v(t) = \frac{90t}{4t + 10}$

$a(t) = \frac{(4t + 10)90 - 90t(4)}{(4t + 10)^2}$

$= \frac{900}{(4t + 10)^2} = \frac{225}{(2t + 5)^2}$

(a) $v(1) = \frac{90}{14} \approx 6.43 \text{ ft/sec}$

(b) $v(5) = \frac{90(5)}{30} = 15 \text{ ft/sec}$

(c) $v(10) = \frac{90(10)}{50} = 18 \text{ ft/sec}$

$a(1) = \frac{225}{49} \approx 4.59 \text{ ft/sec}^2$

$a(5) = \frac{225}{15^2} = 1 \text{ ft/sec}^2$

$a(10) = \frac{225}{25^2} = 0.36 \text{ ft/sec}^2$

61. $g(t) = t^3 - 3t + 2$

$g'(t) = 3t^2 - 3$

$g''(t) = 6t$

62. $f(x) = 12x^{1/4}$

$f'(x) = 3x^{-3/4}$

$f''(x) = \frac{-9}{4}x^{-7/4} = \frac{-9}{4x^{7/4}}$

63. $f(\theta) = 3 \tan \theta$

$f'(\theta) = 3 \sec^2 \theta$

$$\begin{aligned}f''(\theta) &= 6 \sec \theta (\sec \theta \tan \theta) \\&= 6 \sec^2 \theta \tan \theta\end{aligned}$$

64. $h(t) = 4 \sin t - 5 \cos t$

$$h'(t) = 4 \cos t + 5 \sin t$$

$$h''(t) = -4 \sin t + 5 \cos t$$

66. $y = \frac{(10 - \cos x)}{x}$

$$xy + \cos x = 10$$

$$xy' + y - \sin x = 0$$

$$xy' + y = \sin x$$

65. $y = 2 \sin x + 3 \cos x$

$$y' = 2 \cos x - 3 \sin x$$

$$y'' = -2 \sin x - 3 \cos x$$

$$\begin{aligned} y'' + y &= -(2 \sin x + 3 \cos x) + (2 \sin x + 3 \cos x) \\ &= 0 \end{aligned}$$

67. $h(x) = \left(\frac{x-3}{x^2+1}\right)^2$

$$\begin{aligned} h'(x) &= 2\left(\frac{x-3}{x^2+1}\right)\left(\frac{(x^2+1)(1)-(x-3)(2x)}{(x^2+1)^2}\right) \\ &= \frac{2(x-3)(-x^2+6x+1)}{(x^2+1)^3} \end{aligned}$$

68. $f(x) = \left(x^2 + \frac{1}{x}\right)^5$

$$f'(x) = 5\left(x^2 + \frac{1}{x}\right)^4 \left(2x - \frac{1}{x^2}\right)$$

70. $h(\theta) = \frac{\theta}{(1-\theta)^3}$

$$h'(\theta) = \frac{(1-\theta)^3 - \theta[3(1-\theta)^2(-1)]}{(1-\theta)^6}$$

$$= \frac{(1-\theta)^2(1-\theta+3\theta)}{(1-\theta)^6} = \frac{2\theta+1}{(1-\theta)^4}$$

69. $f(s) = (s^2 - 1)^{5/2}(s^3 + 5)$

$$\begin{aligned} f'(s) &= (s^2 - 1)^{5/2}(3s^2) + (s^3 + 5)\left(\frac{5}{2}\right)(s^2 - 1)^{3/2}(2s) \\ &= s(s^2 - 1)^{3/2}[3s(s^2 - 1) + 5(s^3 + 5)] \\ &= s(s^2 - 1)^{3/2}(8s^3 - 3s + 25) \end{aligned}$$

72. $y = 1 - \cos 2x + 2 \cos^2 x$

$$y' = 2 \sin 2x - 4 \cos x \sin x$$

$$= 2[2 \sin x \cos x] - 4 \sin x \cos x$$

$$= 0$$

71. $y = 3 \cos(3x + 1)$

$$y' = -3 \sin(3x + 1)(3)$$

$$y' = -9 \sin(3x + 1)$$

73. $y = \frac{x}{2} - \frac{\sin 2x}{4}$

$$y' = \frac{1}{2} - \frac{1}{4} \cos 2x(2) = \frac{1}{2}(1 - \cos 2x) = \sin^2 x$$

74. $y = \frac{\sec^7 x}{7} - \frac{\sec^5 x}{5}$

$$y' = \sec^6 x (\sec x \tan x) - \sec^4 x (\sec x \tan x)$$

$$= \sec^5 x \tan x (\sec^2 x - 1)$$

$$= \sec^5 x \tan^3 x$$

75. $y = \frac{2}{3} \sin^{3/2} x - \frac{2}{7} \sin^{7/2} x$

$$y' = \sin^{1/2} x \cos x - \sin^{5/2} x \cos x$$

$$= (\cos x) \sqrt{\sin x} (1 - \sin^2 x)$$

$$= (\cos^3 x) \sqrt{\sin x}$$

76. $f(x) = \frac{3x}{\sqrt{x^2+1}}$

$$f'(x) = \frac{3(x^2+1)^{1/2} - 3x^2(x^2+1)^{-1/2}(2x)}{x^2+1}$$

$$= \frac{3(x^2+1) - 3x^2}{(x^2+1)^{3/2}} = \frac{3}{(x^2+1)^{3/2}}$$

77. $y = \frac{\sin \pi x}{x+2}$

$$y' = \frac{(x+2)\pi \cos \pi x - \sin \pi x}{(x+2)^2}$$

78. $y = \frac{\cos(x-1)}{x-1}$

$$\begin{aligned}y' &= \frac{-(x-1)\sin(x-1) - \cos(x-1)(1)}{(x-1)^2} \\&= -\frac{1}{(x-1)^2}[(x-1)\sin(x-1) + \cos(x-1)]\end{aligned}$$

80. $f(x) = \sqrt[3]{x^2 - 1}$

$$\begin{aligned}f'(x) &= \frac{1}{3}(x^2 - 1)^{-2/3}(2x) = \frac{2x}{3(x^2 - 1)^{2/3}} \\f'(3) &= \frac{2(3)}{3(4)} = \frac{1}{2}\end{aligned}$$


82. $y = \csc 3x + \cot 3x$

$$\begin{aligned}y' &= -3 \csc 3x \cot 3x - 3 \csc^2 3x \\y'\left(\frac{\pi}{6}\right) &= 0 - 3 = -3\end{aligned}$$

84. $f(x) = [(x-2)(x+4)]^2 = (x^2 + 2x - 8)^2$

$$\begin{aligned}f'(x) &= 4(x^3 + 3x^2 - 6x - 8) \\&= 4(x-2)(x+1)(x+4)\end{aligned}$$


The zeros of f' correspond to the points on the graph of f where the tangent line is horizontal.



86. $y = \sqrt{3x}(x+2)^3$

$$y' = \frac{3(x+2)^2(7x+2)}{2\sqrt{3x}}$$

y' does not equal zero for any x in the domain.
The graph has no horizontal tangent lines.



79. $f(x) = \sqrt{1-x^3}$

$$\begin{aligned}f'(x) &= \frac{1}{2}(1-x^3)^{-1/2}(-3x^2) = \frac{-3x^2}{2\sqrt{1-x^3}} \\f'(-2) &= \frac{-12}{2(3)} = -2\end{aligned}$$

81. $y = \frac{1}{2} \csc 2x$


$$y' = -\csc 2x \cot 2x$$

$$y'\left(\frac{\pi}{4}\right) = 0$$

83. $g(x) = 2x(x+1)^{-1/2}$

$$g'(x) = \frac{x+2}{(x+1)^{3/2}}$$


g' does not equal zero for any value of x in the domain.
The graph of g has no horizontal tangent lines.



85. $f(t) = \sqrt[3]{t+1} \sqrt[3]{t+1}$

$$\begin{aligned}f(t) &= (t+1)^{1/2}(t+1)^{1/3} = (t+1)^{5/6} \\f'(t) &= \frac{5}{6(t+1)^{1/6}}\end{aligned}$$

f' does not equal zero for any x in the domain.
The graph of f has no horizontal tangent lines.



87. $f(t) = t^2(t-1)^5, (2, 4)$


(a) $f'(t) = t(t-1)^4(7t-2)$

$$f'(2) = 24$$

(b) $y - 4 = 24(t-2)$

$$y = 24t - 44$$

(c)



88. $g(x) = x\sqrt{x^2 + 1}$, $(3, 3\sqrt{10})$


(a) $g'(x) = \sqrt{x^2 + 1} + \frac{x^2}{\sqrt{x^2 + 1}}$

$$g'(3) = \frac{19\sqrt{10}}{10}$$

(b) $y - 3\sqrt{10} = \frac{19\sqrt{10}}{10}(x - 3)$

$$y = \frac{19\sqrt{10}}{10}x - \frac{27\sqrt{10}}{10}$$

(c)



89. $y = \tan \sqrt{1-x}$, $(-2, \tan \sqrt{3})$


(a) $y' = \frac{-1}{2\sqrt{1-x} \cos^2 \sqrt{1-x}}$

$$y'(-2) = \frac{-\sqrt{3}}{6 \cos^2 \sqrt{3}} \approx -11.1983$$

(b) $y - \tan \sqrt{3} = \frac{-\sqrt{3}}{6 \cos^2 \sqrt{3}}(x + 2)$

$$y = \frac{-\sqrt{3}}{6 \cos^2 \sqrt{3}}x + \tan \sqrt{3} - \frac{\sqrt{3}}{3 \cos^2 \sqrt{3}}$$

(c)



90. $y = 2 \csc^3(\sqrt{x}) = \frac{2}{\sin^3(\sqrt{x})}$, $(1, 2 \csc^3(1))$


(a) $y' = \frac{-3 \cos(\sqrt{x})}{\sqrt{x} \sin^4(\sqrt{x})}$

$$y'(1) = \frac{-3 \cos(1)}{\sin^4(1)}$$

(b) $y - 2 \csc^3(1) = \frac{-3 \cos(1)}{\sin^4(1)}(x - 1)$

$$y = \frac{-3 \cos(1)}{\sin^4(1)}x + \frac{2}{\sin^3(1)} + \frac{3 \cos(1)}{\sin^4(1)}$$

(c)



91. $y = 2x^2 + \sin 2x$

$$y' = 4x + 2 \cos 2x$$

$$y'' = 4 - 4 \sin 2x$$

92. $y = x^{-1} + \tan x$

$$y' = -x^{-2} + \sec^2 x$$

$$y'' = 2x^{-3} + 2 \sec x (\sec x \tan x)$$

$$= \frac{2}{x^3} + 2 \sec^2 x \tan x$$

93. $f(x) = \cot x$

$$f'(x) = -\csc^2 x$$

$$f''(x) = -2 \csc x (-\csc x \cdot \cot x)$$

$$= 2 \csc^2 x \cot x$$

94. $y = \sin^2 x$

$$y' = 2 \sin x \cos x = \sin 2x$$

$$y'' = 2 \cos 2x$$

95. $f(t) = \frac{t}{(1-t)^2}$

$$f'(t) = \frac{t+1}{(1-t)^3}$$

$$f''(t) = \frac{2(t+2)}{(1-t)^4}$$

96. $g(x) = \frac{6x-5}{x^2+1}$

$$g'(x) = \frac{2(-3x^2 + 5x + 3)}{(x^2 + 1)^2}$$

$$g''(x) = \frac{2(6x^3 - 15x^2 - 18x + 5)}{(x^2 + 1)^3}$$

97. $g(\theta) = \tan 3\theta - \sin(\theta - 1)$

$$g'(\theta) = 3 \sec^2 3\theta - \cos(\theta - 1)$$

$$g''(\theta) = 18 \sec^2 3\theta \tan 3\theta + \sin(\theta - 1)$$

98. $h(x) = x\sqrt{x^2 - 1}$

$$h'(x) = \frac{2x^2 - 1}{\sqrt{x^2 - 1}}$$

$$h''(x) = \frac{x(2x^2 - 3)}{(x^2 - 1)^{3/2}}$$

99. $T = \frac{700}{t^2 + 4t + 10}$

$$T = 700(t^2 + 4t + 10)^{-1}$$

$$T' = \frac{-1400(t+2)}{(t^2 + 4t + 10)^2}$$

(a) When $t = 1$,

$$T' = \frac{-1400(1+2)}{(1+4+10)^2} \approx -18.667 \text{ deg/hr.}$$

(c) When $t = 5$,

$$T' = \frac{-1400(5+2)}{(25+20+10)^2} \approx -3.240 \text{ deg/hr.}$$

(b) When $t = 3$,

$$T' = \frac{-1400(3+2)}{(9+12+10)^2} \approx -7.284 \text{ deg/hr.}$$

(d) When $t = 10$,

$$T' = \frac{-1400(10+2)}{(100+40+10)^2} \approx -0.747 \text{ deg/hr.}$$

100. $v = \sqrt{2gh} = \sqrt{2(32)h} = 8\sqrt{h}$

$$\frac{dv}{dh} = \frac{4}{\sqrt{h}}$$

(a) When $h = 9$, $\frac{dv}{dh} = \frac{4}{3}$ ft/sec.

(b) When $h = 4$, $\frac{dv}{dh} = 2$ ft/sec.

101. $x^2 + 3xy + y^3 = 10$

$$2x + 3xy' + 3y + 3y^2y' = 0$$

$$3(x + y^2)y' = -(2x + 3y)$$

$$y' = \frac{-(2x + 3y)}{3(x + y^2)}$$

102. $x^2 + 9y^2 - 4x + 3y = 0$

$$2x + 18yy' - 4 + 3y' = 0$$

$$3(6y + 1)y' = 4 - 2x$$

$$y' = \frac{4 - 2x}{3(6y + 1)}$$

103. $y\sqrt{x} - x\sqrt{y} = 16$

$$y\left(\frac{1}{2}x^{-1/2}\right) + x^{1/2}y' - x\left(\frac{1}{2}y^{-1/2}y'\right) - y^{1/2} = 0$$

$$\left(\sqrt{x} - \frac{x}{2\sqrt{y}}\right)y' = \sqrt{y} - \frac{y}{2\sqrt{x}}$$

$$\frac{2\sqrt{xy} - x}{2\sqrt{y}}y' = \frac{2\sqrt{xy} - y}{2\sqrt{x}}$$

$$y' = \frac{2\sqrt{xy} - y}{2\sqrt{x}} \cdot \frac{2\sqrt{y}}{2\sqrt{xy} - x} = \frac{2y\sqrt{x} - y\sqrt{y}}{2x\sqrt{y} - x\sqrt{x}}$$

104. $y^2 = x^3 - x^2y + xy - y^2$

$$0 = x^3 - x^2y + xy - 2y^2$$

$$0 = 3x^2 - x^2y' - 2xy + xy' + y - 4yy'$$

$$(x^2 - x + 4y)y' = 3x^2 - 2xy + y$$

$$y' = \frac{3x^2 - 2xy + y}{x^2 - x + 4y}$$

105. $x \sin y = y \cos x$

$$(x \cos y)' + \sin y = -y \sin x + y' \cos x$$

$$y'(x \cos y - \cos x) = -y \sin x - \sin y$$

$$y' = \frac{y \sin x + \sin y}{\cos x - x \cos y}$$

106. $\cos(x + y) = x$

$-(1 + y') \sin(x + y) = 1$

$-y' \sin(x + y) = 1 + \sin(x + y)$

$y' = -\frac{1 + \sin(x + y)}{\sin(x + y)}$


$= -\csc(x + 1) - 1$

107. $x^2 + y^2 = 20$

$2x + 2yy' = 0$

$y' = -\frac{x}{y}$

At $(2, 4)$: $y' = -\frac{1}{2}$



Tangent line: $y - 4 = -\frac{1}{2}(x - 2)$

$x + 2y - 10 = 0$

Normal line: $y - 4 = 2(x - 2)$

$2x - y = 0$

108. $x^2 - y^2 = 16$

$2x - 2yy' = 0$

$y' = \frac{x}{y}$


At $(5, 3)$: $y' = \frac{5}{3}$

Tangent line: $y - 3 = \frac{5}{3}(x - 5)$

$5x - 3y - 16 = 0$

Normal line: $y - 3 = -\frac{3}{5}(x - 5)$

$3x + 5y - 30 = 0$



109. $y = \sqrt{x}$

$\frac{dy}{dt} = 2$ units/sec

$\frac{dy}{dt} = \frac{1}{2\sqrt{x}} \frac{dx}{dt} \Rightarrow \frac{dx}{dt} = 2\sqrt{x} \frac{dy}{dt} = 4\sqrt{x}$

(a) When $x = \frac{1}{2}$, $\frac{dx}{dt} = 2\sqrt{2}$ units/sec.

(b) When $x = 1$, $\frac{dx}{dt} = 4$ units/sec.

(c) When $x = 4$, $\frac{dx}{dt} = 8$ units/sec.

110. Surface area = $A = 6x^2$, x = length of edge


$\frac{dx}{dt} = 5$

$\frac{dA}{dt} = 12x \frac{dx}{dt} = 12(4.5)(5) = 270 \text{ cm}^2/\text{sec}$

111. $\frac{s}{h} = \frac{1/2}{2}$

$s = \frac{1}{4}h$

$\frac{dV}{dt} = 1$

Width of water at depth h :


$w = 2 + 2s = 2 + 2\left(\frac{1}{4}h\right) = \frac{4+h}{2}$

$V = \frac{5}{2}\left(2 + \frac{4+h}{2}\right)h = \frac{5}{4}(8+h)h$

$\frac{dV}{dt} = \frac{5}{2}(4+h)\frac{dh}{dt}$

$\frac{dh}{dt} = \frac{2(dV/dt)}{5(4+h)}$

When $h = 1$, $\frac{dh}{dt} = \frac{2}{25}$ m/min.



112. $\tan \theta = x$


$$\frac{d\theta}{dt} = 3(2\pi) \text{ rad/min}$$

$$\sec^2 \theta \left(\frac{d\theta}{dt} \right) = \frac{dx}{dt}$$

$$\frac{dx}{dt} = (\tan^2 \theta + 1)(6\pi) = 6\pi(x^2 + 1)$$

When $x = \frac{1}{2}$,

$$\frac{dx}{dt} = 6\pi \left(\frac{1}{4} + 1 \right) = \frac{15\pi}{2} \text{ km/min} = 450\pi \text{ km/hr.}$$



113. $s(t) = 60 - 4.9t^2$

$$s'(t) = -9.8t$$

$$s = 35 = 60 - 4.9t^2$$


$$4.9t^2 = 25$$

$$t = \frac{5}{\sqrt{4.9}}$$

$$\tan 30 = \frac{1}{\sqrt{3}} = \frac{s(t)}{x(t)}$$

$$x(t) = \sqrt{3}s(t)$$

$$\frac{dx}{dt} = \sqrt{3} \frac{ds}{dt} = \sqrt{3}(-9.8) \frac{5}{\sqrt{4.9}} \approx -38.34 \text{ m/sec}$$



Problem Solving for Chapter 2

1. (a) $x^2 + (y - r)^2 = r^2$, Circle

$$x^2 = y$$
, Parabola

Substituting:


$$(y - r)^2 = r^2 - y$$

$$y^2 - 2ry + r^2 = r^2 - y$$

$$y^2 - 2ry + y = 0$$

$$y(y - 2r + 1) = 0$$

Since you want only one solution, let $1 - 2r = 0 \Rightarrow r = \frac{1}{2}$. Graph $y = x^2$ and $x^2 + (y - \frac{1}{2})^2 = \frac{1}{4}$.



(b) Let (x, y) be a point of tangency:

$$x^2 + (y - b)^2 = 1 \Rightarrow 2x + 2(y - b)y' = 0 \Rightarrow y' = \frac{x}{b - y}$$
, Circle


$$y = x^2 \Rightarrow y' = 2x$$
, Parabola

Equating:

$$2x = \frac{x}{b - y}$$

$$2(b - y) = 1$$

$$b - y = \frac{1}{2} \Rightarrow b = y + \frac{1}{2}$$



Also, $x^2 + (y - b)^2 = 1$ and $y = x^2$ imply:

$$y + (y - b)^2 = 1 \Rightarrow y + \left[y - \left(y + \frac{1}{2} \right) \right]^2 = 1 \Rightarrow y + \frac{1}{4} = 1 \Rightarrow y = \frac{3}{4} \text{ and } b = \frac{5}{4}$$

$$\text{Center: } \left(0, \frac{5}{4} \right)$$

Graph $y = x^2$ and $x^2 + (y - \frac{5}{4})^2 = 1$.


2. Let (a, a^2) and $(b, -b^2 + 2b - 5)$ be the points of tangency. For $y = x^2$, $y' = 2x$ and for $y = -x^2 + 2x - 5$, $y' = -2x + 2$. Thus, $2a = -2b + 2 \Rightarrow a + b = 1$, or $a = 1 - b$. Furthermore, the slope of the common tangent line is

$$\begin{aligned} \frac{a^2 - (-b^2 + 2b - 5)}{a - b} &= \frac{(1-b)^2 + b^2 - 2b + 5}{(1-b) - b} = -2b + 2 \\ &\Rightarrow \frac{1 - 2b + b^2 + b^2 - 2b + 5}{1 - 2b} = -2b + 2 \\ &\Rightarrow 2b^2 - 4b + 6 = 4b^2 - 6b + 2 \\ &\Rightarrow 2b^2 - 2b - 4 = 0 \\ &\Rightarrow b^2 - b - 2 = 0 \\ &\Rightarrow (b-2)(b+1) = 0 \end{aligned}$$

$$b = 2, -1$$

For $b = 2$, $a = 1 - b = -1$ and the points of tangency are $(-1, 1)$ and $(2, -5)$.
The tangent line has slope -2 : $y - 1 = -2(x + 1) \Rightarrow y = -2x - 1$

For $b = -1$, $a = 1 - b = 2$ and the points of tangency are $(2, 4)$ and $(-1, -8)$.
The tangent line has slope 4 : $y - 4 = 4(x - 2) \Rightarrow y = 4x - 4$



3. (a)	$f(x) = \cos x$	$P_1(x) = a_0 + a_1x$	(b)	$f(x) = \cos x$	$P_2(x) = a_0 + a_1x + a_2x^2$
	$f(0) = 1$	$P_1(0) = a_0 \Rightarrow a_0 = 1$		$f(0) = 1$	$P_2(0) = a_0 \Rightarrow a_0 = 1$
	$f'(0) = 0$	$P'_1(0) = a_1 \Rightarrow a_1 = 0$		$f'(0) = 0$	$P'_2(0) = a_1 \Rightarrow a_1 = 0$
	$P_1(x) = 1$			$f''(0) = -1$	$P''_2(0) = 2a_2 \Rightarrow a_2 = -\frac{1}{2}$
				$P_2(x) = 1 - \frac{1}{2}x^2$	

(c)	x	-1.0	-0.1	-0.001	0	0.001	0.1	1.0
	$\cos x$	0.5403	0.9950	≈ 1	1	≈ 1	0.9950	0.5403
	$P_2(x)$	0.5	0.9950	≈ 1	1	≈ 1	0.9950	0.5

$P_2(x)$ is a good approximation of $f(x) = \cos x$ when x is near 0.

$$\begin{aligned} (d) \quad f(x) &= \sin x & P_3(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 \\ f(0) &= 0 & P_3(0) &= a_0 \Rightarrow a_0 = 0 \\ f'(0) &= 1 & P'_3(0) &= a_1 \Rightarrow a_1 = 1 \\ f''(0) &= 0 & P''_3(0) &= 2a_2 \Rightarrow a_2 = 0 \\ f'''(0) &= -1 & P'''_3(0) &= 6a_3 \Rightarrow a_3 = -\frac{1}{6} \\ P_3(x) &= x - \frac{1}{6}x^3 \end{aligned}$$

4. (a) $y = x^2$, $y' = 2x$, Slope = 4 at $(2, 4)$

$$\begin{aligned} \text{Tangent line: } y - 4 &= 4(x - 2) \\ y &= 4x - 4 \end{aligned}$$

- (c) Tangent line: $y = 0$

$$\text{Normal line: } x = 0$$

- (b) Slope of normal line: $-\frac{1}{4}$

$$\begin{aligned} \text{Normal line: } y - 4 &= -\frac{1}{4}(x - 2) \\ y &= -\frac{1}{4}x + \frac{9}{2} \\ y &= -\frac{1}{4}x + \frac{9}{2} = x^2 \\ &\Rightarrow 4x^2 + x - 18 = 0 \\ &\Rightarrow (4x + 9)(x - 2) = 0 \end{aligned}$$

—CONTINUED—

$$x = 2, -\frac{9}{4}$$

$$\text{Second intersection point: } \left(-\frac{9}{4}, \frac{81}{16}\right)$$

4. —CONTINUED—

- (d) Let (a, a^2) , $a \neq 0$, be a point on the parabola $y = x^2$. Tangent line at (a, a^2) is $y = 2a(x - a) + a^2$. Normal line at (a, a^2) is $y = -(1/2a)(x - a) + a^2$. To find points of intersection, solve:

$$x^2 = -\frac{1}{2a}(x - a) + a^2$$

$$x^2 + \frac{1}{2a}x = a^2 + \frac{1}{2}$$

$$x^2 + \frac{1}{2a}x + \frac{1}{16a^2} = a^2 + \frac{1}{2} + \frac{1}{16a^2}$$

$$\left(x + \frac{1}{4a}\right)^2 = \left(a + \frac{1}{4a}\right)^2$$

$$x + \frac{1}{4a} = \pm \left(a + \frac{1}{4a}\right)$$

$$x + \frac{1}{4a} = a + \frac{1}{4a} \Rightarrow x = a \quad (\text{Point of tangency})$$

$$x + \frac{1}{4a} = -\left(a + \frac{1}{4a}\right) \Rightarrow x = -a - \frac{1}{2a} = -\frac{2a^2 + 1}{2a}$$

The normal line intersects a second time at $x = -\frac{2a^2 + 1}{2a}$.

5. Let $p(x) = Ax^3 + Bx^2 + Cx + D$

$$p'(x) = 3Ax^2 + 2Bx + C.$$

$$\text{At } (1, 1): \quad A + B + C + D = 1 \quad \text{Equation 1}$$

$$3A + 2B + C = 14 \quad \text{Equation 2}$$

$$\text{At } (-1, -3): \quad -A + B - C + D = -3 \quad \text{Equation 3}$$

$$3A - 2B + C = -2 \quad \text{Equation 4}$$

Adding Equations 1 and 3: $2B + 2D = -2$

Subtracting Equations 1 and 3: $2A + 2C = 4$

Adding Equations 2 and 4: $6A + 2C = 12$

Subtracting Equations 2 and 4: $4B = 16$

Hence, $B = 4$ and $D = \frac{1}{2}(-2 - 2B) = -5$. Subtracting $2A + 2C = 4$ and $6A + 2C = 12$, you obtain $4A = 8 \Rightarrow A = 2$. Finally, $C = \frac{1}{2}(4 - 2A) = 0$. Thus, $p(x) = 2x^3 + 4x^2 - 5$.

6. $f(x) = a + b \cos cx$

$$f'(x) = -bc \sin cx$$

$$\text{At } (0, 1): \quad a + b = 1 \quad \text{Equation 1}$$

$$\text{At } \left(\frac{\pi}{4}, \frac{3}{2}\right): \quad a + b \cos\left(\frac{c\pi}{4}\right) = \frac{3}{2} \quad \text{Equation 2}$$

$$-bc \sin\left(\frac{c\pi}{4}\right) = 1 \quad \text{Equation 3}$$

From Equation 1, $a = 1 - b$. Equation 2 becomes

$$(1 - b) + b \cos\left(\frac{c\pi}{4}\right) = \frac{3}{2} \Rightarrow -b + b \cos\frac{c\pi}{4} = \frac{1}{2}.$$

From Equation 3, $b = \frac{-1}{c \sin(c\pi/4)}$. Thus:

$$\frac{1}{c \sin(c\pi/4)} + \frac{-1}{c \sin(c\pi/4)} \cos\left(\frac{c\pi}{4}\right) = \frac{1}{2}$$

$$1 - \cos\left(\frac{c\pi}{4}\right) = \frac{1}{2} c \sin\left(\frac{c\pi}{4}\right)$$

Graphing the equation

$$g(c) = \frac{1}{2} c \sin\left(\frac{c\pi}{4}\right) + \cos\left(\frac{c\pi}{4}\right) - 1,$$

you see that many values of c will work. One answer:

$$c = 2, b = -\frac{1}{2}, a = \frac{3}{2} \Rightarrow f(x) = \frac{3}{2} - \frac{1}{2} \cos 2x$$


7. (a) $x^4 = a^2x^2 - a^2y^2$

$$a^2y^2 = a^2x^2 - x^4$$

$$y = \frac{\pm\sqrt{a^2x^2 - x^4}}{a}$$

$$\text{Graph: } y_1 = \frac{\sqrt{a^2x^2 - x^4}}{a} \text{ and } y_2 = -\frac{\sqrt{a^2x^2 - x^4}}{a}$$

(b)



$(\pm a, 0)$ are the x -intercepts, along with $(0, 0)$.

(c) Differentiating implicitly:

$$4x^3 = 2a^2x - 2a^2yy'$$

$$y' = \frac{2a^2x - 4x^3}{2a^2y}$$

$$= \frac{x(a^2 - 2x^2)}{a^2y} = 0 \Rightarrow 2x^2 = a^2 \Rightarrow x = \frac{\pm a}{\sqrt{2}}$$

$$\left(\frac{a^2}{2}\right)^2 = a^2\left(\frac{a^2}{2}\right) - a^2y^2$$

$$\frac{a^4}{4} = \frac{a^4}{2} - a^2y^2$$


$$a^2y^2 = \frac{a^4}{4}$$

$$y^2 = \frac{a^2}{4}$$

$$y = \pm \frac{a}{2}$$

$$\text{Four points: } \left(\frac{a}{\sqrt{2}}, \frac{a}{2}\right), \left(\frac{a}{\sqrt{2}}, -\frac{a}{2}\right), \left(-\frac{a}{\sqrt{2}}, \frac{a}{2}\right), \left(-\frac{a}{\sqrt{2}}, -\frac{a}{2}\right)$$

9. (a)



Line determined by $(0, 30)$ and $(90, 6)$:

$$y - 30 = \frac{30 - 6}{0 - 90}(x - 0)$$

$$= -\frac{24}{90}x = -\frac{4}{15}x \Rightarrow y = -\frac{4}{15}x + 30$$

When $x = 100$:

$$y = -\frac{4}{15}(100) + 30 = \frac{10}{3} > 3 \Rightarrow \text{Shadow determined by man}$$

8. (a) $b^2y^2 = x^3(a - x); a, b > 0$

$$y^2 = \frac{x^3(a - x)}{b^2}$$

$$\text{Graph } y_1 = \frac{\sqrt{x^3(a - x)}}{b} \text{ and } y_2 = -\frac{\sqrt{x^3(a - x)}}{b}$$

(b) a determines the x -intercept on the right: $(a, 0)$.
 b affects the height.

(c) Differentiating implicitly:

$$2b^2yy' = 3x^2(a - x) - x^3 = 3ax^2 - 4x^3$$

$$y' = \frac{(3ax^2 - 4x^3)}{2b^2y} = 0$$

$$\Rightarrow 3ax^2 = 4x^3$$

$$3a = 4x$$


$$x = \frac{3a}{4}$$

$$b^2y^2 = \left(\frac{3a}{4}\right)^3 \left(a - \frac{3a}{4}\right) = \frac{27a^3}{64} \left(\frac{1}{4}a\right)$$

$$y^2 = \frac{27a^4}{256b^2} \Rightarrow y = \pm \frac{3\sqrt{3}a^2}{16b}$$

$$\text{Two points: } \left(\frac{3a}{4}, \frac{3\sqrt{3}a^2}{16b}\right), \left(\frac{3a}{4}, -\frac{3\sqrt{3}a^2}{16b}\right)$$

(b)



Line determined by $(0, 30)$ and $(60, 6)$:

$$y - 30 = \frac{30 - 6}{0 - 60}(x - 0) = -\frac{2}{5}x \Rightarrow y = -\frac{2}{5}x + 30$$

When $x = 70$:

$$y = -\frac{2}{5}(70) + 30$$

$= 2 < 3 \Rightarrow \text{Shadow determined by child}$

9. —CONTINUED—

(c) Need $(0, 30), (d, 6), (d + 10, 3)$ collinear.

$$\frac{30 - 6}{0 - d} = \frac{6 - 3}{d - (d + 10)} \Rightarrow \frac{24}{d} = \frac{3}{10} \Rightarrow d = 80 \text{ feet}$$

(d) Let y be the distance from the base of the street light to the tip of the shadow. We know that $dx/dt = -5$.

For $x > 80$, the shadow is determined by the man.

$$\frac{y}{30} = \frac{y - x}{6} \Rightarrow y = \frac{5}{4}x \text{ and } \frac{dy}{dt} = \frac{5}{4} \frac{dx}{dt} = \frac{-25}{4}$$

For $x < 80$, the shadow is determined by the child.

$$\frac{y}{30} = \frac{y - x - 10}{3} \Rightarrow y = \frac{10}{9}x + \frac{100}{9} \text{ and } \frac{dy}{dt} = \frac{10}{9} \frac{dx}{dt} = \frac{-50}{9}$$

Therefore:

$$\frac{dy}{dt} = \begin{cases} -\frac{25}{4}, & x > 80 \\ -\frac{50}{9}, & 0 < x < 80 \end{cases}$$


dy/dt is not continuous at $x = 80$.

10. (a) $y = x^{1/3} \Rightarrow \frac{dy}{dt} = \frac{1}{3}x^{-2/3} \frac{dx}{dt}$

$$1 = \frac{1}{3}(8)^{-2/3} \frac{dx}{dt}$$

$$\frac{dx}{dt} = 12 \text{ cm/sec}$$

(c) $\tan \theta = \frac{y}{x} \Rightarrow \sec^2 \theta \cdot \frac{d\theta}{dt} = \frac{x(dy/dt) - y(dx/dt)}{x^2}$



From the triangle, $\sec \theta = \sqrt{68}/8$. Hence

$$\frac{d\theta}{dt} = \frac{8(1) - 2(12)}{64(68/64)} = \frac{-16}{68} = \frac{-4}{17} \text{ rad/sec.}$$

11. $L'(x) = \lim_{\Delta x \rightarrow 0} \frac{L(x + \Delta x) - L(x)}{\Delta x}$
 $= \lim_{\Delta x \rightarrow 0} \frac{L(x) + L(\Delta x) - L(x)}{\Delta x}$
 $= \lim_{\Delta x \rightarrow 0} \frac{L(\Delta x)}{\Delta x}$

Also, $L'(0) = \lim_{\Delta x \rightarrow 0} \frac{L(\Delta x) - L(0)}{\Delta x}$.

But, $L(0) = 0$ because

$$L(0) = L(0 + 0) = L(0) + L(0) \Rightarrow L(0) = 0.$$

Thus, $L'(x) = L'(0)$ for all x . The graph of L is a line through the origin of slope $L'(0)$.

(b) $D = \sqrt{x^2 + y^2} \Rightarrow \frac{dD}{dt} = \frac{1}{2}(x^2 + y^2) \left(2x \frac{dx}{dt} + 2y \frac{dy}{dt} \right)$
 $= \frac{x(dx/dt) + y(dy/dt)}{\sqrt{x^2 + y^2}}$
 $= \frac{8(12) + 2(1)}{\sqrt{64 + 4}}$
 $= \frac{98}{\sqrt{68}} = \frac{49}{\sqrt{17}} \text{ cm/sec}$

12. $E'(x) = \lim_{\Delta x \rightarrow 0} \frac{E(x + \Delta x) - E(x)}{\Delta x}$

$$= \lim_{\Delta x \rightarrow 0} \frac{E(x)E(\Delta x) - E(x)}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} E(x) \left(\frac{E(\Delta x) - 1}{\Delta x} \right)$$

$$= E(x) \lim_{\Delta x \rightarrow 0} \frac{E(\Delta x) - 1}{\Delta x}$$

But, $E'(0) = \lim_{\Delta x \rightarrow 0} \frac{E(\Delta x) - E(0)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{E(\Delta x) - 1}{\Delta x} = 1$.

Thus, $E'(x) = E(x)E'(0) = E(x)$ exists for all x .

For example: $E(x) = e^x$.

13. (a)

z (degrees)	0.1	0.01	0.0001
$\frac{\sin z}{z}$	0.0174524	0.0174533	0.0174533

$$\begin{aligned}
 (c) \frac{d}{dz}(\sin z) &= \lim_{\Delta z \rightarrow 0} \frac{\sin(z + \Delta z) - \sin z}{\Delta z} \\
 &= \lim_{\Delta z \rightarrow 0} \frac{\sin z \cdot \cos \Delta z + \sin \Delta z \cdot \cos z - \sin z}{\Delta z} \\
 &= \lim_{\Delta z \rightarrow 0} \left[\sin z \left(\frac{\cos \Delta z - 1}{\Delta z} \right) \right] + \lim_{\Delta z \rightarrow 0} \left[\cos z \left(\frac{\sin \Delta z}{\Delta z} \right) \right] \\
 &= (\sin z)(0) + (\cos z)\left(\frac{\pi}{180}\right) = \frac{\pi}{180} \cos z
 \end{aligned}$$

14. (a) $v(t) = -\frac{27}{5}t + 27$ ft/sec

$$a(t) = -\frac{27}{5} \text{ ft/sec}^2$$

(b) $v(t) = -\frac{27}{5}t + 27 = 0 \Rightarrow \frac{27}{5}t = 27 \Rightarrow t = 5$ seconds

$$S(5) = -\frac{27}{10}(5)^2 + 27(5) + 6 = 73.5 \text{ feet}$$

(c) The acceleration due to gravity on Earth is greater in magnitude than that on the moon.

(b) $\lim_{z \rightarrow 0} \frac{\sin z}{z} \approx 0.0174533$

$$\text{In fact, } \lim_{z \rightarrow 0} \frac{\sin z}{z} = \frac{\pi}{180}.$$

(d) $S(90) = \sin\left(\frac{\pi}{180}90\right) = \sin\frac{\pi}{2} = 1$

$$C(180) = \cos\left(\frac{\pi}{180}180\right) = -1$$

$$\frac{d}{dz}S(z) = \frac{d}{dz} \sin(cz) = c \cdot \cos(cz) = \frac{\pi}{180}C(z)$$

(e) The formulas for the derivatives are more complicated in degrees.

15. $j(t) = a'(t)$

(a) $j(t)$ is the rate of change of acceleration.

(b) $s(t) = -8.25t^2 + 66t$

$$v(t) = -16.5t + 66$$

$$a(t) = -16.5$$

$$a'(t) = j(t) = 0$$

The acceleration is constant, so $j(t) = 0$.

(c) a is position.

b is acceleration.

c is jerk.

d is velocity.