


- Let R be the shaded region in the first quadrant enclosed by the y -axis and the graphs of $y = 1 - x^3$ and $y = \sin(x^2)$, as shown in the figure above.
 - Find the area of R .
 - A horizontal line, $y = k$, is drawn through the point where the graphs of $y = 1 - x^3$ and $y = \sin(x^2)$ intersect. Find k and determine whether this line divides R into two regions of equal area. Show the work that leads to your conclusion.
 - Find the volume of the solid generated when R is revolved about the line $y = -3$.

Free Response Section 1: Calculator Active



2. A planetary rover travels on a flat surface. The path of the rover for the time interval $0 \leq t \leq 2$ hours is shown in the rectangular coordinate system above. The rover starts at the point with coordinates $(6, 5)$ at time $t = 0$. The coordinates $(x(t), y(t))$ of the position of the rover change at rates given by

$$\begin{aligned}x'(t) &= -12 \sin(2t^2) \\y'(t) &= 10 \cos(1 + \sqrt{t}),\end{aligned}$$

where $x(t)$ and $y(t)$ are measured in meters and t is measured in hours.

- (a) Find the acceleration vector of the rover at time $t = 1$. Find the speed of the rover at time $t = 1$.
- (b) Find the total distance that the rover travels over the time interval $0 \leq t \leq 1$.
- (c) Find the y -coordinate of the position of the rover at time $t = 1$.
- (d) The rover receives a signal at each point where the line tangent to its path has slope $\frac{1}{2}$. At what times t , for $0 \leq t \leq 2$, does the rover receive a signal?

Free Response Section 2: Non Calculator

t (days)	0	10	22	30
$W'(t)$ (GL per day)	0.6	0.7	1.0	0.5

3. The twice-differentiable function W models the volume of water in a reservoir at time t , where $W(t)$ is measured in gigaliters (GL) and t is measured in days. The table above gives values of $W'(t)$ sampled at various times during the time interval $0 \leq t \leq 30$ days. At time $t = 30$, the reservoir contains 125 gigaliters of water.
- (a) Use the tangent line approximation to W at time $t = 30$ to predict the volume of water $W(t)$, in gigaliters, in the reservoir at time $t = 32$. Show the computations that lead to your answer.
- (b) Use a left Riemann sum, with the three subintervals indicated by the data in the table, to approximate $\int_0^{30} W'(t) dt$. Use this approximation to estimate the volume of water $W(t)$, in gigaliters, in the reservoir at time $t = 0$. Show the computations that lead to your answer.
- (c) Explain why there must be at least one time t , other than $t = 10$, such that $W'(t) = 0.7$ GL/day.
- (d) The equation $A = 0.3W^{2/3}$ gives the relationship between the area A , in square kilometers, of the surface of the reservoir, and the volume of water $W(t)$, in gigaliters, in the reservoir. Find the instantaneous rate of change of A , in square kilometers per day, with respect to t when $t = 30$ days.

Free Response Section 2: Non Calculator


5. Let f be the function satisfying $f'(x) = 4x - 2xf(x)$ for all real numbers x , with $f(0) = 5$ and $\lim_{x \rightarrow \infty} f(x) = 2$.

(a) Find the value of $\int_0^{\infty} (4x - 2xf(x)) dx$. Show the work that leads to your answer.

(b) Use Euler's method to approximate $f(-1)$, starting at $x = 0$, with two steps of equal size.

(c) Find the particular solution $y = f(x)$ to the differential equation $\frac{dy}{dx} = 4x - 2xy$ with the initial condition $f(0) = 5$.

Free Response Section 2: Non Calculator



5. The graph of the continuous function f , consisting of three line segments and a semicircle, is shown above.

Let g be the function given by $g(x) = \int_{-2}^x f(t) dt$.

(a) Find $g(-6)$ and $g(3)$.

(b) Find $g'(0)$.

(c) Find all values of x on the open interval $-6 < x < 3$ for which the graph of g has a horizontal tangent. Determine whether g has a local maximum, a local minimum, or neither at each of these values. Justify your answers.

(d) Find all values of x on the open interval $-6 < x < 3$ for which the graph of g has a point of inflection. Explain your reasoning.

Free Response Section 2: Non Calculator

6. The function f satisfies the equation

$$f'(x) = f(x) + x + 1$$

and $f(0) = 2$. The Taylor series for f about $x = 0$ converges to $f(x)$ for all x .

(a) Write an equation for the line tangent to the curve $y = f(x)$ at the point where $x = 0$.

(b) Find $f''(0)$ and find the second-degree Taylor polynomial for f about $x = 0$.

(c) Find the fourth-degree Taylor polynomial for f about $x = 0$.

(d) Find $f^{(n)}(0)$, the n th derivative of f at $x = 0$, for $n \geq 2$. Use the Taylor series for f about $x = 0$ and the Taylor series for e^x about $x = 0$ to find a polynomial expression for $f(x) - 4e^x$.